
# **PCT**





### INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6: C12N 15/12, 15/85, 5/10, C07K 14/705, 16/28, C12Q 1/68, A61K 31/70, 38/17, 48/00

(11) International Publication Number: **A2** 

WO 98/31799

(43) International Publication Date:

23 July 1998 (23.07.98)

(21) International Application Number:

PCT/US98/00959

(22) International Filing Date:

21 January 1998 (21.01.98)

(30) Priority Data:

60/034,204 21 January 1997 (21.01.97) US 60/034,205 21 January 1997 (21.01.97)

US

(71) Applicant (for all designated States except US): HUMAN GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US).

(72) Inventors; and

- (75) Inventors/Applicants (for US only): NI, Jian [CN/US]; 5502 Manorfield Road, Rockville, MD 20853 (US). GENTZ, Reiner, L. [DE/US]; 13404 Fairland Park Drive, Silver Spring, MD 20904 (US). ROSEN, Craig, A. [US/US]; 22400 Rolling Hill Road, Laytonsville, MD 20882 (US).
- (74) Agents: BROOKES, A., Anders et al.; Human Genome Sciences, Inc., 9410 Key West Avenue, Rockville, MD 20850 (US).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

#### Published

Without international search report and to be republished upon receipt of that report.

- (54) Title: POLYNUCLEOTIDES AND POLYPEPTIDES ENCODING RECEPTORS
- (57) Abstract

Receptor polypeptides and polynucleotides and methods for producing such polypeptides by recombinant techniques are disclosed. Also disclosed are methods for utilizing receptor polypeptides and polynucleotides in the design of protocols for the treatment of diseases and diagnostic assays for such conditions.

## FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AΤ	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	ТJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
ВJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

# POLYNUCLEOTIDES AND POLYPEPTIDES ENCODING RECEPTORS

## FIELD OF INVENTION

5

10

15

20

25

30

35

This invention relates to newly identified polynucleotides and the polypeptides encoded by them, the use of such polynucleotides and polypeptides, and their production. More particularly, the polynucleotides and polypeptides of the present invention relate to specific receptor families described in the specification and known in the art. The invention also relates to inhibiting or activating the action of such polynucleotides and polypeptides.

## BACKGROUND OF THE INVENTION

Receptor proteins are found on the membrane of the cells and are generally involved in signal transduction. There are many types of receptor proteins, and for convenience, these proteins are grouped in families based on similarity in structure and function.

For example, the TM4SF superfamily of cell surface proteins, also known as the tetraspan receptor superfamily, is comprised of at least seventeen individual gene products (these include CD9, CD20, CD37, CD53, CD63, CD81, CD82, A15, CO-029, Sm23, RDS, Uro B, Uro A, SAS, Rom-1, PETA3, and YKK8). The TM4SF superfamily is the second largest group in the CD antigen superfamily. Each member of the TM4SF superfamily can be characterized by several putative physical features including four highly conserved transmembrane domains, two divergent extracellular loops, and two short and highly divergent cytoplasmic tails. Expression patterns for members of the TM4SF superfamily tend to be rather broad and can vary widely between members. The functional roles of TM4SF superfamily members are primarily associated with signal transduction events and pathways, but also include cell adhesion in platelets and other lymphocytic and non-lymphocytic cell lines, as well as cell motility, proliferation, and metastasis. In addition, recent evidence suggests that a subset of the members of the TM4SF superfamily may function as potassium channel molecules.

One member of the TM4SF family, CD20, is a four membrane spanning domain cell surface phosphoprotein expressed exclusively on B lymphocytes. Although the precise functional role of CD20 has yet to be determined, it is thought to function primarily as a receptor during B-cell activation. Furthermore, a large number of experimental observations suggest several additional speculative roles for the CD20 molecule. For example, CD20-specific immunoprecipitation of biochemically cross-linked plasma membrane proteins suggests that CD20 assumes a multimeric structural

conformation characteristic of other previously described membrane channel proteins. Further experimentation has revealed that expression of exogenous CD20 on the cell surface specifically increases Ca<sup>2+</sup> conductance across the plasma membrane. Together, these results suggest that CD20 complexes may function as B-cell specific Ca<sup>2+</sup> ion channels. In addition, monoclonal antibodies raised against CD20 have been used to stimulate resting B-cells to transition out of the G0/G1 segment of the cell cycle. It has also been demonstrated that CD20 is associated with both serine and tyrosine kinases and, more specifically, that CD20 is associated, although not directly, with the Src family of tyrosine kinases including p56/53lyn, p56lck, and p59fyn.

A second example of a receptor subfamily, called sialoadhesin molecules, belongs to the Ig superfamily of receptor-like molecules. The more than 100 members of the Ig superfamily are generally considered to engage in specific cell-cell interactions through which intercellular communication may occur. In addition to classical protein-protein interactions, intercellular communication may also be mediated through protein-carbohydrate interactions. In fact, all members of the sialoadhesin family of the Ig superfamily are capable of mediating protein-sialic acid binding interactions. To date, only a small number of proteins have been assigned to the sialoadhesin family including sialoadhesin, CD33, CD22, the myelin-associated glycoprotein (MAG), and the Schwann cell myelin protein (SMP). Each of these proteins is expressed in a restricted subset of cell types. For example, CD22 and CD33 are expressed exclusively by B-lymphocytes and cells of the myelomonocytic lineage, respectively.

Similarly, galectins are a family of the lectin superfamily of carbohydrate-binding proteins which have a high affinity for b-galactoside sugars. Although a large number of glycoproteins containing b-galactoside sugars are produced by the cell, only a few will bind to known galectins *in vitro*. Such apparent binding specificity suggests a highly specific functional role for the galectins. Galectin 1 (conventionally termed LGALSI for lectin, galactoside-binding, soluble -1) is thought to specifically bind laminin, a highly polylactosaminated cellular glycoprotein, as well as the highly polylactosaminated lysosome-associated membrane proteins (LAMPs). Galectin 1 has also been shown to bind specifically to a lactosamine-containing glycolipid found on olfactory neurons and to integrin  $a_7b_1$  on skeletal muscle cells. Galectin 3 has also been observed to bind specifically to laminin, immunoglobulin E and its receptor, and bacterial lipopolysaccharides.

Various galectins have been shown to function in the mechanisms of intercellular communication. For example, depending on cell type, galectin 1 has been observed to modulate cell adhesion either positively or negatively. More specifically, galectin 1 appears to inhibit cell adhesion of skeletal muscle presumably by galectin 1 mediated disruption of laminin-integrin  $a_7b_1$  interactions. Alternatively, galectin 1 appears to promote cell adhesion in several non-skeletal muscle cell types examined

5

10

15

20

25

30

presumably by a glycoconjugate cross-linking mechanism. Galectin 3 has also been observed to function in modulating cell-adhesion, as well as in the activation of certain immune cells by cross-linking IgE and IgE receptors. In addition, galectins have been observed to be involved in the regulation of immune cell activity, as well as in such diverse processes as cell adhesion, proliferation, inflammation, autoimmunity, and metastasis of tumor cells. Furthermore, a galectin-like antigen designated HOM-HD-21 was recently found to be highly expressed in a Hodgkin's Disease cDNA library. Very recently, a novel galectin, termed PCTA-1, was identified as a specific cell surface marker on human prostate cancer cell lines and patient-derived carcinomas. Galectins have also been found to function intracellularly as a component of ribonucleoprotein complexes. Finally, galectins 1 and 3 have each been found to modulate T-cell growth and apoptosis by interaction with CD45 and possibly Bcl2, respectively.

A relatively new family of cell-surface proteins has been identified and termed the Ly6 superfamily. The members of this family include murine and human SCA-2, rat Ly-6 (also termed ThB), human CD59 [also known as protectin or membrane attack complex inhibition factor (MACIF)], and E48 antigen. The determination of an initial functional role for SCA-2 may lie in an analysis of its expression profile with regard to the complex process of hematopoiesis. SCA-2 is highly expressed in early thymic precusor cells. In turn, progeny of the intrathymic precusor population continue to express SCA-2, but only until the point of transition occurs from blast cell to small cell. Further experimental evidence demonstrates that mature thymocytes and peripheral T-cells do not express detectable levels of SCA-2, whereas mature, peripheral B-cells do continue to express SCA-2. As a result, it seems very likely that SCA-2 plays an important role in thymocyte maturation and differentiation. A plausible explanation for this functional hypothesis is that SCA-2 may act as a receptor for a unknown cytokine which regulates thymocyte maturation and differentiation.

In addition, CD59 is a recently identified integral membrane protein which appears to be involved in the regulation of complement. Recent studies show that the CD59 antigen may prevent damage from complement C5b-9 and protect astrocytes during inflammatory and infectious disorders of the nervous system. Expression of recombinant human CD59 on porcine donor organs have been shown to prevent complement-mediated lysis and activation of endothelial cells that leads to hyperacute rejection. Recently, researchers at Alexion Pharmaceuticals (New Haven, CT) reported on the production of transgenic pigs which expressed human CD59. In these animals, xenogeneic organs were resistant to hyperacute rejection. (Fodor, et al., "Expression of a functional human complement inhibitor in a transgenic pig as a model for the prevention of xenogeneic hyperacute organ rejection," Proc. Natl. Acad. Sci., 91:1153-11157 (1994).) The same company also reported that expression of recombinant transmembrane CD59 in paroxysmal nocturnal hemoglobinuria (PNH) B-

5

10

15

20

25

30

cells confers resistance to human complement. (Rother et al., "Expression of recombinant transmembrane CD59 in paroxysmal nocturnal hemoglobinuria B-cells confers resistance to human complement," Blood, 84:2604-2611 (1994).) PNH is an acquired hematopoietic disorder characterized by complement-mediated hemolytic anemia, pancytopenia, and venous thrombosis. It is thought that retroviral gene therapy with this molecule could provide a treatment for PNH patients.

A final Ly6 superfamily member, the E48 antigen, is involved in intercellular adhesion between keratinocyte cells of the squamous epithelium. Such keratinocytes are attached to adjoining cells by large numbers of desmosomes, which are thought to play a role in the transition of transformed keratinocytes to metastatic tumor cells. Treatment with a monoclonal antibody raised against the E48 antigen has been successful in the eradication of residual, postoperative squamous cell carcinoma cells of the upper aerodigestive tract in several *in vivo* models and, to some degree, in humans. (van Dongen, et al., "Progress in radioimmunotherapy of head and neck cancer," Oncol. Rep. 1:259-264 (1994).) The gene encoding the E48 antigen has been mapped to the q24-qter region of human chromosome 8. Interestingly, a number of human diseases have been mapped to this region of chromosome 8 including Langer-Giedion syndrome, brachio-otorhinolaryngeal syndrome, trichorhinolaryngeal syndrome, and epidermolysis bullosa simplex.

A further example of a receptor family includes the prohibitin receptors. The prohibitin gene product is expressed in a wide variety of tissues and has been implicated as a component of a number of anti-proliferative mechanisms. The prohibitin gene encodes a 30 kD postsynthetically modified polypeptide located primarily in the mitochondria, but also may be associated with the IgM receptor on the B-cell plasma membrane. The protein functionally inhibits DNA synthesis and entry into S phase of the cell cycle by an unknown mechanism. Interestingly, although the prohibitin gene product is hypothesized to be involved in the maintenance of senescence and the prevention of cancer, one study found that, although somatic mutations in the prohibitin gene were present in a small number of breast cancers, no mutations were identified in any other breast, ovary, liver, and lung cancers examined. (Sato et al., Genomics 17:762-764 (1993).) However, the prohibitin gene has been mapped to human chromosome 17q12-21, the same region thought to contain the gene involved in sporadic breast cancer. Furthermore, DNA sequence analysis of the prohibitin gene identified somatic mutation in 4 of 23 cases of sporadic breast cancer examined. Thus, prohibitin family members may be involved in the development of cancer.

Moreover, the EGFR family of plasma membrane proteins are an integral component of normal cellular proliferation and in the pathogenesis of the cancerous state. The family is relatively small and includes the EGFR, c-erbB-2, c-erbB-3, and others. Various cancers are correlated with aberrant expression of one or more of these

5

10

15

20

25

30

10

15

20

25

30

35

genes. A number of ligands have been identified which bind to the EGFR-like receptors listed above including TGF-a, heparin-binding EGF, amphiregulin, criptoregulin, heregulin, and others. A large fraction of adenocarcinomas examined to date, especially those of the breast, colon, and pancreas, are typified by the amplification or overexpression of the c-erbB-2 gene. EGF, or an analogous ligand, initiates the cellular growth factor response by binding to the EGFR, or EGFR-related, receptor. Following the binding event, the receptor molecule dimerizes activating its intracellular tyrosine kinase domain. This event results in the phosphorylation of specific tyrosine residues near the carboxy terminus of the receptor. The diversity of signals able to be transduced through the relatively small number of EGFR-related receptor molecules is amplified considerably by the recent finding that EGFR-like receptor molecules can function when dimerized with other EGFR family members forming heterodimers.

Members of the EGFR-related family of integral membrane proteins have been implicated in the pathogenesis of a number of human disease-states. For example, a mutation in the EGFR itself appears to play an important role in the development of glioblastomas. (Sang et al., J. Neurosurg 82:841-846 (1995).) The EGFR gene is amplified or overexpressed in the majority of primary human glioblastomas. Although not conferring a distinct advantage on cell growth, an increase in EGFR expression was found to confer an increase in the ability of glioma cells to maintain anchorage-independent growth in soft agar especially in response to EGF and retinoic acid. Anchorage-independent growth *in vitro* correlates highly with tumorigenicity *in vivo*, therefore, it is likely that cells which express abnormally high levels of EGFR in human glioblastoma cells may be involved in the high potential for these cells to cause tumors *in vivo*.

Moreover, overexpression or amplification of c-erbB-2 has been reported to be involved in a high number adenocarcinomas, particularly of the breast, colon, and pancreas, and in a small proportion of ovarian carcinomas.

Thus, there is a clear need for identifying and exploiting novel members of the receptor families, such as those described above. Although structurally related, these receptors will likely possess diverse and multifaceted functions in a variety of cell and tissue types. Receptor type molecules should prove useful in target based screens for small molecules and other such pharmacologically valuable factors. Monoclonal antibodies raised against such receptors may prove useful as therapeutics in an antitumor, diagnostic, or other capacity. Furthermore, receptors described here may prove useful in an active or passive immunotherapeutical role in patients with cancer or other immunocompromised disease states.

#### SUMMARY OF THE INVENTION

In one aspect, the invention relates to receptor polypeptides and polynucleotides, as well as the methods for their production. Another aspect of the invention relates to methods for using such receptor polypeptides and polynucleotides.

5 Such uses include the treatment of the specified diseases, among others. In still another aspect, the invention relates to methods to identify agonists and antagonists using the materials provided by the invention, and treating conditions associated with receptor imbalance with the identified compounds. Yet another aspect of the invention relates to diagnostic assays for detecting diseases associated with inappropriate receptor activity or levels.

# **DESCRIPTION OF THE INVENTION Definitions**

The following definitions are provided to facilitate understanding of certain terms used frequently herein.

"Receptor" refers, among others, to a polypeptide comprising the amino acid sequence set forth in SEQ ID NO:Y, or an allelic variant thereof.

"Receptor Activity" or "Biological Activity of the Receptor" refers to the metabolic or physiologic function of said receptor including similar activities or improved activities or these activities with decreased undesirable side-effects. Also included are antigenic and immunogenic activities of said receptor.

"Receptor gene" refers to a polynucleotide comprising the nucleotide sequence set forth in SEQ ID NO:X or allelic variants thereof and/or their complements.

"SEQ ID NO:X" comprises all or a substantial portion of the polynucleotide encoding each receptor of the invention. The value X for the nucleotide sequence is an integer specified in Table 1. This nucleotide sequence was translated into the receptor polypeptide identified in Table 1 as "SEQ ID NO:Y," where the value of Y for each receptor polypeptide is an integer defined in Table 1.

The invention further provides a composition of matter comprising a nucleic acid molecule which comprises a human cDNA clone identified by a cDNA Clone ID (Identifier) in Table 1, which DNA molecule is contained in the material deposited with the American Type Culture Collection ("ATCC") and given the ATCC Deposit Number shown in Table 1 for that cDNA clone. The ATCC is located at American Type Culture Collection (ATCC), 12301 Park Lawn Drive, Rockville, Maryland 20852, USA. The deposit has been made under the terms of the Budapest Treaty on the international recognition of the deposit of micro-organisms for purposes of patent procedure. The strain will be irrevocably and without restriction or condition released to the public upon the issuance of a patent. The deposit is provided merely as convenience to those of skill in the art and is not an admission that a deposit is required for enablement, such as that

15

20

25

30

required under 35 U.S.C. §112. The nucleotide sequence of the polynucleotides contained in the deposited material, as well as the amino acid sequence of the polypeptide encoded thereby, are controlling in the event of any conflict with any description of sequences herein.

"Antibodies" as used herein includes polyclonal and monoclonal antibodies, chimeric, single chain, and humanized antibodies, as well as Fab fragments, including the products of an Fab or other immunoglobulin expression library.

"Isolated" means altered "by the hand of man" from the natural state. If an "isolated" composition or substance occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living animal is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein.

"Polynucleotide" generally refers to any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotides" include, without limitation single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, "polynucleotide" refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term polynucleotide also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications has been made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. "Polynucleotide" also embraces relatively short polynucleotides, often referred to as oligonucleotides.

-

"Polypeptide" refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres. "Polypeptide" refers to both short chains, commonly referred to as peptides, oligopeptides or oligomers, and to longer chains, generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene-encoded amino acids. "Polypeptides" include amino acid sequences modified either by natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications can

5

10

15

20

25

30

occur anywhere in a polypeptide, including the peptide backbone, the amino acid sidechains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched as a result of ubiquitination, and they may be cyclic. with or without branching. Cyclic, branched and branched cyclic polypeptides may result from posttranslation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphotidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. (See, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York, 1993 and Wold, F., Posttranslational Protein Modifications: Perspectives and Prospects, pgs. 1-12 in POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic Press, New York, 1983; Seifter et al., "Analysis for protein modifications and nonprotein cofactors", Meth Enzymol (1990) 182:626-646 and Rattan et al., "Protein Synthesis: Posttranslational Modifications and Aging", Ann NY Acad Sci (1992) 663:48-62.)

"Variant" as the term is used herein, is a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide respectively, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to

5

10

15

20

25

30

10

15

20

25

30

35

sequence.

occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis.

"Identity" is a measure of the identity of nucleotide sequences or amino acid sequences. In general, the sequences are aligned so that the highest order match is obtained. "Identity" per se has an art-recognized meaning and can be calculated using published techniques. (See, e.g.: COMPUTATIONAL MOLECULAR BIOLOGY, Lesk, A.M., ed., Oxford University Press, New York, 1988; BIOCOMPUTING: INFORMATICS AND GENOME PROJECTS, Smith, D.W., ed., Academic Press, New York, 1993; COMPUTER ANALYSIS OF SEQUENCE DATA, PART I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; SEOUENCE ANALYSIS IN MOLECULAR BIOLOGY, von Heinje, G., Academic Press, 1987; and SEQUENCE ANALYSIS PRIMER, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991.) While there exist a number of methods to measure identity between two polynucleotide or polypeptide sequences, the term "identity" is well known to skilled artisans. (Carillo, H., and Lipton, D., SIAM J Applied Math (1988) 48:1073.) Methods commonly employed to determine identity or similarity between two sequences include, but are not limited to, those disclosed in Guide to Huge Computers, Martin J. Bishop, ed., Academic Press, San Diego, 1994, and Carillo, H., and Lipton, D., SIAM J Applied Math (1988) 48:1073. Methods to determine identity and similarity are codified in computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, GCS program package (Devereux, J., et al., Nucleic Acids Research (1984) 12(1):387), BLASTP, BLASTN, FASTA (Atschul. S.F. et al., a J Molec Biol (1990) 215:403.)

least, for example, 95% "identity" to a reference nucleotide sequence of SEQ ID NO:X is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence of SEQ ID NO: X. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference

As an illustration, by a polynucleotide having a nucleotide sequence having at

Similarly, by a polypeptide having an amino acid sequence having at least, for example, 95% "identity" to a reference amino acid sequence of SEQ ID NO:Y is intended that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid of SEQ ID NO:Y. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

15

20

10

5

## Polypeptides of the Invention

In one aspect, the present invention relates to receptor polypeptides (or receptor proteins). The receptor polypeptides include the polypeptide of SEQ ID NO:Y; as well as polypeptides comprising the amino acid sequence of SEQ ID NO:Y; and polypeptides comprising the amino acid sequence which have at least 80% identity to that of SEQ ID NO:Y over its entire length, and still more preferably at least 90% identity, and even still more preferably at least 95% identity to SEQ ID NO:Y. Furthermore, those with at least 97-99% identity to SEQ ID NO:Y are highly preferred. Also included within receptor polypeptides are polypeptides having the amino acid sequence which have at least 80% identity to the polypeptide having the amino acid sequence of SEQ ID NO:Y over its entire length, and still more preferably at least 90% identity, and even still more preferably at least 95% identity to SEQ ID NO:Y. Furthermore, those with at least 97-99% are highly preferred. Preferably receptor polypeptides exhibit at least one biological activity of the receptor.

30

25

The receptor polypeptides may be in the form of the "mature" protein or may be a part of a larger protein such as a fusion protein. It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, prosequences, sequences which aid in purification such as multiple histidine residues, or an additional sequence for stability during recombinant production.

35

Fragments of the receptor polypeptides are also included in the invention. A "fragment" is a polypeptide having an amino acid sequence that entirely is the same as part, but not all, of the amino acid sequence of the aforementioned receptor polypeptides. As with receptor polypeptides, fragments may be "free-standing," or comprised within a larger polypeptide of which they form a part or region, most

11

preferably as a single continuous region. Representative examples of polypeptide fragments of the invention, include, for example, fragments from about amino acid number 1-20, 21-40, 41-60, 61-80, 81-100, and 101 to the end of receptor polypeptide. In this context "about" includes the particularly recited ranges larger or smaller by several, 5, 4, 3, 2 or 1 amino acid at either extreme or at both extremes.

Preferred fragments include, for example, truncation polypeptides having the amino acid sequence of receptor polypeptides, except for deletion of a continuous series of residues that includes the amino terminus, or a continuous series of residues that includes the carboxyl terminus or deletion of two continuous series of residues. one including the amino terminus and one including the carboxyl terminus.

Also preferred are fragments characterized by structural or functional domains, such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions. The "domains" of each receptor polypeptide are illustrated in the Figures. The Figures compare SEQ ID NO:Y to the closest know homologue. Identical amino acids shared between the two polypeptides are shaded, while conservative amino acid changes are boxed. By examining the regions or amino acids shaded and/or boxed, the skilled artisan can readily identify conserved domains between the two polypeptides. The amino acids sequences of SEQ ID NO:Y falling within these conserved domains are "fragments" and are specifically contemplated by the present invention. Especially preferred is the extracellular domains of a receptor of the invention. Soluble extracellular domains have antagonist activity mediated by competition with a receptor ligand.

Other preferred fragments are biologically active fragments. Biologically active fragments are those that mediate receptor activity, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also included are those that are antigenic or immunogenic in an animal, especially in a human.

Preferably, all of these polypeptide fragments retain a biological activity of the receptor, including antigenic activity. Variants of the defined sequence and fragments also form part of the present invention. Preferred variants are those that vary from the referents by conservative amino acid substitutions -- i.e., those that substitute a residue with another of like characteristics. Typical such substitutions are among Ala. Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and Glu; among Asn and Gln; and among the basic residues Lys and Arg; or aromatic residues Phe and Tyr. Particularly preferred are variants in which several, 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination.

5

10

15

20

25

30

The receptor polypeptides of the invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

## Polynucleotides of the Invention

5

10

15

20

25

30

35

Another aspect of the invention relates to receptor polynucleotides. Receptor polynucleotides include isolated polynucleotides which encode the receptor polypeptides and fragments, and polynucleotides closely related thereto. More specifically, a receptor polynucleotide of the invention includes a polynucleotide comprising the nucleotide sequence contained in SEQ ID NO:X encoding a receptor polypeptide of SEQ ID NO:Y, and polynucleotide having the particular sequence of SEQ ID NO:X.

Receptor polynucleotides further include a polynucleotide comprising a nucleotide sequence that has at least 80% identity over its entire length to a nucleotide sequence encoding the receptor polypeptide of SEQ ID NO:Y, and a polynucleotide comprising a nucleotide sequence that is at least 80% identical to that of SEQ ID NO:X over its entire length. In this regard, polynucleotides at least 90% identical are particularly preferred, and those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred and those with at least 98-99% are most highly preferred, with at least 99% being the most preferred. Also included under receptor polynucleotides are a nucleotide sequence which has sufficient identity to a nucleotide sequence contained in SEQ ID NO:X, or contained in the cDNA insert in the plasmid deposited with ATCC, to hybridize under conditions useable for amplification or for use as a probe or marker. Moreover, the receptor polynucleotide includes a nucleotide sequence having at least 80% identity to a nucleotide sequence encoding the receptor polypeptide expressed by the cDNA insert deposited at the ATCC, and a nucleotide sequence comprising at least 15 contiguous nucleotides of such cDNA insert. In this regard, polynucleotides at least 90% identical are particularly preferred, and those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred and those with at least 98-99% are most highly preferred, with at least 99% being the most preferred. The invention also provides polynucleotides which are complementary to all the above receptor polynucleotides.

The receptors of the invention are structurally related to other proteins of specified receptor families, as shown by the results in the Figures. The cDNA sequence of SEQ ID NO:X encodes a polypeptide as described in Table 1 as SEQ ID NO:Y. Because the receptor polypeptides contain domains similar in structure to other receptor family members, the receptors of the present invention are expected to have,

inter alia, similar biological functions/properties to their homologous polypeptides and polynucleotides, and their utility is obvious to anyone skilled in the art.

Table 1

5

Clone ID Name	SEQ ID NO:X	SEQ ID NO:Y	ATCC Deposit No.	ATCC Deposit Date	Receptor Family	Homology
HMACR70			209054	05/16/97	Ig	Sialoadhesin
	l	18	#####	01/21/98	_	OB-1
HTEDK48			209054	05/16/97	TM4SF	MRC-OX44 PETA-3
1-1849 bp	2					
160-900 bp	3	19				
HTPED39			209054	05/16/97	TM4SF	NAG-2
HPWAE25	4	20	#####	1/21/98		TALLA-1
HTPEF86	5	21	209053	05/16/97	TM4SF	CD20
	,					B1 Antigen
HSBBF02	6	22	209054	05/16/97	TM4SF	TALLA-I
HLTAH80	7	23	97242	08/02/95	TM4SF	TALLA-1
			209054	05/16/97		
HTPBA27	8	. 24	97242	08/02/95	TM4SF	NAG-2
			209054	05/16/97		·
HAIDQ59			209054	05/16/97	TM4SF	CD9
						Antigen
5' Sequence	9	25				
3' Sequence	10					
HHFEK40	11	26	209054	05/16/97	TM4SF	PETA-3
HGBGV89	12	27	209125	06/09/97	TM4SF	L6H
			209054	05/16/97		₩.
HUVBB80	13	28	209054	05/16/97	TM4SF	L6
HJACE54	14	29	209053	05/16/97	Lectin	Galectin-3
		1	[			Galectin-5
	<u> </u>		<u> </u>			Galectin-8
HROAD63	15	30	209053	05/16/97	Ly6	E48 splice
				ļ		variant
HMWGS46	16	31	209053	05/16/97	Prohibitin	BAP-37
HNFGW06	17	32	209053	05/16/97	EGFR	EGFR

The novel full-length cDNA clone designated **HMACR70** may be a member of the sialoadhesin family of the Ig superfamily of receptor-like molecules and a CD33 homologue. HMACR70 contains a 1497 nucleotide cDNA insert encoding a 315 amino acid ORF and was cloned from a GM-CSF-treated human macrophage cDNA library. The only additional cDNA libraries in the HGS database which include this clone are human eosinophils and possibly human gall bladder. A BLAST analysis of the amino acid sequence of HMACR70 demonstrates that this clone exhibits approximately 50% identity and 69% similarity over a 300 amino acids stretch of a gene termed human

10

differentiation antigen, and 38% identity and 62% similarity of the human myelin-associated glycoprotein precursor CD33 gene.

A more recent BLAST analysis confirms HMACR70's designation as a sialoadhesin family member. HMACR70 is homologous to two recently identified sialoadhesin family members, human OB binding protein (OB) 1 and 2. (See, Genbank Accession No. U71382; see Figure 1.) It is thought that OB-1 and OB-2 may bind leptin. Thus, HMACR70, as a sialoadhesin family member, may act to attenuate or even amplify intercellular routes of communication, including binding to leptin or modulating the activity of immune cells, such as macrophages. Clearly, any diseases affected by these processes could be treated by the polypeptide or fragment of HMACR70.

The full-length nucleotide sequences of ten novel human cDNA clones which potentially belong to the TM4SF superfamily are disclosed in the table above and will be addressed sequentially.

The cDNA clone HTEDK48 contains a 1849 nucleotide cDNA insert encoding a 245 amino acid ORF that was cloned from a human testes cDNA library. The coding sequence of HTEDK48 (SEQ ID NO: 3) may be fused to other human proteins, such as 3-hydroxyacyl-CoA dehydrogenase. BLAST analysis of the amino acid sequence of HTEDK48 demonstrates that this clone exhibits approximately 30% identity and 51% similarity over a 245 amino acid stretch of the CD82 molecule. Recent studies have shown that CD82 can associate with CD4 or CD8 and deliver costimulatory signals for the TCR/CD3 pathway. CD82 has also been found to be involved in syncytium formation in HTLV-I-infected T-cells. And finally, in a recently published study in which the expression of the CD82 gene by tumors of the lung was examined retrospectively, it was reported that CD82 may be linked to the suppression of tumor metastasis of prostate cancer. The study also reported that decreased CD82 expression may be involved in malignant progression of such cancers. Thus, HTEDK48 may also be involved in the development of cancer.

A more recent BLAST analysis shows that HTEDK48 is homologous the rat leukocyte antigen, MRC OX-44, and the platelet endothelial tetraspan antigen -3 (PETA-3). (See Figure 2X.) MRC OX-44, a member of a new family of cell surface proteins, appears to be involved in growth regulation. (See, Bellacosa, A., et al., "The Rat Leukocyte antigen MRC OX-44 is a Member of a New Family of Cell Surface Proteins which Appear to be Involved in Growth Regulation," Mol. Cell. Bio. 11: 2864-2872 (1991).) Similarly, PETA-3 has been located to platelet endothelial cells, and an anti-PETA-3 antigen monoclonal antibody can stimulate platelet aggregation and mediator release. (See, Fitter, S., "Molecular Cloning of cDNA Encoding a Novel Platelet-Endothelial Cell Tetra-Span Antigen, PETA-3," Blood, 86(4):1348-1355 (1995).) Thus, HTEDK48 may function similar to MRC OX-44 or PETA-3 to affect

10

15

20

25

30

growth of blood cells. Administering polypeptides or fragments of HTEDK48 may be an effective treatment of blood disorders.

The cDNA clone **HPWAE25** contains a 1288 nucleotide cDNA insert encoding a 273 amino acid ORF that was cloned from a human pancreas tumor cDNA library, while clone **HTPED39** represents a truncated cDNA sequence. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including keratinocytes, ulcerative colitis, striatum depression, lymph node breast cancer, ovarian cancer, stage B2 prostate cancer, kidney medulla, and others. Northern blot analysis of HLTAH80 also shows expression in a variety of human cell lines including U937, MM96, WM115, and MDAMB231. A BLAST analysis of the amino acid sequence of HTPED39 demonstrates that this clone exhibits approximately 35% identity and 50% similarity over the entire length of the CD37 molecule. The CD37 antigen is expressed on B cells and on a subpopulation of T cells, but not on pre-B or plasma cells. It has been reported that CD37 expression is downregulated in conjunction with B-cell activation, suggesting that CD37 may be involved in the processes which dictate the activation state of the B-cell.

Moreover, HPWAE25 is also homologous to recently identified TM4SF members, NAG-2 and TALLA-1. (See Figure 3.) NAG-2 is thought to complex with integrins and other TM4SF proteins, while TALLA-1 is a highly specific marker of T-cell acute lymphoblastic leukemia and neuroblastoma. (See, Tachibana, I., et al., "NAG-2, A Novel Transmembrane-4 Superfamily (TM4SF) Protein that Complexs with Integrins and Other TM4SF Proteins," J. Biol. Chem., 272:29181-29189 (1997); Takagi, S., "Identification of a Higly Specific Surface Marker of T-cell Acute Lymphoblastic Leukemia and Neuroblastoma as a New Member of the Transmembrane 4 Superfamily," Int. J. Cancer 61(5):706-715 (1995).) Thus, HPWAE25 may be involved the development of cancer, particularly leukemia, lymphoma, and neuroblastoma. HPWAE25 may be used as an effective treatment of these cancers, as well as a diagnostic marker.

A subfamily of TM4SF receptors include CD20 proteins. A CD20-like cDNA clone was obtained from a human pancreas tumor cDNA library and contains a 1236 nucleotide insert which encodes a 250 amino acid ORF. A BLAST analysis of the deduced amino acid sequence of HTPEF86 exhibits approximately 41% identity and 61% similarity to the CD20 gene, also known as B1 antigen. (See Figure 4.) Expression of this gene is detected in only two additional HGS human cDNA libraries; amygdala depression and 9 week early stage human. Although the precise functional role of CD20 has yet to be determined, it is clear that CD20 plays a key role in the regulation of B-cell activation. Based primarily on sequence identity, the novel CD20-like molecule presented herein may also be involved in cell cycle activation. Potential therapeutic and/or diagnostic applications for HTPEF86 may include such clinical

5

10

15

20

25

30

presentations as juvenile rheumatoid arthritis, Graves' Disease, and a number of B-cell lymphomas or other lymphoid tumors.

The clone **HSBBF02** contains a 1115 nucleotide cDNA insert encoding a 245 amino acid ORF and was cloned from an HSC 172 cell line cDNA library. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including brain amygdala depression, endothelial cells, fetal liver and heart, osteoblasts, testes, and others. A BLAST analysis of the amino acid sequence of HSBBF02 demonstrates that this clone exhibits approximately 64% identity and 80% similarity with the A15 molecule over a 131 amino acid stretch (A15 is composed of 244 amino acids). A more recent BLAST search shows that HSBBF02 is similar to the TALLA-1 protein and may in fact be a closely related family member. (See Figure 5.)

In addition, a second cDNA clone, designated **HLTAH80**, exhibits sequence similarity to the A15 molecule and TALLA-1. (See Figure 6.) This clone contains a 1662 nucleotide cDNA insert encoding a 253 amino acid ORF and was cloned from a human T-cell lymphoma cDNA library. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including B-cell lymphoma, corpus collosum, endometrial tumor, osteosarcoma, testes, and others. Northern blot analysis of HLTAH80 also shows expression in a variety of human tissues including spleen, lymph node, thymus, PBLs, heart, and a particularly strong signal in skeletal muscle and pancreas. A BLAST analysis of the amino acid sequence of HLTAH80 demonstrates that this clone exhibits approximately 35% identity and 55% similarity over the entire length of the A15 molecule.

Since expression of A15 drops to undetectable levels when comparing immature T-cells to peripheral blood lymphocytes, it is thought that A15 may play a role in the development of T-cells. Furthermore, the MXS1(CCG-B7) gene which codes for A15 contains a number of triplet nucleotide repeats which have been associated with neuropsychiatric diseases such as Huntington's chorea, fragile X syndrome, and myotonic dystrophy. In addition, A15 appears to be expressed exclusively on T-cell acute lymphoblastic leukemia cell lines, including several derived from adult T-cell leukemia and those established by immortalization with human T-cell leukemia virus type I or Herpesvirus saimiri. Thus, clones HLTAH80 and/or HSBBF02 may also be involved in diseases caused by the expansion of repeats or chromosomal instability.

The cDNA clone **HTPBA27** contains a 1345 nucleotide cDNA insert encoding a 238 amino acid ORF and was cloned from a human tumor pancreas cDNA library. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including cerebellum, breast lymph node, osteosarcoma, adult testes, RS4;11 bone marrow cell line, microvascular endothelial cells, and others. A BLAST analysis of the amino acid sequence of HTPBA27 demonstrates that this

5

10

15

20

25

30

clone exhibits approximately 40% identity and 64% similarity with a glycoprotein termed CD53 over its entire length. CD53 is thought to be involved in thymopoiesis, since rat CD53 can be detected on immature CD4-8-thymocytes and the functionally mature single-positive subset, but cannot be detected on the intermediate CD4+8+ thymocytic subset of cells. The CD53 molecule has also been implicated as a component of signal transduction pathways in B cells, monocytes and granulocytes, rat macrophages; NK, and T cells. Moreover, as illustrated in Figure 7, HTPBA27 was recently confirmed as a TM4SF receptor. (See, Tachibana, I., et al., "NAG-2, A Novel Transmembrane-4 Superfamily (TM4SF) Protein that with Integrins and Other TM4SF Proteins," J. Biol. Chem., 272:29181-29189 (1997).) Calling the HTPBA27 polypeptide NAG-2, this group confirmed HTPBA27's status as a TM4SF receptor by showing that NAG-2 complexes with integrin and other TM4SF receptors. Thus, diseases caused by the failure of HTPBA27 to complex with integrin and other TM4SF receptors can be treated by administering HTPBA27. HTPBA27 can also be used to diagnose these diseases.

The cDNA clone **HAIDQ59** contains cDNA insert encoding a 221 amino acid ORF that was cloned from a human epithelial cell induced with TNFa and INF cDNA library. The 5' end of HAIDQ59 is represented by the SEQ ID NO: 9, while the 3' end is represented by SEQ ID NO: 10. This clone appears in only two additional cDNA libraries in the HGS database. These two libraries were constructed from the human Jurkat T-cell line and human microvascular endothelial cells. A BLAST analysis of the amino acid sequence of HAIDQ59 demonstrates that this clone exhibits approximately 53% identity and 69% similarity over 226 amino acids of the CD9 TM4SF molecule. (See Figure 8.) It has been demonstrated that the CD9 molecule is involved in signal transduction pathways in platelets, as well as in cell adhesion in both platelets and pre-B-cell lines. Intriguingly, a monoclonal antibody (vpg15), which recognizes the feline homologue of CD9, has been shown to block infection by feline immunodeficiency virus (FIV). Furthermore, a recent study shows that cells expressing high levels of CD9 exhibited suppressed cell motility. Thus, HAIDQ59 may also be involved in signal transduction of blood cells.

The cDNA clone **HHFEK40** contains a 936 nucleotide cDNA insert encoding a 252 amino acid ORF and was cloned from a human fetal heart cDNA library. This clone appears once in the human fetal heart cDNA library and possibly in a hemangiopericytoma cDNA library. A BLAST analysis of the amino acid sequence of HHFEK40 demonstrated that this clone exhibits approximately 60% identity and 75% similarity over the entire length of a molecule designated PETA-3. (See Figure 9.) PETA-3 was originally identified as a novel human platelet surface glycoprotein termed gp27. Although PETA-3 is present in low abundance on the platelet surface, an anti-PETA-3 monoclonal antibody can stimulate platelet aggregation and mediator release.

5

10

15

20

25

30

Thus, HHFEK40 may function similar to PETA-3 to affect growth of blood cells. Administering polypeptides or fragments of HHFEK40 may be an effective treatment of blood disorders.

The cDNA clone HGBGV89 contains a 738 nucleotide cDNA insert encoding a 197 amino acid ORF and was cloned from a human gall bladder cDNA library. The only two additional appearances of this clone in the HGS database are in a normalized fetal liver cDNA library and in a fetal liver/spleen cDNA library. The cDNA clone HUVBB80 contains a 1071 nucleotide cDNA insert encoding a 201 amino acid ORF and was cloned from a human umbilical vein cDNA library. This clone appears in several additional cDNA libraries in the HGS database including prostate BPH, thyroid, and fetal liver/spleen. BLAST analyses of the amino acid sequences of HGBGV89 and HUVBB80 demonstrate that these clones exhibit approximately 49% identity and 65% similarity and 47% identity and 68% similarity, respectively, over the entire length of a molecule designated L6 surface protein or human tumor-associated antigen L6. (See Figures 10 & 11.) Moreover, another group has confirmed the TM4SF receptor homology of HGBGV89 by describing the protein as a putative transmembrane protein L6H. (See Genbank Accession No 2587054; see Figure 10.) The L6 cell surface antigen is highly expressed on lung, breast, colon, and ovarian carcinomas. Promising results of phase 1 clinical studies have been reported with an anti-L6 monoclonal antibody, or its humanized counterpart, suggesting that the L6 antigen may be an attractive target for monoclonal antibody-based cancer therapy.

In summary, there is a clear need for identifying and exploiting novel members of the TM4SF superfamily such as those described herein. Although structurally related, these factors will likely possess diverse and multifaceted functions in a variety of cell and tissue types. Receptor type molecules, such as the novel potential members of the TM4SF superfamily detailed here, should prove useful in target based screens for small molecules and other such pharmacologically valuable factors. Monoclonal antibodies raised against such factors may prove useful as therapeutics in an anti-tumor, diagnostic, or other capacity. Furthermore, factors such as the nine novel TM4SF superfamily-like molecules described here may prove useful in an active or passive immunotherapeutical role in patients with cancer or other immunocompromised disease states.

Besides TM4SF receptors, receptors from other families are also described. For example, clone **HJACE54**, also called galectin 11, exhibits significant sequence identity to the rat galectin 5, the chicken galectin 3 gene, and the human galectin 8 genes. (See Figure 12.) The galectin 11 cDNA clone contains an 865 nucleotide insert which encodes a 133 amino acid ORF. The clone was obtained from a Jurkat T-cell G1 phase cDNA library. A BLAST analysis of the deduced amino acid sequence of HJACE54 demonstrates approximately 35% identity and 57% similarity to the amino

5

10

15

20

25

30

acid sequence of the rat galectin 5 gene. Expression of galectin 11 is quite limited in the HGS database. In fact, the only two additional ESTs in the HGS database which contain the HJACE54 sequence were found in human neutrophil and human infant adrenal gland cDNA libraries. Northern blot analyses have not been performed to examine expression patterns of the galectin 11 gene.

Various galectins have been shown to function in the mechanisms of intercellular communication. For example, depending on cell type, galectin 1 has been observed to modulate cell adhesion either positively or negatively. More specifically, galectin I appears to inhibit cell adhesion of skeletal muscle presumably by galectin Imediated disruption of laminin-integrin a<sub>2</sub>b<sub>1</sub> interactions. Alternatively, galectin 1 appears to promote cell adhesion in several non-skeletal muscle cell types examined presumably by a glycoconjugate cross-linking mechanism. Galectin 3 has also been observed to function in modulating cell-adhesion, as well as in the activation of certain immune cells by cross-linking IgE and IgE receptors. In addition, galectins have been observed to be involved in the regulation of immune cell activity, as well as in such diverse processes as cell adhesion, proliferation, inflammation, autoimmunity, and metastasis of tumor cells. Furthermore, a galectin-like antigen designated HOM-HD-21 was recently found to be highly expressed in a Hodgkin's Disease cDNA library. Very recently, a novel galectin, termed PCTA-1, was identified as a specific cell surface marker on human prostate cancer cell lines and patient-derived carcinomas. Galectins have also been found to function intracellularly as a component of ribonucleoprotein complexes. Finally, galectins 1 and 3 have each been found to modulate T-cell growth and apoptosis by interaction with CD45 and possibly Bcl2, respectively. As a result, the discovery of a novel galectin, such as that encoded by HJACE54, is likely to be a valuable asset both diagnostically and therapeutically.

Additionally, a full-length nucleotide sequence of a novel human cDNA clone which encodes an apparent splice variant of the previously described human E48 antigen has recently been determined. (See Figure 13.) Clone HROAD63 contains a 441 nucleotide cDNA which encodes a 70 amino acid polypeptide. This novel clone exhibits significant sequence identity to several members of a relatively new family of cell-surface proteins termed the Ly6 superfamily. These members include murine and human SCA-2, rat Ly-6 (also termed ThB), and human CD59 [also known as protectin or membrane attack complex inhibition factor (MACIF)]. The novel E48 splice variant was obtained from the HGS human stomach cDNA library. The clone is present in only a limited number of other HGS cDNA libraries including kidney cancer, keratinocyte, and tongue. An alignment of the nucleotide sequences of the human E48 and HROAD63 cDNAs demonstrates that the initial 168 and 178 nucleotides of E48 and HROAD63, respectively, are identical, with the exception of an additional 10 nucleotides of sequence at the extreme 5' end of the HROAD63 sequence. The

10

15

20

25

30

sequence of the two clones is also identical for an additional 229 nucleotides including the 3' end of the coding sequences and the entire 3' untranslated regions. The only divergence of nucleotide sequence in this region of the clones is the deletion of a single thymidine residue in the 3' UTR of the E48 cDNA. The major difference between the two nucleotide sequences is a 329 nucleotide deletion from the HROAD63 sequence. This deletion causes a shift in the HROAD63 reading frame and encompasses the translational stop signal used in the E48 clone. As a result, the carboxy terminal sequence of HROAD63 is radically altered with regard to that of E48 (as illustrated in Figure 13 by the obvious differences between amino acids 56-128 of E48 and 56-70 of HROAD63 in the amino acid alignment). The clinical presentation of disorders, including abnormal skin and hair phenotypes, may be attributed, at least in part, to a non-functional Ly6 superfamily member such as E48 or HROAD63. HROAD63 may also be involved in blood disorders, as seen with its homologues SCA-2 and CD59.

A novel prohibitin cDNA clone presented herein was originally identified in a human bone marrow cell line (RS4;11) cDNA library. The clone contains a 1066 nucleotide insert which encodes a 299 amino acid polypeptide. BLAST and BestFit analyses of the predicted amino acid sequence of HMWGS46 demonstrate a highly significant sequence identity to a murine protein termed IgM B-cell receptor associated protein (BAP)-37 (Genbank accession number X78683). The HMWGS46 amino acid sequence exhibits nearly perfect identity and similarity over the entire length of the murine BAP-37 sequence. (See Figure 14.) In addition, the full-length nucleotide sequences of HMWGS46 and BAP-37 exhibit at least 87% identical. The HMWGS46 clone also exhibits approximately 49% sequence identity and 85% sequence similarity to a human gene designated prohibitin. Finally, the HMWGS46 cDNA appears in a substantial number of HGS human cDNA libraries in addition to the bone marrow cell line cDNA library from which it was cloned. Some of the cDNA libraries in which this clone appears include keratinocytes, induced endothelial cells, activated neutrophils, synovial sarcoma, colon carcinoma cell line, Jurkat cell line membrane bound polysomes, epileptic frontal cortex, primary dendritic cells, and a number of others. The novel gene related to prohibitin and BAP-37 may prove quite useful as a diagnostic for tumorigenesis, as well as a target for therapeutic intervention of such an event. Thus, although the precise functional role of the prohibitin family members are less than clear, it is quite likely that such homologues are involved in such complex processes as development, senescence, and tumor suppression. Therefore a novel gene, such as HMWGS46, may prove quite useful as a diagnostic for tumorigenesis, as well as a target for therapeutic intervention of such an event.

A human cDNA clone encoding a novel epidermal growth factor receptor (EGFR)-like molecule is also disclosed. The novel EGFR-like cDNA clone presented herein was originally identified in an activated human neutrophil cDNA library. The

5

10

15

20

25

30

clone contains a 704 nucleotide insert which encodes a 168 amino acid polypeptide. A BLAST analysis of the predicted amino acid sequence of **HNFGW06** demonstrates that this novel clone exhibits approximately 85% identity and 90% similarity to a protein designated epidermal growth factor receptor-related protein [Homo sapiens]. (See Figure 15.) The expression profile of the HNFGW06 clone in the HGS database indicates the existence of a fairly highly restricted expression pattern. In addition to the activated neutrophil library from which this clone was obtained, it also appears in the following HGS human cDNA libraries: synovial sarcoma, smooth muscle, placenta, and possibly primary dendritic cells.

The novel EGFR-like cDNA clone HNFGW06 may lead to a number of exciting possibilities for therapeutic and/or diagnostic treatments or reagents. For example, HNFGW06 may be involved in the onset of human breast cancers as well. In addition, due to the fact that TGF-a acts through binding to the EGFR, it is possible that HNFGW06 may also play a role in a variety of gastric processes including regulation of acid secretion, regulation of mucous cell growth, and protection against ethanol- and aspirin-induced injury to gastric tissues.

#### GENERATING POLYNUCLEOTIDES

Polynucleotides of the present invention encoding a receptor may be obtained using standard cloning and screening, from a cDNA library derived from mRNA in cells specified in Table 1 using the expressed sequence tag (EST) analysis (Adams, M.D., et al. Science (1991) 252:1651-1656; Adams, M.D. et al., Nature, (1992) 355:632-634; Adams, M.D., et al., Nature (1995) 377 Supp:3-174.) Polynucleotides of the invention can also be obtained from natural sources such as genomic DNA libraries or can be synthesized using well known and commercially available techniques.

The nucleotide sequence encoding a receptor polypeptide of SEQ ID NO:Y may be identical to the polynucleotide encoding SEQ ID NO:Y, or it may be a sequence, which as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:Y.

When the polynucleotides of the invention are used for the recombinant production of a receptor polypeptide, the polynucleotide may include the coding sequence for the mature polypeptide or a fragment thereof, by itself; the coding sequence for the mature polypeptide or fragment in reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro- protein sequence, or other fusion peptide portions. For example, a marker sequence which facilitates purification of the fused polypeptide can be encoded. In certain preferred embodiments of this aspect of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in

5

10

15

20

. 25

30

10

15

20

25

30

35

Gentz et al., Proc Natl Acad Sci USA (1989) 86:821-824, or is an HA tag. The polynucleotide may also contain non-coding 5' and 3' sequences, such as transcribed, non-translated sequences, splicing and polyadenylation signals, ribosome binding sites and sequences that stabilize mRNA.

Further preferred embodiments are polynucleotides encoding receptor variants comprising the amino acid sequence of receptor polypeptide of Table 1 (SEQ ID NO:Y) in which several, 5-10, 1-5, 1-3, 1-2 or 1 amino acid residues are substituted, deleted or added, in any combination.

The present invention further relates to polynucleotides that hybridize to the herein above-described sequences. In this regard, the present invention especially relates to polynucleotides which hybridize under stringent conditions to the herein above-described polynucleotides. As herein used, the term "stringent conditions" means hybridization will occur only if there is at least 80%, and preferably at least 90%, and more preferably at least 95%, yet even more preferably 97-99% identity between the sequences.

Polynucleotides of the invention, which are identical or sufficiently identical to a nucleotide sequence contained in SEQ ID NO:X or a fragment thereof, or to the cDNA insert in the plasmid deposited at the ATCC, or a fragment thereof, may be used as hybridization probes for cDNA and genomic DNA, to isolate full-length cDNAs and genomic clones encoding the receptor and to isolate cDNA and genomic clones of other genes (including genes encoding homologs and orthologs) that have a high sequence similarity to the receptor gene. Such hybridization techniques are known to those of skill in the art. Typically these nucleotide sequences are 80% identical, preferably 90% identical, more preferably 95% identical to that of the referent. The probes generally will comprise at least 15 nucleotides. Preferably, such probes will have at least 30 nucleotides and may have at least 50 nucleotides. Particularly preferred probes will range between 30 and 50 nucleotides.

In one embodiment, to obtain a polynucleotide encoding the receptor polypeptide, including homologs and orthologs from other species, comprises the steps of screening an appropriate library under stringent hybridization conditions with a labeled probe having the SEQ ID NO:X or a fragment thereof; and isolating full-length cDNA and genomic clones containing said polynucleotide sequence. Such hybridization techniques are well known to those of skill in the art. Stringent hybridization conditions are as defined above or, alternatively, conditions under overnight incubation at 42°C in a solution comprising: 50% formamide. 5xSSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10 % dextran sulfate, and 20 microgram/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65°C.

The polynucleotides and polypeptides of the present invention may be employed as research reagents and materials for discovery of treatments and diagnostics to animal and human disease.

# 5 Vectors, Host Cells, Expression

10

15

20

25

30

35

The present invention also relates to vectors which comprise a polynucleotide or polynucleotides of the present invention, and host cells which are genetically engineered with vectors of the invention and to the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention.

For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof for polynucleotides of the present invention. Introduction of polynucleotides into host cells can be effected by methods described in many standard laboratory manuals, such as Davis et al., BASIC METHODS IN MOLECULAR BIOLOGY (1986) and Sambrook et al., MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989) such as calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction or infection.

Representative examples of appropriate hosts include bacterial cells, such as streptococci, staphylococci, E. coli, Streptomyces and Bacillus subtilis cells; fungal cells, such as yeast cells and Aspergillus cells; insect cells such as Drosophila S2 and Spodoptera Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, HEK 293 and Bowes melanoma cells; and plant cells.

A great variety of expression systems can be used. Such systems include, among others, chromosomal, episomal and virus-derived systems, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression systems may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides to produce a polypeptide in a host may be used. The appropriate nucleotide sequence may be inserted into an expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook et al., MOLECULAR CLONING, A LABORATORY MANUAL (supra).

For secretion of the translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the desired polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

If the receptor polypeptide is to be expressed for use in screening assays, generally, it is preferred that the polypeptide be produced at the surface of the cell. In this event, the cells may be harvested prior to use in the screening assay. If the receptor polypeptide is secreted into the medium, the medium can be recovered in order to recover and purify the polypeptide; if produced intracellularly, the cells must first be lysed before the polypeptide is recovered.

Receptor polypeptides can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during isolation and or purification.

## Diagnostic Assays

5

10

15

20

25

30

35

This invention also relates to the use of receptor polynucleotides or polypeptides for use as diagnostic reagents. Detection of a mutated form of the receptor gene associated with a dysfunction will provide a diagnostic tool that can add to or define a diagnosis of a disease or susceptibility to a disease which results from underexpression, over-expression or altered expression of the receptor. Individuals carrying mutations in the receptor gene may be detected at the DNA level by a variety of techniques.

Nucleic acids for diagnosis may be obtained from a subject's cells, such as from blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification techniques prior to analysis. RNA or cDNA may also be used in similar fashion. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to labeled receptor nucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in electrophoretic mobility of DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing. (See, e.g., Myers et al., Science (1985) 230:1242.) Sequence changes at specific locations may also be revealed by

nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method. (See Cotton et al., Proc Natl Acad Sci USA (1985) 85: 4397-4401.) In another embodiment, an array of oligonucleotides probes comprising receptor nucleotide sequence or fragments thereof can be constructed to conduct efficient screening of e.g., genetic mutations. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability. (See for example: M.Chee et al., Science, Vol 274, pp 610-613 (1996).)

The diagnostic assays offer a process for diagnosing or determining a susceptibility to specific diseases through detection of mutation in the receptor gene by the methods described.

In addition, specific diseases can be diagnosed by methods comprising determining from a sample derived from a subject an abnormally decreased or increased level of receptor polypeptide or receptor mRNA. Decreased or increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example, PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods. Assay techniques that can be used to determine levels of a protein in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

Thus in another aspect, the present invention relates to a diagnostic kit for a disease or susceptibility to a disease which comprises:

- (a) a receptor polynucleotide, preferably the nucleotide sequence of SEQ ID NO:X, or a fragment thereof;
  - (b) a nucleotide sequence complementary to that of (a);
- (c) a receptor polypeptide, preferably the polypeptide of SEQ ID NO:Y, or a fragment thereof; or
- (d) an antibody to a receptor polypeptide, preferably to the polypeptide of SEQ ID NO: Y.
- It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

#### Chromosome Assays

The nucleotide sequences of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome. The mapping of relevant sequences to chromosomes according to the present invention is an important first step in correlating those sequences with gene associated disease. Once a sequence has been mapped to a precise chromosomal location, the physical position of the

5

10

15

20

25

sequence on the chromosome can be correlated with genetic map data. Such data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes).

The differences in the cDNA or genomic sequence between affected and unaffected individuals can also be determined. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

10

15

20

25

30

5

#### Antibodies

The polypeptides of the invention or their fragments or analogs thereof, or cells expressing them can also be used as immunogens to produce antibodies immunospecific for the receptor polypeptides. The term "immunospecific" means that the antibodies have substantially greater affinity for the polypeptides of the invention than their affinity for other related polypeptides in the prior art.

Antibodies generated against the receptor polypeptides can be obtained by administering the polypeptides or epitope-bearing fragments, analogs or cells to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler, G. and Milstein, C., Nature (1975) 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor et al., Immunology Today (1983) 4:72) and the EBV-hybridoma technique (Cole et al., MONOCLONAL ANTIBODIES AND CANCER THERAPY, pp. 77-96, Alan R. Liss, Inc., 1985).

Techniques for the production of single chain antibodies (U.S. Patent No. 4,946,778) can also be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms including other mammals, may be used to express humanized antibodies.

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or to purify the polypeptides by affinity chromatography. Antibodies against receptor polypeptides may also be employed to treat diseases.

## Vaccines

Another aspect of the invention relates to a method for inducing an immunological response in a mammal which comprises inoculating the mammal with a receptor polypeptide, or a fragment thereof, adequate to produce antibody and/or T cell immune response to protect said animal from a disease. Yet another aspect of the invention relates to a method of inducing immunological response in a mammal which

comprises, delivering a receptor polypeptide via a vector directing expression of the receptor polynucleotide in vivo in order to induce such an immunological response to produce antibody to protect said animal from diseases.

Further aspect of the invention relates to an immunological/vaccine formulation (composition) which, when introduced into a mammalian host, induces an immunological response in that mammal to a receptor polypeptide wherein the composition comprises a receptor polypeptide or receptor gene. The vaccine formulation may further comprise a suitable carrier. Since a receptor polypeptide may be broken down in the stomach, it is preferably administered parenterally (including subcutaneous, intramuscular, intravenous, intradermal etc. injection). Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation instonic with the blood of the recipient; and aqueous and nonaqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

#### Screening Assays

5

10

15

20

25

30

35

The receptor polypeptide of the present invention may be employed in a screening process for compounds which bind the receptor and which activate (agonists) or inhibit activation of (antagonists) the receptor polypeptide of the present invention. Thus, polypeptides of the invention may also be used to assess the binding of small molecule substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural substrates and ligands or may be structural or functional mimetics. See Coligan et al., Current Protocols in Immunology 1(2):Chapter 5 (1991).

The receptor polypeptides are responsible for many biological functions, including many pathologies. Accordingly, it is desirous to find compounds and drugs which stimulate the receptor on the one hand and which can inhibit the function of the receptor on the other hand. In general, agonists are employed for therapeutic and prophylactic purposes for such conditions and diseases. Antagonists may be employed for a variety of therapeutic and prophylactic purposes for such conditions and diseases.

In general, such screening procedures involve producing appropriate cells which express the receptor polypeptide of the present invention on the surface thereof.

10

15

20

25

30

Such cells include cells from mammals, yeast, Drosophila or E. coli. Cells expressing the receptor (or cell membrane containing the expressed receptor) are then contacted with a test compound to observe binding, or stimulation or inhibition of a functional response.

The assays may simply test binding of a candidate compound wherein adherence to the cells bearing the receptor is detected by means of a label directly or indirectly associated with the candidate compound or in an assay involving competition with a labeled competitor. Further, these assays may test whether the candidate compound results in a signal generated by activation of the receptor, using detection systems appropriate to the cells bearing the receptor at their surfaces. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed.

Further, the assays may simply comprise the steps of mixing a candidate compound with a solution containing a receptor polypeptide to form a mixture, measuring receptor activity in the mixture, and comparing the receptor activity of the mixture to a standard.

The receptor cDNA, protein and antibodies to the protein may also be used to configure assays for detecting the effect of added compounds on the production of receptor mRNA and protein in cells. For example, an ELISA may be constructed for measuring secreted or cell associated levels of receptor protein using monoclonal and polyclonal antibodies by standard methods known in the art, and this can be used to discover agents which may inhibit or enhance the production of the receptor (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues. Standard methods for conducting screening assays are well understood in the art.

Examples of potential receptor antagonists include antibodies or, in some cases, oligonucleotides or proteins which are closely related to the ligand of the receptor, e.g., a fragment of the ligand, or small molecules which bind to the receptor but do not elicit a response, so that the activity of the receptor is prevented.

Thus in another aspect, the present invention relates to a screening kit for identifying agonists, antagonists, ligands, receptors, substrates, enzymes, etc. for receptor polypeptides; or compounds which decrease or enhance the production of receptor, which comprises:

- (a) a receptor polypeptide, preferably that of SEQ ID NO:Y;
- (b) a recombinant cell expressing a receptor polypeptide, preferably that of SEQ35 ID NO:Y;
  - (c) a cell membrane expressing a receptor polypeptide; preferably that of SEQ ID NO: Y; or
    - (d) antibody to a receptor polypeptide, preferably that of SEQ ID NO: Y.

SUBSTITUTE SHEET (RULE 26)

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

# Prophylactic and Therapeutic Methods

5

10

15

20

25

30

35

This invention provides methods of treating an abnormal conditions related to both an excess of and insufficient amounts of receptor activity.

If the activity of the receptor is in excess, several approaches are available. One approach comprises administering to a subject an inhibitor compound (antagonist) as described along with a pharmaceutically acceptable carrier in an amount effective to inhibit activation by blocking the binding of ligands to the receptor or by inhibiting a second signal, and thereby alleviating the abnormal condition.

In another approach, soluble forms of the receptor polypeptides still capable of binding the ligand in competition with endogenous receptor may be administered. Typical embodiments of such competitors comprise fragments of the receptor polypeptide.

In still another approach, expression of the gene encoding endogenous receptor can be inhibited using expression blocking techniques. Known such techniques involve the use of antisense sequences, either internally generated or separately administered. (See, for example, O'Connor, J Neurochem (1991) 56:560 in Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca-Raton, FL (1988).) Alternatively, oligonucleotides which form triple helices with the gene can be supplied. (See, for example, Lee et al., Nucleic Acids Res (1979) 6:3073; Cooney et al., Science (1988) 241:456; Dervan et al., Science (1991) 251:1360.) These oligomers can be administered per se or the relevant oligomers can be expressed in vivo.

For treating abnormal conditions related to an under-expression of the receptor and its activity, several approaches are also available. One approach comprises administering to a subject a therapeutically effective amount of a compound which activates the receptor, i.e., an agonist as described above, in combination with a pharmaceutically acceptable carrier, to thereby alleviate the abnormal condition. Alternatively, gene therapy may be employed to effect the endogenous production of the receptor by the relevant cells in the subject. For example, a polynucleotide of the invention may be engineered for expression in a replication defective retroviral vector, as discussed above. The retroviral expression construct may then be isolated and introduced into a packaging cell transduced with a retroviral plasmid vector containing RNA encoding a polypeptide of the present invention such that the packaging cell now produces infectious viral particles containing the gene of interest. These producer cells may be administered to a subject for engineering cells in vivo and expression of the polypeptide in vivo. For overview of gene therapy, see Chapter 20, Gene Therapy and

15

20

25

30

35

other Molecular Genetic-based Therapeutic Approaches, (and references cited therein) in Human Molecular Genetics, T Strachan and A P Read, BIOS Scientific Publishers Ltd (1996).

#### 5 Formulation and Administration

Peptides, such as the soluble form of receptor polypeptides, and agonists and antagonist peptides or small molecules, may be formulated in combination with a suitable pharmaceutical carrier. Such formulations comprise a therapeutically effective amount of the polypeptide or compound, and a pharmaceutically acceptable carrier or excipient. Such carriers include but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. Formulation should suit the mode of administration, and is well within the skill of the art. The invention further relates to pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

Polypeptides and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds. Preferred forms of systemic administration of the pharmaceutical compositions include injection, typically by intravenous injection. Other injection routes, such as subcutaneous, intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transmucosal and transdermal administration using penetrants such as bile salts or fusidic acids or other detergents. In addition, if properly formulated in enteric or encapsulated formulations, oral administration may also be possible. Administration of these compounds may also be topical and/or localized, in the form of salves, pastes, gels and the like.

The dosage range required depends on the choice of peptide, the route of administration, the nature of the formulation, the nature of the subject's condition, and the judgment of the attending practitioner. Suitable dosages, however, are in the range of 0.1-100 µg/kg of subject. Wide variations in the needed dosage, however, are to be expected in view of the variety of compounds available and the differing efficiencies of various routes of administration. For example, oral administration would be expected to require higher dosages than administration by intravenous injection. Variations in these dosage levels can be adjusted using standard empirical routines for optimization, as is well understood in the art.

Polypeptides used in treatment can also be generated endogenously in the subject, in treatment modalities often referred to as "gene therapy" as described above. Thus, for example, cells from a subject may be engineered with a polynucleotide, such as a DNA or RNA, to encode a polypeptide ex vivo, and for example, by the use of a retroviral plasmid vector. The cells are then introduced into the subject.

SUBSTITUTE SHEET (RULE 26)

All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

						-																		11.	771	ĿD	110	ノ		
CAC	TT	CCT	GAG/	GA.	AGA	ACC	TGA	GG.	AAC	AGAC	GTI	rcc	CTC	GCG	GCCC	TG	GCA	CCT	CCA	ACC	CCA	GAT,	ATGO	TGC	TG	CTG	TGC	TGC	TG	90
GTO	AA	GGA	CTC	CT	TCT	TGG	SACI	СС	TTG	тст	GC A A	AGG	GAG	cgc	CGGC	AC	CGT	GGA	GGŤ.	TGG	GGT	СТА	TAC	GACO	AC	GAC	GAC	SAC	AC	30
																							М	L	L	Ļ	L	L	L	
						GAG			~ ^ ^	C C A /	- A C		4 C T	4 4 C			C A T	TAC	T C C I	CTC	۸.C.C	ATC	C V C :	л <del>г</del> т т		CIC	۸۲۲		- ^ ^	
						CTC							+			<del></del>										<del></del> -				180
GGG	AC	GAG	ALCE	ما ما د	166	t I C	1666	.AL	CII	CCI	يا ا د	116	ILA	116	666	110	LIA	AIG.	AGC	GAC	166	IAC	<b>.</b>	ICAA	466	CAC	1661	ACI	111	
Р	L	L	W	G	R	Ε	R	٧	ξ	G	Q	K	S	N	R	K	0	Υ	S	L	Т	М	0	S	S	٧	T	٧	a	
GAG	GGC	ATG	ŢĠŦſ	GTC	CAT	GTG	cgc:	rgc	TCC	TTC	TCC	TAC	ÇCA	GTG	GAC	AGC	CAG	ACT	GAC	тст	GAC	CCA	GTT	CATO	GGC	TAC	TGG	TTC	GG	270
CTC	CCG	TAC	ACA	CAG	GTA	CAC	GC G	AC G	AGG	AAG	AGG	ATG	GGT	CAC	CTG	TÇG	GTC	TGA	CTG	AGA	CTG	GGT	CAA	GTA	CCG	ATG	ACC	AAG	3CC	270
Ε	G	М	С	٧	н	٧	R	С	s	F	S	Y	Р	٧	0	s	Q	Т	D	s	D	Ρ	٧	н	G	Y	W	F	R	
			C 4 T	A T A	400	TGG	A A C	- с т	C C A	CTC	000	۸ ۲ ۸	4 A C	A A C	CCAI	cc t	TCC	ירר א	CTC	CAC	C 4 C	CAA	۸CT	caai	3 4 C	rca.	TTC	۲۵۲۱	רדר	
			+			ACC		+										+					+			<del></del>				360
<u>,                                    </u>		. I I A	CIA	1 A :	100	ACC	110	LUA		CAC	CGG		110	. , , ,	aa.	Cun	ACC		٠٨٠							_	_			
Α	G	N	D	i	S	W	K	Α	Р	٧	A	T	N	N	P	А	₩	Α	٧	Q	Ε	Ε	٢	R	0	R	F	н	L	
CTT	GGG	GAC	ÇCA	CAG	ACC	AAA	AAT	TGC	ACC	CTG	AGC	ATC	AGA	GAT	GCC	AGA	ATG	AGT	GAT	GCG	GGG	AGA	TAC	TTC	TTT	CGT	ATG	GAG.	AAA	450
GAA	ccc	ÇTG	ĠGT	GTC	TGG	TTT	TTA	ACG	TGG	GAC	TCG	TAG	TCT	CTA	CGG	TÇT	TAC	TCA	CTA	CGC	CCC	TCT	ATG	AAG	AAA	AGCA	TAC	CTC	TTT	
L	G	D	P	a	Ţ	κ	N	С	T	L	s	l	R	D	Α	R	М	s	ם	Α	G	R	Υ	F	F	R	M	Ε	ĸ	
GGA	AAT	ΓΑΤΑ	AAA	TGG	AAT	TAT	AAA	TAT	'GAC	CAG	СТС	TCI	rgre	AAC	GTG	ACA	TAC	CCT	CCT	CAG	AAC	TTG	ACT	GTG	ACT	rgro	TTC	CAA	GGA	
						ATA										-		<del></del>	<del>+</del>											540
						Y				•																				
						GCT																							<del>+</del>	630
CTT	CCC	STGT	CGT	AGG	STGT	CGA	GAC	CCC	CTTC	STCG	AGT	AGA	AGAA	AAGT	CAG	GAI	СТС	CCG	GTC	AGA	GAC	GCC	IAAC	CAG	ACA	ACGA	LAA	(16	106	
٤	G	Τ,	Α	S	T	Α	L	G	N	S	S	S	Ļ	S	٧	L	Ε	G	Q	S	L	R	L	٧	С	Α	٧	0	S	
AAT	cco	CCI	rgcc	AGO	SCTO	GAGC	TGG	ACC	CTG	GAGG	AGT	СТ	GACO	сто	TAC	cçc	TCA	ACAG	ccc	TCA	AAC	CCT	ĊTG	GTA	CTO	GGAC	CTG	CAA	GTG	720
TTA	GG	GGG	/C G G	TC	CGAC	TCG	ACC	TG	GAC	TCC	TCA	GA	CTG	GGAC	ATG	GGG	SAG	TGTO	GGC	AGT	TTO	GGG/	GAC	CAT	GA	CCTO	GAC	GTT	CAC	720
N	Ρ	Ρ	Α	R	L	s	W	τ	W	R	S	L	T	L	Υ	Р	s	Q	Р	s	N	Ρ	L	٧	L	Ε	L	Q	٧	
					_																	- c + /		·	T C .	+ <i>i</i>		C 4 C		
						GAA				-			-+-				—-													810
						CTI																						316		
u	1	C	n	F	C	E	F	τ	٦	R	Δ	Λ	N	S	ŧ	G	S	Ð	н	V	S	L	N	L	S	L	a	a	Ε	

Page 1

																							H	TE	D	KΥ	18			
ÇGGC	AC	GAG	TGG	ACA,	ACC.	ATC.	AGG	GAG	CCĂ	GGA	CAC	AGA	GGG	GCA	GAG	:AA(	GTC/	AGCA	ATTO	GC	GC C	CCT	TEE	TCA	GAT	CC.	TATO	ATC	TŢ	
GCC	TG	CTC.	ACC	TGT	TGG	TAG	TCC	CTC	GGT	CCT	GTG	TCT	CCC	CGT	CTC	GTT(	CAG	TCG	ΓΑΑ	CG	CGG	GGA.	AGG	AGT	CTA	GGG	ATA	STAG		90
GGG						-+-			<del></del>				+	<del></del>		<del></del>							<del></del>	<del></del>						180
CCC.	TTT	GTC.	ATE	GGG	TCT	CCA.	AGT	CCT	TCT	ACA	ATT	GAA	TTT	ACA.	AGC	200	ACG	GGG.	TCA	GAC.	AAG	TCG	TAC	CGA	CTT	TAG	GTG'	TGAG	GC	
																							М	Α	Ε	i	Н	T	Р	
TAT	гст	TCC	TTG	AAG.	ААА	CŢG	TTA	TCT	TTA	стс	AAT	GGC	TTC	GTG	GCT	GTG	TCT	GGC	ATC.	ATC	СТА	GTT	ggc	CTG	GGC.	ATT	GGT	GGTA	AAA	270
ATA	AGA.	AGG	AAC	TTC	TTT	GAC	AAT	AGA	AAT	GAG	TTA	CCG	AAG	CAC	CGA	CAC	AGA	CCG	TAG	TAG	GAT	CAA	cca	GAC	CCG	TAA	CCA	CCA1	ΓΤΤ	210
Y	s	s	L	Κ	κ	L	L.	S	L	L	N	G	F	٧	Α	٧	S	G	I	I	L	٧	S	L	G	į	G	G	K	
TGT	GGA	GGG	GCC	TCT	CTG	ACG	AAT	GTC	стс	GGG	CTG	TCC	TCC	GCA	TAC	стс	CTT	CAC	GTT	GGC	AAC	CTG	TGC	CTG	GTG	AŢG	GGA	TGCA	ATÇ	
ACA	CCT	ССС	cgg	AGA	GAC	TGC	TTA	CAG	GAG	ССС	GAC	AGG	AGG	CGT	ATG	GAG	GAA	GTG	CAA	CCG	TTG	GAC	ACG	GAC	CAC	TAC	CCT	ACG1	TAG	360
С	G	G	А	S	L	T	N	٧	L	G	Ļ	s	s	A	Υ	L	L	Н	٧	G	N	L	С	L	٧	м	G	С	Ī	
ACG	GTA	CTG	CTT	GGC	TGT	GCC	GGG	TGG	TAT	GGA	GCG	ACT	AAA	GAG	AGC	AGA	GGC	ACG	стс	TTG	ттт	TGC	ĄTC	CTG	TCA	ATG	GTT	ATT(	зтç	
TGC	CAT	GAC	GAA	CCG	ACA	CGG	ccc	ACC	ATA	CCT	CGC	TGA	TTT	стс	TCG	TCT	CCG	TGC	GAG	AAC	AAA	ACG	TAG	GAC	AGT	TAC	CAA	TAA	CAG	450
Т	٧	L	Ļ	G	С	Α	G	W	Υ .	G	Α	T	к	Ε	s	R	G	т	L	L	F	С	ī	L	s	М	٧	i	٧	
стс	ATC	ATG	GAA	GTT	ACA	GCT	GCC	ACA	GTG	GTC	CTT	CTT	TTC	TTT	CCA	ATT	GTT	GĞA	GAT	GTG	GCC	TTG	GAA	CAC	ACC	TTC	GTG	ACC	CTG	
GAG	TAG	TAC	CTT	CAA	TGT	CGA	CGG	TGT	CAC	CAG	GAA	GAA	AAG	AAA	GGT	TAA	CAA	CCT	CTA	CAC	CGG	AAC	CTI	GTG	TGG	AAG	CAC	TGG	GAC	540
L	I	М	Ε	٧	T	A	Α	T	٧	٧	L	L	F	F	Р	i	٧	G	0	٧	Α	L	٤	н	Ţ	F	٧	T	L	
AGG	AAG	AAT	TAC	AGA	GGT	TAC	AAC	GAG	CCA	GAC	GAC	TAT	TCT	ACA	CAG	TGG	AAC	TTG	GTC	ATG	GAG	AAG	CTA	AAG	TGC	TGT	GGG	GTG.	AAT	630
TCC	TTC	TTA	ATG	TCT	CCA	ATG	TTG	CTC	GGT	CTG	CTG	ATA	AGA	TGT	GTC	ACC	TTG	AAC	CAG	TAC	СТС	TTC	GAT	TTC	ACG	ACA	ccc	CAC	TTA	-
R	K	N	Y	R	G	Y	N	Ε	Ρ	D	D	Υ	s	T	Q	W	N	L	٧	М	Ε	Κ	L	Κ	С	С	G	٧	N	
AAC	TAC	ACA	GAT	777	тст	GGC	TCT	тсс	TTÇ	GAA	ATG	ACA	ACG	GGC	CAC	AÇC	TAC	ccc	AGĢ	AGT	TGC	TGT	AAA	TCC	ATC	GGA	AGT	GTG	TCC	720
TTG	ATG	TGT	CTA	AAA	AGA	CCG	AGA	AGC	SAAG	CTT	TAC	TGT	TGC	CCG	GTG	TGG	ΑTG	GGG	TCC	TCA	ACG	ACA	TT	AGG	TAG	ССТ	TCA	CAC	AGĞ	720
N	Y	T	0	F	S	G	S	S	F	٤	М	Т	Т	G	н	T	Y	Р	R	S	С	С	Κ	S	l	G	S	٧	S	
TGT	GAC	GGA	, ÇGC	GAT	GTG	TÇT	CCA	AAC	CGTC	ATC	CAC	CAC	SAAG	GGC	TGT	TTC	CAT	AAA	СТС	СТА	AAA	ATC	ACC	AAG	ACT	CAG	AGC	TTC	ACC	810
ACA	CTG	CCT	GCG	CTA	CAC	AGA	GGT	TTC	CAG	TAG	GTG	GTC	TTC	CCG	ACA	AAG	GTA	TTT	GAG	GAT	TTT	TAG	TGC	TTC	TGA	GTC	TCG	AAG	TGĠ	010
С	D	G	R	0	٧	s	Ρ	N	٧	i	Н	Q	κ	G	С	F	н	К	L	L	ĸ	Į	ī	ĸ	T	Q	s	F	T /c	es ID
CTG	AGT	GGG	ĄGO	TCT	сто	GGA	AGC 1	rgc/	AGTG	SATA	CAC	AGC	TGC	GGG	TCT	cĢc	TAT	GTT	GCG	CAC	GCT	GGT	rc T1	GAA	CTO	CTO	GCC	TAA	AĞC	900
GAC	TCA	CCC	TCC	AGA	GAC	:001	C G A	\CGT	TCAC	TAT	GTO	CTCC	ACC	ccc	AGA	GCG	ATA	CAA	CGC	GTC	CGA	ACC/	AGA	ACT1	rgac	GAC	CGG	ATT		
L	S	G	s	s	L	G	А	A	٧	1.	Q	R	.W	G	s	R	Y	٧	Α	۵	Α	G	L	ε	Ļ	L	A	. (		DID 0.19

HFWA E 45	
GGCGTCCCTCTGCCTGCCCACTCAGTGGCAACACCCGGGAGCTGTTTTGTCCTTTGTGGAGCCTCAGCAGTTCCCTCTTTCAGAACTC	
CCGCAGGGAGACGGACGGTGAGTCACCGTTGTGGGCCCTCGACAAAACAGGAAACACCTCGGAGTCGTCAAGGGAGAAAGTCTTGAG	.C + 90
TGCCAAGAGCCCTGAACAGGAGCCACCATGCAGTGCTTCAGCTTCATTAAGACCATGATGATCCTCTTCAATTTGCTCATCTTTCTGT	
ACGGTTCTCGGGACTTGTCCTCGGTGGTACGTCACGAAGTCGAAGTAATTCTGGTACTACTAGGAGAAGTTAAACGAGTAGAAAGACA	+ 180 :A
	•
MQCFSFIKTMMILFNLLIFL	v
GGTGCAGCCCTGTTGGCAGTGGGCATCTGGGTGTCAATCGATGGGGCATCCTTTCTGAAGATCTTCGGGCCACTGTCGTCCAGTGCCA	rg → 270
CCACGTCGGGACAACCGTCACCCGTAGACCCACAGTTAGCTACCCCGTAGGAAAGACTTCTAGAAGCCCGGTGACAGCAGGTCACGGT	
G A A L L A V G I W V S I D G A S F L K I F G P L S S S A	1
CAGTITGTCAACGTGGGCTACTTCCTCATCGCAGCCGGCGTTGTGGTCTTTGCTCTTGGTTTTCCTGGGCTGCT	<del>-+</del> 360
GTCAAACAGTTGCACCCGATGAAGGAGTAGCGTCGGCCGCAACACCAGAAACGAGAACCAAAGGACCCGACGA	. <b>L</b>
Q F V N V G Y F L I A A G V V V F A L G F L G C Y G A K T	Ē
AGCAAGTGTGCCCTCGTGACGTTCTTCTTCATCCTCCTCCTCATCTTCATTGCTGAGGTTGCAGCTGCTGTGGTCGCCTTGGTGTACA	
TCGTTCACACGGGAGCACTGCAAGAAGAAGTAGGAGGAGGAGTAGAAGTAACGACTCCAACGTCGACGACACCAGCGGAACCACATGT	→ 450 3G
	Т
SKEALVIPPILLLIPIAEVAAAVVALVI	ı
ACAATGGCTGAGCACTTCCTGACGTTGCTGGTAGTGCCTGCC	4C -+ 540
TGTTACCGACTCGTGAAGGACTGCAACGACCATCACGGACGG	rG
T.M. A.E.H.F.L.T.L.V.V.P.A.I.K.D.Y.G.S.Q.E.D.F.T.Q.V.W.	N
ACCACCATGAAAGGGCTCAAGTGCTGTGGCTTCACCAACTATACGGATTTTGAGGACTCACCCTACTTCAAAGAGAACAGTGCCTTTC	rŕ
TGGTGGTACTTTCCCGAGTTCACGACACCGAAGTGGTTGATATGCCTAAAACTCCTGAGTGGGATGAAGTTTCTCTTGTCACGGAAAG	++ 630
-	
T T M K G L K C Ć G F T N Y T D F E D S P Y F K E N S A F	P
CCATTCTGTTGCAATGACAACGTCACCAACACACCCAATGAAACCTGCACCAAGCAAAAAGGCTCACGACCAAAAAGTAGAGGGTTGCT	TC 720
GGTAAGACAACGTTACTGTTGCAGTGGTTGTGTCGGTTACTTTGGACGTGGTTCGTTTTCCGAGTGCTGGTTTTTCATCTCCCAACGA	<del>-+</del> 720 AG
	F
·	
AATCAGCTTTTGTATGACATCCGAACTAATGCAGTCACCGTGGGTGTGGCAGCTGGAATTGGGGGCCTCGAGCTGCCATGA	<del>-+</del> 810
TTAGTCGAAAACATACTGTAGGCTTGATTACGTCAGTGGCACCCACC	ΓΑ
NOLLYDIRTNAVTVGGVAAGIGGLELAAM	N
TGTGTCCATGTATCTGTACTGCAATCTACAATAAGTCCACTTCTGCCTCTGCCACTACTGCTGCCACATGGGAACTGTGAAGAGGCAC	СС
ACACAGGTACATAGACATGACGTTAGATGTTATTCAGGTGAAGACGGAGACGGTGATGACGACGGTGTACCCTTGACACTTCTCCGTG	<del>-+</del> 900
·	
C V L V C V L O C T L C D L L D L L L D H G N C F F A	ρ

Päge 1

																						t	T	٢		- 7	صا		
AAAAA	AACA	AGG	TCC	CCA	CAG	CAA	AGAA	AAA	GGA	ATA	GGA1	T C A	AGA	GAT.	ACG	TGC	сто	CTG	GCA	GAC	CAA	GCA	TGA	ΑT	TCG	ATGA	CTT	CA	90
 																													-
																							М	N	S	М	Т	s	
GCAGTT															A T.C	- C T (	^ T C /	\ <b>C</b> C C	r A G	: C A /	1 T T Z	TOT	ጉተገ	- ۵ ۲	G T G	cccc	:TG	TAT	
GCAGTT CGTCAA	CCG	STGG	CCA	ATT	CTG	TGT	IGG	1 G G	1166	CAC		A C A		CAA	# T A C	C A	C V C .	1000	o T O	CT	ΓΔΔ	. 4.0	GAG	STG	CAC	6660	GAC	<del></del>	180
CGTCAA	AGGC	CACC	GGI	IAA	GAC	ALA	ALL	ALL	ALL	. (, ) (,	666	16	1140			307													
A V	Ρ	Ą	A	N	S	٧	L	٧	٧	A	Ρ	Н	N	G	Y	ρ	٧	T	Р	G	I	М	5	Н	۷	Р	L	Y	
CCAAAC	CAGC	CAGC	CGC	AAG	ŢCC	ACC	TAG	TŢC	сто	GGA	ACC	CAI	CCTA	AGTT	TG	GTG	TCG.	AATO	TG	AAT	GGG	CAGO	CT	GTG	CAG	AAAI	GCT	CTG	270
GGTTT	STCG	GTCG	GCG	TTC	AGO	TGG	ATC	AAC	GGAC	CCT	TGG	GT	GGA	ГСАА	AC	CAC	AGC	TTAC	AC.	TTA	CCC	GTCC	GGA	CAC	GTC	TTT	CGA	GAC	
P N	S	Q	ρ	۵	٧	н	L	٧	ρ	G	N	Ρ	Р	s	L	٧	S	N	٧	N	G	G	Р	٧	۵	κ	A	L	
AAAGA			. c c T	· <b>+</b> ^ ^	2000	A	TCC	ΔG	۸۲۲	Λ T C Δ	TTC	נהר	CTG	GCTO	CAC	ATC	GGC	CTC	GGC	TCC	ATC	ATG	GCG	ACC	GTT	стс	GTA	.ggg	
TTTCT																													360
																											٧	G	
	G																												
GAATA																													450
CTTAT	GGAC	AGA	TAAA	AG T	AAG.	ATG	CCTC	:CG	AAA	GGG	AAGA	4 C C	CCT	CCG.	AAC	ACC	: A A A	TAG	TAA	AG I	CCI	AGA	GAG	AG	١Α٠	افاتاء	List	C 1 1	
E Y	L	s	i	s	F	Y	G	G	F	Ρ	F	W	G	G	L	W	F	I	Į	S	G	S	L,	S	٧	Α	Α	Ε	
AATCA	GCCA	TAT	TCT	TAT	TGC	CTG	CTG1	гст	GGC	AGT	TTG	GGC	TTG	AAC	AŢC	GTO	CAGI	GCA	ATC	TGO	TCT	GEA	GT1	GG	AGT	CATA	CTO	TTC	540
TTAGT	CGG	TATA	AGA	ATA	ACG	GAC	GAC	4GA	CCG	TCA	AAC	ccc	SAAC	TTG	TAC	C A	GTC	CGT	TAG	ACC	AG/	\CGT	CAA	ACC	TCA	GTAI	[GA(	BAAG	-
N O	) P	Y	S	Y	С	L	L	s	G	s	L	G	L	N	l	٧	S	Α	I	С	S	А	٧	G	٧	I	L	F	
ATCAC																										GAT.	rtc'	TGGC	
TAGTO																													
IAGIG	21617	AGAI	ILA	IAA	.666																					. !			
[ ]		L	S	I	₽		ρ	e.																					_
GTGCT																													
CACGA	ACGA	CCAG	SAAG	ACG	GAC	GAC	CTC	AAA	4000	TAC	CGT	AC	GCG	TAGA	AAG	GGT	GAA	ACC	SAC	GGT	CAA	CCA	GAC	GAC	AGT	TAG	iii	GILF	4
νı	LL	٨	F	С	L	L	Ε	F	G	1	Α	С	Α	S	S	Н	F	G	С	۵	L	Ą	С	C	: C	s S	S	N	
GTGA	GTGT	CATO	CTAT	CCA	<b>1</b> AA(	CATO	CTAT	GC	AGC	444(	CCCA	GT	GAT	CAC	ccc	AGA	ACC	GGT	GAC	стс	ACC	ACC.	AAG	TTA	ATTO	CAG	TGA	GAT	: + 810
CACT																													
	s v															. E		٧				Р	S	, ,		s s	_		

OANX. (SEQIDNO: 21)

GTCGTTCCTGTTCTGACAGCTGAGGAAACGTCTCTCCCACTGTTTGTACTCTCACCTTCATTCTCAATTCAGTCTAGGAAACCATGCTG CAGCAAGGACAAGACTGTCGACTCCTTTGCAGAGAGGGTGACAAACATGAGAGTGGAAGTAAGAAGTTAAGTCAGATCCTTTGGTACGAC

39

TTTCTCTATCAAGAAGAAGACAGAGATTTTAAACAGATGTTAACCAAGAGGGACTCCCTAGGGCACATGCATCAGCACATATGTGGGCAT 1080 AAAGAGATAGTTCTTCTGTCTCTAAAATTTGTCTACAATTGGTCTCCCTGAGGGATCCCGTGTACGTAGTCGTATACACCCGTA

CCAGCCTCTGGGGCCTTGGCACACCCATTCGTGTGCTCTGCTGCTGCATGTGAGCTTGTGGGTTAGAGGAACAAATATCTAGACATTCAATCT GGTCGGAGACCCCGGAACCGTGTGGGTAAGCACACGAGACGACGACGTACACTCGAACACCCAATCTCCTTGTTTATAGATCTGTAAGTTAGA

BNSDOCID: <WO\_\_\_9831799A2\_I\_>

Päge 1

HSRRF02	_
CACGAGCAGGGTCTCGGGCTAGTCATGGCGTCCCCGTCTCGGAGACTGCAGACTAAACCAGTCATTACTTGTTTCAAGAGCGTTCTGCTA	
GTGCTCGTCCCAGAGCCCGATCAGTACCGCAGGGGCAGAGCCTCTGACGTCTGATTTGGTCAGTAATGAACAAAGTTCTCGCAAGACGAT	
M A S P S R R L Q T K P V I T C F K S V L L	
ATCTACACTITTATTTTCTGGATCACTGGCGTTATCCTTCTTGCAGTTGGCATTTGGGGCAAGGTGAGCCTGGAGAATTACTTTTCTCTT	
TAGATGTGAAAATAAAAGACCTAGTGACCGCAATAGGAAGAACGTCAACCGTAAACCCCGTTCCACTCGGACCTCTTAATGAAAAGAGAA	
I Y T F I F W I T G V I L L A V G I W G K V S L E N Y F S L	
TTAAATGAGAAGGCCACCAATGTCCCCTTCGTGCTCATTGCTACTGGTACCGTCATTATTCTTTTGGGCACCTTTGGTTGTTTTGCTACC	
AATTTACTCTTCCGGTGGTTACAGGGGAAGCACGAGTAACGATGACCATGGCAGTAATAAGAAAACCCGTGGAAACCAACAAAACGATGG	
LNEKATŅ V PF V LIATGT V I I LLGT F G C F A T	
TGCCGAGCTTCTGCATGGATGCTAAAACTGTATGCAATGTTTCTGACTCTCGTTTTTTTGGTCGAACTGGTCGCCATCGTAGGATTT	)
ACGGCTCGAAGACGTACCTACGATTTTGACATACGTTACAAAGACTGAGAGCAAAAAAACCAGCTTGACCAGCGACGGTAGCATCCTAAA	
C D A S A W M I K I Y A M F L T L V F L V E L V A A I V G F	
C R A S A W II C R C I W W I E	
GTTTTCAGACATGAGATTAAGAACAGCTTTAAGAATAATTATGAGAAGGCTTTGAAGCAGTATAACTCTACAGGAGATTATAGAAGCCAT	)
CAAAAGTCTGTACTCTAATTCTTGTCGAAATTCTTATTAATACTCTTCCGAAACTTCGTCATATTGAGATGTCCTCTAATATCTTCGGTA	
V F R H. E I K M S F K N N Y E K A L K Q Y N S T G D Y R S H	
GCAGTAGACAAGATCCAAAATACGTTGCATTGTTGTGGTGTCACCGATTATAGAGATTGGACAGATACTAATTATTACTCAGAAAAAGGA	^
CGTCATCTGTTCTAGGTTTTATGCAACGTAACAACACCACAGTGGCTAATATCTCTAACCTGTCTATGATTAATAATGAGTCTTTTTCCT	,
AVUXIUNIENCOGVISINO	
TTTCCTAAGAGTTGCTGTAAACTTGAAGATTGTACTCCACAGAGAGATGCAGACAAAGTAAACAATGAAGGTTGTTTTATAAAGGTGATG	0
AAAGGATTCTCAACGACATTTGAACTTCTAACATGAGGTGTCTCTCTACGTCTGTTTCATTTGTTACTTCCAACAAAATATTTCCACTAC	
F P K S C C K L E D C T P Q R D A D K V N N E G C F I K V M	
ACCATTATAGAGTCAGAAATGGGAGTCGTTGCAGGAATTTCCTTTGGAGTTGCTTGC	
TGGTAATATCTCAGTCTTTACCCTCAGCAACGTCCTTAAAGGAAACCTCAACGAACG	O
TILESEMGVVAGISFGVACFULIGI, CAL	
CTCTCTCGTGCCATAACAAATAACCAGTATGAGATAGTGTAACCCAATGTATCTGTGGGCCTATTCCTCTACCTTTAAGGACATTTAG 81	0
GAGAGAGCACGGTATTGTTTATTGGTCATACTCTATCACATTGGGTTACATAGAGACCCGGATAAGGAGAGATGGAAATTCCTGTAAATC	
LSRAITNNOYEIV. CSCG ID NO: 22)	
Z	
GGTCCCCCTGTGAATTAGAAAGTTGCTTGGCTGGAGAACTGACAACACTACTTACT	00
COACCCOCACACTTAATCTTTCAACCAALLGALLICITGALLGITGITGITGAALUACTATCTGCTTTTCAACCAALGACTAACCAACTTAATCTTTCAACCAALLGALLGITGITGITGAACCAACTTAATCTTTCAACCAACTAACAACAACAACAAC	

41 HSBBF0Z	Page 2
CAATCAAGATGTATGTAGACCTAAAACTACACCAATAGGCTGATTCAATCAA	GTA 
GTTAGTTCTACATACATCTGGATTTTGATGTGGTTATCCGACTAAGTTAGTT	
TGTTTGCTATGTTCTAAGTCCACCTTCTATCCCATTCATGTTAGATCGTTGAAACCCTGTATCCCTCTGAAACACTGGAAGAGCTAG	TAA 
ACAAACGATACAAGATTCAGGTGGAAGATAGGGTAAGTACAATCTAGCAACTTTGGGACATAGGGAGACTTTGTGACCTTCTCGATC	
ATTGTAAATGAAGTAAAAAAAAAAAAAAAAAAAAAAAA	

BNSDOCID: <WO\_\_\_9831799A2\_I\_>

HLTAH8O	
CACGAGCATTGCCGCTCTCTCGGTGAGCGCAGCCCCGCTCTCCGGGCCGGGCCTTCGCGGGCCACCGGCGCCATGGGCCAGTGCGGC	ATC + 90
DE LA COMENZA DE	TAG
. M G Q C G	ŧ
	•
ACCTCCTCCAAGACCGTGCTGGTCTTTCTCAACCTCATCTTCTGGGGGGCAGCTGGCATTTTATGCTATGTGGGAGCCTATGTCTTC	ATC 180
TGGAGGAGGTTCTGGCACGACCAGAAAGAGTTGGAGTAGAAGACCCCCCGTCGACCGTAAAATACGATACACCCTCGGATACAGAAG	
	t
T S S K T V L V F L N L I F W G A A G I L C Y V G A Y V F	
ACTTATGATGACTATGACCACTTCTTTGAAGATGTGTACACGCTCATCCCTGCTGTAGTGATCATAGCTGTAGGAGCCCTGCTTTTC	ATC 270
TGAATACTACTGATACTGGTGAAGAAACTTCTACACATGTGCGAGTAGGGACGACATCACTAGTATCGACATCCTCGGGACGAAAAG	
	ſ
	•
ATTGGGC TAATTGGCTGCTGTGCCACAATCCGGGAAAGTCGCTGTGGACTTGCCACGTTTGTCATCATCCTGCTCTTGGTTTTTGTC	ACA 
TAACCCGATTAACCGACGACACGGTGTTAGGCCCTTTCAGCGACACCTGAACGGTGCAAACAGTAGTAGGACGAGAACAAAAACAC	TGT
I G L I G C C A T I R E S R C G L A T F V I I L L L V F V	т
GAAGTTGTTGTAGTGGTTTTGGGATATGTTTACAGAGCAAAGGTGGAAAATGAGGTTGATCGCAGCATTCAGAAAGTGTATAAGACC	<del></del>
CTTCAACAACATCACCAAAACCCTATACAAATGTCTCGTTTCCACCTTTTACTCCAACTAGCGTCGTAAGTCTTTCACATATTCTG	SATG
F V V V V L G Y V Y R A K V E N E V D R S I Q K V Y K T	Y
	CTCC
AATGGAACCAACCCTGATGCTGCCAGCCGGGCTATTGATTATGTACAGAGCAGCTGCATTGTTGTGGGAATTCACAACTACTCAGA	
TTACCTTGGTTGGGACTACGACGATCGGCCCGATAACTAATACATGTCTCTGTCGACGTAACAACACCTTAAGTGTTGATGAGTCT	BALL ,
NGTNP.D. AASRAIDYVOROLHCCGIHNYSD	W
GAAAATACAGATTGGTTCAAAGAAACCAAAAACCAGAGTGTCCCTCTTAGCTGCTGCAGAGAGACTGCCAGCAATTGTAATGGCAG	CCTG
	. 030
CTTTTATGTCTAACCAAGTTTCTTTGGTTTTTGGTCTCACAGGGAGAATCGACGACGTCTCTCTGACGGTCGTTAACATTACCGTC	3373
ENT D'W FKETKNOS V PLSCCRETAS NCNGS	L
GCCCACCCTTCCGACCTCTATGCTGAGGGGTGTGAGGCTCTAGTTGTGAAGAAGCTACAAGAAATCATGATGCATGTGATCTGGGC	CGCA
CGGGTGGGAAGGCTGGAGATACGACTCCCCACACTCCGAGATCAACACTTCTTCGATGTTCTTTAGTACTACGTACACTAGACCCG	
CGG TGGGAAGGC TGGAGATACGAC TCCCCACAC TCCGAGATCAACAC TCTTCGATGTTCTTAGATGTTCTTTAGATGTTCTTAGATGTTTTAGATGTTTTAGATGTTTTAGATGTTTTAGATGTTTTAGATGTTTTAGATGTTTTAGATGTTTTAGATGTTTTAGATGTTTTAGATGTTTTTAGATGTTTTAGATGTTTTTAGATGTTTTTAGATGTTTTTT	
A H P S D L Y A E G C E A L V V K K L Q E I M M H V I W A	. А
CTGGCATTTGCAGCTATTCAGCTGCTGGGCATGCTGTGTGCTTGCATCGTGTTGTGCAGAAGGAGTAGAGATCCTGCTTACGAGCT	CTTC
GACCGTAAACGTCGATAAGTCGACGACCCGTACGACACACGAACGTAGCACAACRCGTCTTCCTCATCTCTAGGACGAATGCTCGA	, 0,0
	c
	, г

																						н	1 }	73	$\mathbb{C}\mathcal{A}$ $\mathcal{E}$	77			
CACGAG	CGC	AGA	GCT	TGG	GGG	. Ť T C	CTT	GGI	rcgc	ACC	CAC	CAC	CCTO	CCI	rgc(	CAC	TGG	TC/	AGCC	TTC	AGG	GAC				cgc	CTG	GTC	90
GTGCTC	GCG	TCT	CGA	ACC	cc	AAG	GA/	ACC.	AGC	TGO	GGT	GGT	3GA(	GG	4CG(	GG T (	GACO	CAG	rcgg	AAC	TCC	CTG	GGA	4C T C	cgic	GCG	GAC	CAG	<b>J U</b>
TCTTTC	CTG	TGG	CCA	GCC	CĄC	JAAC	TGA	4AG(	GCT	rgc	GGC.	ATG	GCG	CGC	GCC.	TGC	стс	CAG	SCCC	TCA	AGT	ACC	TCA	ATGI	TTC	3001	TCA	ACC	180
AGAAAG	GAC	ACC	GGT	CGG	GT	STTO	GAC?	FTC	GCG/	ACG	CCG	TAC	CGC	GCG	c G G	ACG	GAG	GTC(	GGG	AGI	TTCA	TGG	AG.	TAC	AAG(	:GGA	۱AG٦		. • •
												М	Α	R	А	С	L	Q	Α	٧	K	Υ	L	М	F	А	F	N	
														T 0 0	c T C		000	A C A	C 4 C 1		۸.C.C.1	T C C		۸۵۵۱	C T C	TC T .	TCT.	TCCT	
TGTTC																						- 1							270
ACAAGA	AGA	CCG	ACC	CTO	CG.	ACA	CCGC	LAC	GAL	LLA	CAG	LLG	IAG	ACC	GAC	Cuu	CGG	IGI	G 1 C (		, cur		.00	, 60,	GAC.	AGA/	10/1/		
L F	F	W	L	G .	. C	С	G	٧	L	G	٧	G	l	W	L	Α	Α	T	Q	G	S	F	. A	·T	L	S	S	S	
TCCCG	гссс	TGT	CGC	CTO	GCC	AAC	CTG.	стс	ATC	ATC	ACC	GĢC	GCC	TTT	GTC	ATG	GCC	ATC	GGC	TTC	GTG	GC	TGC	CTG	GGŢ	GCC.	ATC.	AAGG	360
AGGGC	AGGG	ACA	GCC	GA	CGG	TTG	GAC	GAG	TAG	TAG	TGG	CCG	CGG	ΑΑΑ	CAG	TAC	CGG	TAG	ccg.	AAG	CAC	ccg.	ACG	GAC	CCA	CGG	TAG	TTCC	000
E P	S	1	S	Α	А	N	L	L	1	1	T	G	А	F	٧	М	А	ı	G	F	٧	G	С	L	G	А	ī	K	
	J	_	J	•												· C T C		C 4 C	000	۸۲۲	ATC.	ררר	ATC	· ר ד ר	. т т с	TTC	פרר	ΤΔΓΔ	
AGAAC																													450
TCTTG	TTC	ACGO	SAG	SAC	GAG	TGA	.AAG	AAG	IGAL	GAL	.GAL	.GAL	, GAL	. <b>L</b> A L	.AAG	G A C	.GAC			100	1 4 3		174	.444	IAAG		-		
E N	K	С	L	L	L	ī	F	F	L	·L	L	L	L	٧	F	L	L	Ε	Α	T	i	Α	I	L	F	F	А	Y	
CGGAC	AAG	<b>4 T</b> T (	GAC	AGG	TAT	GCC	CAG	CAA	GAC	сто	AA(	GAAA	AGGO	TTO	CAC	сто	TAC	GGC	ACG	CAG	GGC	AAC	GTG	GGC	CTC	ACC	AAC	GCCT	540
GCCTG	TTC	TAA	TG	TCC	ATA	CGG	GTC	GTT	CTG	GAC	TTO	CTTI	rcc	SAAC	GTO	GAG	CATO	ccc	TGC	GTC	CCG	TTG	CAC	:ccg	GAC	TGG	TTG	CGGA	
тΩ	K	ī	D	R	Υ	А	a	a	D	L	K	к	G	L	Н	L	Y	G	T	Q	G	N	٧	G	L	T	N	A	
GGAGC											~ ^ T !	CTC	~ ^ ^ ^	~ T A (	~ A C ·	T C A I	TCC	: ŤTſ	10 A C	CTO	: T A C	ΔΔΩ	ect.	CACC	sese	GGTA	ACC T	rgac t	
CCTCG		1																											630
CCTCG	IAG	IAG	616	, 66	il 11	AAU	الماداد	)AL	3ACF	•••	3CA	GAG	3111	3411	313/		3700		_					_	_	.,			
w s	i	I	0	T	D	F	R	С	С	G	٧	S	N	Y	T	0	W	F	Ε	٧	Y	N	А	ī	R	٧	۲	D	
CCTGC	TGC	TŢG	GAG	TTC	AG	rga(	GAGO	TG.	TGG	SCT	GCA	CGC	ccc	CGG	CAC	CTG	GTG	GAA	sgco	3000	TGC	TAC	GA	GAC	GGT	GAAC	3GT0	STGGC	: - 720
GGAC	ACG	AAC	СТС	AAC	STC.	ACTO	STC	GAC	ACC	GA	CGT	GCG	GGG	GCC	GTG	GAC	CAC	CTT	cègo	CGG	CACC	ATO	GCT	CTG	CCA	CTT	CAC	CACCO	ì
s c	С	L	Ε	F	S	Ξ	s	С	G	L	Н	А	Р	G	Т	W	W	K	А	Ρ	С	Y	Ε	Ţ	٧	K	٧	W	
TTCAC						TO T	000	C A T	CTT	***	CCT	CTC	C 4 C	ccc	CCT	сст	GC A	CAT	rr Tí	ace.	ר דו	SACE	CTT	cac	CAT	GACI	CATI	GTAC <sup>1</sup>	ī
AAGT																													
AAGIO	3C 1 C	.116	GAL	.GAL	د ل ل	ALA		J I A	GAA	ALC	CGA	CAL	uiu	CCG	CGA	CCA		C 1 A	<b>.</b>										
L Q	Ε	N	L	L	A	٧	G	!	F	G	L	. с	ī	А	L	٧	a	I	L	G	L	Т	F	Α	. m	T	п	Y	
GCCA																													
CGGT	TCAC	CAG	TTO	CG	TCT	GTG	GAT	GAC	GCG	CAT	ccc	GCG	GGT	GGC	GGC	CGA	AGA	GAC	GGT	TTT	CCT	GCG	GGT	GCC	CCT	CTA	CCG	GCGT	G
сα	v	٧	K	А	D	Т	Y	С	. A		6	Sec	<u>ئ</u> چ	ID	. <i>\</i>	0	.2	4)	)										

45 HTPRA27	Page 2
ACAGC TGC TTTTCCCACCACCAGC TTCGG TG TTCTGCCCCATGC TGGGAGGGAGGGAGGGAGGGACAGG TGCC TGGAGCCCCCGGAA	CC -+ 990
TGTCGACGAAAAGGGTGGTGGAGCCACAAGACGGGGTACGACCTCCTCCCTC	
GTTTCTGGAAGGCCCTAGCTCAGGTGGCTTCAGGGCCTCCGGACCCCCCTGGGAGGGGTGGCCACGTGCTGGCTG	GG -+ 1080
CAAAGACCTTCCGGGATCGAGTCCACCGAAGTCCCGGAGGCCTGGGGGGGACCCTCCCCACCGGTGCACGACCGAC	
GGGGTGGGAGGGGCCTCCAGCACTTTTATATTTACGTATTCTCCAAAGCAGTGTTCACACGGGAGCCAGCC	TC
CCCCACCCTCCCGGAGGTCGTGAAAAATATAAATGCATAAGAGGTTTCGTCACAAGTGTGCCCTCGGTCGG	
GGAAAACAGGTTGGCGCTGGAGGAGCCGGGTCTTGGCATCCTGGAGGTGGCCCCACTGGTECTGGTGCTCCAGGCGGGGCCGTGGA	CC 1260
CCTTTGTCCAACCGCGAEETCCTCGGCCCAGAACCGTAGGACCTCCACCGGGGTGACCAGGACCACGAGGTCCGCCCGGCACCT	
(SEQ II	2:0VIC
TCACCTACATTCCATAGTGGGCCCGTGGGGCTCCTGGTGCATCTTAATAAAGTGTGAGCAGCAAAAAAAA	n 5
AGTGGATGTAAGGTATCACCCGGGCACCCCGAGGACCACGTAGAATTATTTCACACTCGTCGTTTTTTTT	40

									_													ł	AF		20	59	١			
ĢCG	CCG	CCG	GGC	CGC	AGC.	ATG	GGG	CGC	TTC	cgc	GGG	GGC	ÇTG	CGG	TGC	ATC.	AAG	TAC	CTG	CTG	CTT	GGC	TTC	AAC	стд	CTC	TTC	TGGC	TG	90
cgc	GGC	GGC	ĊCG	GCG.	rcg	TAC	ccc	GCG	AAG	GCG	CCC	CCG	GAC	GCC	ACG	TAG	TTC	ATG	GAĊ	GAC	GAA	CCG	AAG	TTG	GAC	GAG	AAG.	ACC		30
						М	G	R	F	R	G	G	Ļ	R	С	i	κ	Y	L	L	L	G	F	N	L	L	F	W	L	
GCT	GGA	TCG	GCC	GTC	ATT	GÇT	ттт	GGA	CTA	TGG	TTT	CGG	TTC	GGA	GGT	GÇC	ΑΤΑ	AAG	GAG	TTA	TCA	TCA	ĢAG	GAC	AAG	TCC	CCA	GAG	TAT	180
CGA	CCT	AGC	cGG	CAG	TAA	CGA	AAA	CCŢ	GAT	ACC	AAA	GCC	AAG	CCT	CCA	CGG	TAT	TTC	стс	AAT	AGT	AGT	стс	CTG	TTC	AGG	GGT	CTC	ATA	100
A	G	s	Α	٧	I	Α	F	G	L	w	F	R	F	G	G	A	I	ĸ	ε	L	s	s	Ε	D	К	s	Ρ	Ε	Y	
TTC	TAT	GTO	GGG	CTG	TAT	GTT	CTG	GTT	GGA	GCC	GGG	GCC	ÇTG	ATO	ATG	GÇC	GTG	GGG	TTC	TTC	GGA	TGC	ŢGC	GGA	GCC	ATG	CGG	GAG	TCĢ	070
AAG	ATA	CAC	CCC	GAC	ATA	CAA	GAC	CAA	CCI	CGG	ccc	CGG	GAC	TAC	TAC	CGG	CAC	ccc	AAG	AAG	CCT	ACG	ACG	CCT	CGG	TAC	GCC	CTC.	AGC	270
F	Υ	٧	G	L	Y	Ÿ.	L	٧	G	А	G	Α	L	M	М	A	٧	G	F	F	G	С	С	G	Α	М	R	Ε	s	
CAA	TGI	GTO	CTT	GGA	TCA	TTT	TTT	ACC	TG	сто	CTG	GTG	ATA	TTI	GCT	GCT	GAA	GTA	ACC	ACT	GG/	AGTA	TTT	GCT	777	· A T A	GGC	AAG	GGG	360
GTT	ACA	CAC	GAA	CCT	AGT	AAA	AAA	TGG	ACC	GGAG	GAC	CAC	TA1	TAA/	ACGA	CĠA	CTT	CAT	TGG	TGA	CC1	CAT	AAA	CGA	ΑΑΑ.	TAT	CCG	TTC	ccc	555
Q	ε	٧	L	G	S	F	F	T	С	L	L	٧	I	F	Α	Α	ε	٧	T	T	G	٧	F	Α	F	ı	G	K	G	
GTA	GCT	ΓΑΤ	ÇGA	CAT	GTT	CAG	ACC	ATG	TA.	TGAA	GAG	GCT	TAC	CAAT	TGA <b>T</b>	TAC	CTT	AAA	GAC	AGG	GGA	AAA	GGC	AAT	GGC	AÇA	CTC	ATC	ACC	450
CAT	CG/	ATAC	GCT	GTA	CAA	GTC	TGG	TAC	AT.	ACTI	стс	CGA	ATO	TT/	ACTA	ATG	GAA	TT	СТС	TCC	CC	ודדו	ccc	TTA	CCC	TGT	GAG	TAG	TGG	730
٧	A	i	R	н	٧	۵	T	М	Y	Ε	ε	A	Y	N	D	Y	L	κ	D	R	G	κ	G	N	G	Т	L	I	T	
TTC	CAC	CTC	AACA	TTT	CAG	TGC	TGT	GGA	AAA,	ĄGA#	AGC	TCC	GAA	ACA	GCTC	CĄA	CCT	ACA	TGC	CCA	AAA	GGAC	CTT	CTA	GG/	AC AC	AAG	AAT	TGÇ	540
AAG	GT	GAG	TTGT	AAA	GTC	ACG	ACA	CCI	TT	TCTI	TCG	AGG	CT.	rg r	CCAG	GTT	GGA	TG	TAC	GGT	TTT	CCTC	GAA	GAT	.001	rgte	TTC	TTA	ACĠ	340
F	н	s	τ.	F	Q	С	С	G	κ	Ε	s	S	Ε	Q	٧	a	P	τ	c	ρ	κ	ε	L	L	G	н	K	N	С	
ATO	GA.	TGA	AATI	GAG	ACC	ΑΤΑ	AATC	AGI	rgt	TAA	CTO	CAC	СТ	CAT	TGGA	AŢŢ	GTO	:GG1	[AT]	GGA	AAT	TGC	AGGT	СТО	ACC	GAŢC	:111	GGC	ATG	620
TAC	CT	ACT	TTAA	CTC	TGG	TAT	ГТАС	TCA	ACA.	ATTO	GAG	GTO	GA	GTA.	ACCT	TAA	CAC	CC/	ATA	VC C 1	ГТА	ACG	rcc.	GAC	:TG	CTAC	SAAA	CCG	TAC	630
i	D	Ε	I	Ε	T	1	I	s	٧	K	L	0	L	I	G	1	٧	G	I	G	Ţ	А	G	L	Ŧ	i	F	G	M	
ATA	TT	CAG	CATO	GTC	СТС	TĢC	TGT	GC	GAT.	ACG/	AAA	TCA	VÇG,	AGA	TGTG	ATA	TG	AAG	CTAC	TTC	TA	CAT	AAA	AATI	rgc/	AAT(	TAA	AGC	TTT	720
TA	AA	GTC	GTA(	CAG	GAG	ACC	GACA	CGC	CTA	TGC	TTT	SAGI	GC	TCT.	ACAC	TAT	AC.	rtc	GAT(	AA(	GAT	GTA	TT	ГТА	₹C G .	TTAC	GATI	TCG		720
ł	F	s	М	٧	L	С	С	А	i	R	N	s	R	0	٧	ſ		(	500	Ţ	L [	1	\C	)'.o	15	)	)			
CA GT			ATG		734	+ (	Sc	Q	J		N	O,	9	$\bigcup$												~				

HAIDO59	
AGTGTTTATGGGACTAAAAAACTTTTAACACCTTTTTAGGGGAAATATTTTGGTCCTATACAAAACATGTAAATATGCTTTATTACTTTC	90
TCACAAATACCCTGATTTTTTGAAAATTGTGGAAAAATCCCCTTTATAAAACCAGGATATGTTTTGTACATTTATACGAAATAATGAAAG	
ATTTTCTGACCCTGCTGTAAACTACTGCAACCCTCACATCCCTCAAAGGGACTTTTATGTCAAACTCTTCTGTTTCTCCAAATATAAGGA	180
TAAAAGACTGGGACGACATTTGATGACGTTGGGAGTGTAGGGGAGTTTCCCTGAAAATACAGTTTGAGAAGACAAAGAGGTTTATATTCCT	
AAAAAGACTAAAGCAAGAGATCTGGCAGTTGAAAATTGTGGGGAAAGAGAATTTGTATGGGCACTGTATCTATGAAATACCTCATACTTAC	270
TTTTTCTGATTTCGTTCTCTAGACCGTCAACTTTTAACACCCTTTCTCTTAAACATACCCGTGACATAGATACTTTATGGAGTATGAATG	
GTTTACATGTTTTCCTAACTTTTTGTATTTTCTTGTATAGCCACCTAGAGAATTCTTCATAGATTAAGAACTACAGTTTTCACCACTTA	360
CAAATGTACAAAAGGATTGAAAAAACATAAAAAGAACATATCGGTGGATCTCTTAAGAAGTATCTAATTCTTGATGTCAAAAGTGGTGAAT	
ACATAAGTAAAACAAAGTCCTTCATAATTTAACCATTAGCATCTTTGGCCAAACCAAAATAAAGAAAAGCATCTTCTCCTAGTTGTGTGT	450
TGTATICATTTGTTTCAGGAAGTATTAAATTGGTAATCGTAGAAACCGGTTTGGTTTTATTTCTTTTCGTAGAAGAGGATCAACACACA	
GGGCAACAGAAACAAGTTAAGGAAACAAAAATACTTATATATA	- 540
CCCGTTGTCTTGTTCAATTCCTTTGTTTTTATGAATATATGTGTCTTGTTTTTATTACAAGAAAAATACGTTTAGGGGACACTTTTA	
AAAATTTTCAATGTTTAAAAAAAAAAAAAAAAAAAAAAA	-54°
TTTTAAAAGTTACAAATTTTTTTTTTTTTTTTTTTT	car.

																									llΓ	T	-14	<b>ሩ ነተ</b> ር	رن	
TTCC	GC	AC GA	AGC.	TGC	GGG	GG.	TGGC	CGG	GC T	GGGC	GGC	ccc	GGG	AG	ccac	GCT	сто	AGI	сто	TCI	AGO	CGC	AGT	ccc	TT(	CGC	CGCT	TTC	GĞ	90
AAGO	CG	TGC	TCG,	ACG	CCC	300	ACCC	GCC	CGA	ccc	3000	GGG	3000	CTC	GCC	CGA	GAG	TCA	GAC	AGA	TCC	CGCC	GTC#	vege.	SAAC	acci	GCGA	AAGO	3CC	90
AGC	CC	TGG	CAG	GGC	CCA	GAAI	GCC/	ATG	GCC	CACT	FATA	AAGA	ACTO	SAG	CAGO	ACG	ACT	GG	TGA	TCA	TCI	rac i	TTGA	AGI	ΑΤ	TTA	CTC.	TTT	GTC	180
TCG	GG.	ACCI	GTC	CCG	GGT	CTT	CGG	TAC	CGG	GTGA	ATA	TTC	rgac	CTC	GTC	TGC	TGA	CCC	ACI	[AG]	AGA	ATGA	AAC	TTCA	ATA	AAT	GAGA	<b>AAA</b> (	CAG	100
								М	Α	н	Y	K	Т	ε	۵	0	D	W	L	I	t	Y	L	Κ	Υ	L	L	F	٧	
						0 T C	000	3 C A /		000	• T.C./	- T.C.	~ C T (	2 T C	GGCA		/	۱.C.C.I	T C (	TO	: A C /	4 A C	ACTI	3 C C 3	τ A C :	CTC	۸۵۲	CTCI	CTC	
															CCG.				+-				+						<del></del> +	270
AAG	116	AAG.	AAG	AAG.	ALL	LAG			LGI	Culi	LAG	GAC	JUAL	JAL	CCG	I A G A	446	1 6 6	340	ACI		, , ,	1 CA		410	GAG	, cu	CAG	GAC.	
F	N	F	F	F	W	٧	G	G	А	Α	٧	L	A	٧	G	i	W	T	L	٧	Ε	K	S	G	Y	١	S	٧	L	
GCC	TCC	AGC	ACC	ттт	GCC	GÇC	TCC	GCC	TAÇ	ATC	стс.	ATC	TTT	GCG	GGC	STAC	TT	GTC.	ATG	GTG.	ACC	GGC	TTC	CTG	GGC	TTC	GGT	GCC.	ATC	360
CGG.	AGG	TCG	TGG	AAA	CGG	CGG	AGG	CGG.	ATG	TAG	GAG	TAG	AAA	CGC	CCG	CATO	3AA	CAG	TAC	CAC	TGG	CCG	AAG	GAC	CCG	A A G	CCA	ĊGG	TAG	500
А	s	S	T	F	Д	А	s	А	Υ	1	L	ı	F	Α	G	٧	L	٧	М	٧	Т	G	F	L	G	F	G	Α	I	
			660		000	T00	C T C	TCC	۸	T A T	T T C	TCC	CTC	TTC	СТС	~ T.C./	A T C :	TTC	ста	CTT	c v c	CTG	стс	מרמ	GG A	GTC	CTG	פרר	СΔТ	
								+					<del></del>	$\rightarrow$	GAG	<del></del>		+	<del></del>		<del></del>		+			<del></del>				450
GAG	ALL	LIL	uct	116	CLG	ALG	IGAG.	AGG	1 44		AAG	ACG	GAC	~~ <b>.</b>	GAG	. AG	1 7 0	AAG	unc	<b>C</b>		<u> </u>			_					
Ĺ	W	Ε	R	к	G	С	L	S	Τ	Y	F	С	L	L	Ļ	٧	i	F	L	٧	E	L	٧	Α	G	٧	٠ ـ	А	н	
0.7.0																														
616	TAT	TAC	ÇAG	AGG	CTG	AGT	GAT	GAA	CTG	AAG	CAG	CAC	TTG	AAC	CGG	ACT!	CTG	GCT	GAG	AAC	TAC	GGG	CAG	ccg	GAG	CAC	GCA	GAT	CAC	540
											-+-		+		CGG	-		+					+			-		+		540
											-+-		+			-		+	CTC		ATG		+			-		+		540
CAC	ATA Y	A T G	GTC Q	TCC R	GAC	TCA S	CTA D	CTT E	GAC L	TTC K	атс	GTG H	AAC L	TTG	GCC R	TGA:	GAC L	CGA A	CTC E	TTG	A T G	ccc G	GTC a	GGC P.	CTC	GTG H	CGT	CTA	GTG H	540
V GCC	ATA Y TCA	ATG Y	GTC Q	TCC R	GAC L	S	CTA D GCAG	CTT E GAT	GAC	TTC K	GTC a TGC	GTG H TGC	AAC L GGA	TTG N AGC	GCC R	TGA	GAC L TCA	CGA A GCC	CTC E GAC	TTG N TGG	ATG Y CAG	G G CAC	GTC Q	GGC P. ACG	CTC E TAC	GTG H	A CCTG	CTA D TTG	GTG H	540 630
V GCC	ATA Y TCA	ATG Y	GTC Q	TCC R	GAC L	S	CTA D GCAG	CTT E GAT	GAC	TTC K	GTC a TGC	GTG H TGC	AAC L GGA	TTG N AGC	GCC R	TGA	GAC L TCA	CGA A GCC	CTC E GAC	TTG N TGG	ATG Y CAG	G G CAC	GTC Q	GGC P. ACG	CTC E TAC	GTG H	A CCTG	CTA D TTG	GTG H	
V GCC	ATA Y TCA	ATG Y	GTC Q	TCC R	GAC L	S	CTA D GCAG	CTT E GAT	GAC	TTC K	GTC a TGC	GTG H TGC	AAC L GGA	TTG N AGC	GCC R	TGA	GAC L TCA	CGA A GCC	CTC E GAC	TTG N TGG	ATG Y CAG	G G CAC	GTC Q	GGC P. ACG	CTC E TAC	GTG H	A CCTG	CTA D TTG	GTG H	
CAC V GCC CGG A GAG	ATA Y TCA AGT	ATG Y GTG CAC V	GAC	R CGA GCT R	GAC L CTC GAG	S CAG	CTA  D  GCAG  GTC  Q	GAT CTA	GAC L TTC AAC	K CAAG GTTC K	GTC Q TGC ACG	GTG H TGC ACG	GGA CCT G	AGC TCG	R AAC TTG N	TGAI	GAC L TCA AGT S	CGA  GCC CGG	GAC CTG	TTG N TGG ACC W	ATG Y CAG GTC G	CCC G CAC GTG H	AGC TCG	GGC P. ACG TGC	E TAC	H	A CCTG GGAC L	D STTG AAC	GTG H GCGG GCC R	
CAC V GCC CGG	ATA Y TCA AGT	ATG Y GTG CAC V	GAC	R CGA GCT R	GAC L CTC GAG	S CAG	CTA  D  GCAG  GTC  Q	GAT CTA	GAC L TTC AAC	K CAAG GTTC K	GTC Q TGC ACG	GTG H TGC ACG	GGA CCT G	AGC TCG	R AAC TTTG	TGAI	GAC L TCA AGT S	CGA  GCC CGG	GAC CTG	TTG N TGG ACC W	ATG Y CAG GTC G	CCC G CAC GTG H	AGC TCG	GGC P. ACG TGC	E TAC	H	A CCTG GGAC L	D STTG AAC	GTG H GCGG GCC R	630
CAC V GCC CGG	ATA Y TCA AGT	ATG Y GTG CAC V	GAC	R CGA GCT R	GAC L CTC GAG	S CAG	CTA  D  GCAG  GTC  Q	GAT CTA D	GAC L TTC AAC	K CAAG GTTC K	GTC Q TGC ACG	GTG H TGC ACG	GGA CCT G	AGC TCG	R AAC TTG N	TGAI	GAC L TCA AGT S	CGA  GCC CGG	GAC CTG	TTG N TGG ACC W	ATG Y CAG GTC G	CCC G CAC GTG H	AGC TCG	GGC P. ACG TGC	E TAC	H	A CCTG GGAC L	D STTG AAC	GTG H GCGG GCC R	630
CAC V GCC CGG A GAG CTC	ATA Y TCA AGT S GCC	Y GTG V CGAG GCTC	GGAC	R CCGA	GAC L CTC GAG L CAG	S CAG	D GCAG CGTC Q GCCCCCGGGG	CTT  E  GAT  CTA  D  GAC	GAC L TTC	K CAAGGTTC K	GTC  GTC  GTC  ACG  ACG  C  GACG	GTG H TGC ACG C GAAG	AAC L GGA CCT G ACA TGT	AGC TCC S	R AAC TTG N GGTG CCAC	T AGC TCG.	GAC L TCA AGT S CGC GCG	GGA A GCC CGG A TGC	GAC CTG  CTG  GGC CCG  GGC	TTG N TGG ACC W CAG GTC	Y CAGGTC	GCC GTG H	GTC  AGC  TCG  S  CCAC	GGC P.ACG TGC T	E TAC ATG	H :ATC	A CCTG GGAC L CATC	D GITG	GTG H GCGG R AAG	630
CAC V GCC CGGG A GAG CTC	ATA Y TCA AGT S GCC CGC A	Y AGTG CAC V CGAG E GGGGA	GGCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCC	R CCGA	GAC  CTC  GAG  L  CAG  GCTC	S CAG	D GCAG	CTT  E  GAT  CTA  D  GAC  CTG	F S S S S G A C	K CAAG	GTC  Q  TGC  ACG  C  GACG  C	H TGC ACG C AAGG TTC K	AAC L GGA CCT G ACA TGT T	N AGC TCG S GTC	R AAC TTTG N GGTG CCAC	T AGC TCG S GCG CGC A CTG	GAC L TCA AGT S CGC GCG CTG	A GCC CGG A TGC C	GAC CTG	TTG N TGG ACC W CAG GTC	Y CAG GTC GCG GCC	GCCGGGAA	GTC  AGC  TCG  S  CAC  GTG	GGC P ACG TGC T CCCC	E TAC ATG	H GATO	A CCTG GGAC L CATC	D STTG	GTG H GCGG R AAG ATTC K CCTG	630
CAC  V  GCC  CGG  A  GAG  CTC  E	ATA Y TCA AGT S GCC CGC A	Y AGTG CAC V CGAG E GGGGA	GGAC  GGGC  GGGC  GGGC  G	R CCGA	GAC  CTC  GAG  L  CAG  GCTC	S CAG	D GCAG CGTC Q GCCC CGGGG P	CTT E GAT CTA D GGAC	F S S S S S S S S S S S S S S S S S S S	K K CTGC K C C C C C C C C C C C C C C C C C C	GTC  GTGC  ACG  C  GACG  C  GACG  C	H TGC ACG C AAGG TTC K	AAC L GGA CCT G ACA TGT T	AGC TCG S GTC CAC	R R R R R R R R R R R R R R R R R R R	TGAI	GAC L TCA AGT S CGC GCG CTG	A GCC CGG A TGC C	GAC; CTG D GGC; CCG	TTG N TGG ACC W CAG GTC Q GGGG	Y CAGGGCC R GCA	CCC G CAC GTG H GCC CGG	AGC + C G G G G G G G G G G G G G G G G G	ACG TGC T CCC GGGG	TAC ATG Y STCC	H GATO	A CCTG GGAC L CATC	D STTG	GTG H GCGG R AAG ATTC K CCTG	630
CAC V GCC CGGG A GAG CTC	ATA Y TCA AGT S GCC CGC A	Y AGTG CAC V CGAG E GGGGA	GGAC  GGGC  GGGC  GGGC  G	R CCGA	GAC  CTC  GAG  L  CAG  GCTC	S CAG	D GCAG CGTC Q GCCC CGGGG P	CTT E GAT CTA D GGAC	F S S S S G A C	K K CTGC K C C C C C C C C C C C C C C C C C C	GTC  Q  TGC  ACG  C  GACG  C	H TGC ACG C AAGG TTC K	AAC L GGA CCT G ACA TGT T	N AGC TCG S GTC	R AAC TTTG N GGTG CCAC	TGAI	GAC L TCA AGT S CGC GCG CTG	CGA A GCC CGG A TGC C CCTI	GAC; CTG D GGC; CCG	TTG N TGG ACC W CAG GTC	Y CAGGGCC R GCA	GCCGGGAA	AGC + C G G G G G G G G G G G G G G G G G	ACG TGC T CCC GGGG	E TAC ATG	H GATO	A CCTG GGAC L CATC	D STTG	GTG H GCGG R AAG ATTC K CCTG	630
CAC V GCC CGG A GAG CTC E GTG CAC V	Y TCA AGT S GGCC CGC A GGAC CCTC	Y GGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGG	GGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGG	R CCGC R CCGC R CCGC CCGC CCGC CCGC CCGC	GAC  GAG  CAG  GGTC  GGGAC  L  GGGTC	S S S S S S S S S S S S S S S S S S S	D GCAG  GGAG  GGCCC  GGGG  P  CAAG  K  CACC	GAT CTA D GAC CTG CTG CTG CTG CTG CTG CTG CTG CTG CT	F S S S S S S S S S S S S S S S S S S S	K K CTGC C C C C C C C C C C C C C C C C C	GTC  Q  TGC  ACG  C  GACG  C  GACG  F	H TGC ACG C AAAG TTC K CCTG GGAC	GGA CCT GACA TGT T	AGC TCG S GTC CAC V GGAC GCTC	R AAC TTTG N GGTG CCAC V CCAC H CCAC	T AGC T CG. S GCGC CGC A CTG	GAC L TCA AGT S CGC GCG R CTG GAC L CCAT	GGAA	GAC+CCG  GGC+CCG  GGC+CCG  GGC+CCG  GTAC	TTGG N TGG ACC W CAG GTC Q GGGG CCC G	Y CAGGGCC R GCA	GCCGGGAA	AGCCA	GGC P ACG TGC T CCC GGGG	TACO ATG	H STACE N N SGT(CA)	A CCTG GGAC L CATC GGGCC A TCCA	CTA  D  STTG  AAC  L  TAT  Y  C  C  C  AAC  AAC  T  C  C  C  C  C  C  C  C  C  C  C  C	GTG H GCGG GCC R AAG ATTC K CCTG L	720 810
CAC V GCC CGG A GAG CTC E GTG CAC V	Y TCA AGT S GGCC CGC A GGAC CCTC	Y GGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGG	GGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGG	R CCGC R CCGC R CCGC CCGC CCGC CCGC CCGC	GAC  GAG  CAG  GGTC  GGGAC  L  GGGTC	S S S S S S S S S S S S S S S S S S S	D GCAG  GGAG  GGCCC  GGGG  P  CAAG  K  CACC	GAT CTA D GAC CTG CTG CTG CTG CTG CTG CTG CTG CTG CT	F S S S S S S S S S S S S S S S S S S S	K K CTGC C C C C C C C C C C C C C C C C C	GTC  Q  TGC  ACG  C  GACG  C  GACG  F	H TGC ACG C AAAG TTC K CCTG GGAC	GGA CCT GACA TGT T	AGC TCG S GTC CAC V GGAC GCTC	R AAC TTTG N GGTG CCAC V CCAC H CCAC	T AGC T CG. S GCGC CGC A CTG	GAC L TCA AGT S CGC GCG R CTG GAC L CCAT	GGAA	GAC+CCG  GGC+CCG  GGC+CCG  GGC+CCG  GTAC	TTGG N TGG ACC W CAG GTC Q GGGG CCC G	Y CAGGGCC R GCA	GCCGGGAA	AGCCA	GGC P ACG TGC T CCC GGGG	TACO ATG	H STACE N N SGT(CA)	A CCTG GGAC L CATC GGGCC A TCCA	CTA  D  STTG  AAC  L  TAT  Y  C  C  C  AAC  AAC  T  C  C  C  C  C  C  C  C  C  C  C  C	GCGG R AAG	720 810

CCTCAAGACAACATGTGGCACATGCCATCTGCAAGG

GGAGTTCTGTTGTACACCGTGTACGGTAGACGTTCC

936 (SCQ IDNO: 11)

BNSDOCID: <WO\_\_\_9831799A2\_I\_>

																									<u>احا</u>	<u> </u>	<u>, V )</u>	<u>४प</u>		
AGC 1	TA	CTT	TCA	стс	ACC	GCC	TGT	CCT	TCC	TGA	CAC	CTC	ACC	ATG	TGT	AÇG	GGA	AAA	TGT	GCC	GC	TGT	GTG	GGG	тс	TCC	CTC	ATTA		90
TCGA	AAT	GAA	AGT	GAG	TGG	CGG	ACA	GGA	AGG	ACT	GTG	GAG	TGG	TAC	ACA	TĠC	ССТ	TTT	ACA	CGG	GCG.	ACA	CAC	ccci	3AG.	AGG	GAG.	TAA		-
														М	С	T	G	κ	С	Α	R	С	٧	G	L	S	L	I	Т	
CTC.		c T C	0.7.0	T.C.C	4 T T	CTC		4 A C		· C T C	CTC	CTC	CTA		- A A T		0.40	· ^ C C	TCC	TOO	۸۲۲	4 A C	۸۲۲	A A C:	C 4 T	CIC	A.C.C.	TTC	~ ^ ^	
GAG			!			<del></del>			+		-+-		+			<del></del>			+	<del></del> -			+			<del></del>		<del></del>	<del>+</del>	180
GAG	ACG	GAG	LAG	ALG	IAA	LAL	CGG	116	Lui	UAU	GAL	GAL	LAI	G G F	A I I A			. 166	АСС	ALL	166	116	166	116	۱Α۱ د	GAG	166	AACI	וונ	
L	С	L	٧	С	I	٧	Α	N	Α	L	L	L	٧	Ρ	N	G	ε	T	S	W	T	N	T	N	Н	L	S	L	a	
GTC:	TGG	стс	ATG	GGC	GGC	TŢC	ATT	GGC	GGG	GGC	СТА	ATG	GT/	ACTO	STGT	CÇA	GGG	ATT	GCĄ	GCC.	GTT	CGG	GCA	GGG	GGC	AAG	GGC	TGC		270
CAG	ACC	GAG	TAC	CCG	CCG	AAG	TAA	CCG	CCC	ccc	GAT	TAC	CAT	r G A C	CACA	GGT	ccc	TAA	CGT	CGG	CAA	GCC	CGT	ССС	CCG	TTC	CCG.	ACG.		270
v	w	L	М	G	G	۶	ı	G	G	G	L	М	٧	L	С	Р	G	ı	Α	Α	٧	R	A	G	G	K	G	С	ε	
·		-		_	-																									
			<del>! -</del>	+		<del>-+-</del>		<del></del>	<del></del> i		-+-		+		TTC	-+-			!				+	+		-+-				
CCA	CGA	CCC	ACG	ACA	CCT	TTG	GCG	ACG	TCC	TAC	GAC	GCG	AGC	CA	GAAG	AGG	AGC	CGC	AAG	יייי	CAC	GAA	CCA	CGG	TAG	AIG	ALG	Ŀ		
G	A	G	С	С	G	N	R	С	R	М	L	R	S	٧	F	S	S	Α	F	G	٧	L	G	Α	į	Υ	С	L		
GTG	TCT	GGA	GCT	GGG	стс	CGA	AAT	GGA	ccc	CAGA	TGC	TTA	AT	GAAG	CGGC	GAC	STGO	GGG	TAC	CAC	TTC	GAA	ĢAC	ACC	GCG	GGA	GC T	TAC	TTG	"60
CAC	AGA	ĊCT	CGA	CCC	GAC	GC 1	TTA	CCT	GGC	STCT	ACG	AAT	TA	CTT	GCCC	CTO	CAC	ccc	ATO	GTG	AAG	CTT	CTG	TGG	CGC	CCT	CGA	ATG	AAC	450
٧	s	G	Α	G	L	R	N+	G	Р	R	С	L	М	N	G	Ε	W	G	Υ	Н	F	Ε	D	Т	Α	G	А	Υ	L	
														20 T	,		- + o (				C T C	T T C	Ŧ C C	CTO	CTC		• • • •		TCC	
			+					+					+		GGTC	<del></del>		<del></del>					+	+				+		540
GAG	116	1666	1166	(GA)	ACC		4666	ACG	1616		. 444	1 G G F	460	G C A	CCAC	uu	3ACI	-111	16 A C	, 100	GAG	AAG	AGC	GAC	GAC	CAC		-		
L	N	Ŗ	Τ	L	W	D	R	С	Ε	Α	ρ	Ρ	R	٧	٧	ρ	W	N	٧	T	L	F	S	L	L	٧	Α	Α	S	
TGC	CTG	GAG	ATA	GT/	ACTO	TGT	rggg	ATC	CAC	GCTG	GTG	SAAC	GC	GAC	CATI	GG.	TGT	CTTO	CTG	GGC	GAT	TGC	AGG	AAA	AAA	CAC	GAC	ACA	CCT	630
ACG	GAC	СТС	TAI	CAT	rgad	CACA	ACCC	TAG	GT	CGAC	CAC	TTC	SCG	CTG	GTA	ACC/	ACA	GAA(	GAC	CCG	CTA	ACG	TCC	TTT	TTT	GTC	CTG	TGT	GGÀ	000
С	L.	E	1	٧	L	С	G	I	a	L	٧	N	A	T	ı	G	٧	F	С	G	D	С	R	Κ	K	Q	D	T	Ρ	
a											-001			<b>.</b>	0.4.07	. c T	~ . ~ .	T.C.C.	<b>.</b>				A T A	A A C	TOC		tere	CTC	TCA	
			<del></del>			<del></del>		<del></del>		+					GAC(	<del></del>		<del></del>					+							720
GIG											4662	4667	AAG	GAC	Liu	. GAI	روا ا دا ا	AGG	JAA!	JGAU	CGA	(161	IAI	116	ACC	1447	1000		741	
Н	٠ (	S	eC	J		N	0:	$\mathcal{A}$	. +	)																				
AAA	AAA	AAA	AAAA	AAA	AAA	2 4,	20	/<	-	G) .	<b>T</b> i	$\supset i$	\[(	ኅ .	12	)	ı													
TTT	TTI	TTT	TTT:	TTT	TTT(	- /. G	30	ٺ	ہدر	. ب	いし	ノ I	10	·	100	- )	,													

																								<u>H (</u>	<u>۷۷</u>	$\mathcal{B}_{\mathcal{I}}$	88	0_		
C.	AC G	AGA	GAT	TGT	CGG	CTG	CGG	GTA	TAT	TCC	AATI	rcc	CG.	CTO	сто	ATO	SAAT	FATG	AAC	TGA	AGO	GCT	CTO	GAC	сто	GA.	GTO	GGTT	TCT	90
G	rgc	TCT	CTA.	ACA	GCC	GAC	GCC	CAT	ATA	AGG	TTA	4GG(	GGC.	AGA	GGAC	TAC	:TT/	ATAC	TTO	ACT	TC	CGA	AGA	CTG	GGAC	CT	CAC	CCAA	AGA	
٩G	CAG	GGC.	AAA.	ATG	GGG	TCT	CGG	AAG	TGT	GGA	GGC.	TGC	ÇTA.	AGT	TGT	TTG	CTG/	ATTO	CĠ	7.70	CAC	CTT	rgg.	AGT.	ATAA	ATC(	STG	AACA	ATA	180
TC	GTC	CCG	TTT	TAC	CCC	AGA	GCC	TTC	ACA	CCT	CCG	ACG	GAT	TCA	ACA	AAC(	GAC.	TAAC	GC	GAAC	CGT	GAA	ACC	TCA	TAT	TAG	CAC	TTGT	TAT	
				м	G	S	R	κ	С	G	G	С	L	s	С	L	L	į	Р	٤	Α	L	W	\$	ı	ſ	٧	N	ĺ	
ТΛ	TTG	ΤΔΤ	TTC	ררה	ΔΔΤ	ດດດ	CAA	ACT	TCC	TAT	GCA	TCC	AGC	AAT	ΔΑΑΙ	CTC.	ACC	AACI	TAC	GTG:	TGG	TAT	TTT	GAA	GGA.	ATC	TGT	TTC	TCA	
			·										+					+										AAG	<del></del>	270
			_			•	•	<b>.</b>	c	v	٨	c	c	·N	v	1	т	N	Y	V	W	Υ	F	F	G	ı	С	F	s	
L	L	Y	۲	۲	N	٦	u		,Σ	i	А	3	3		^	_	•	.,	•	•			•	_	J	•	_	•		
									;				+					<del></del>					-					AGC.		360
CG	TAG	TAC	TAC	GAA	TAT	CAT	TGT	TGT	CAA	GAA	GAC	CAT	GAC	CTC	TTA	TTA	TTG	TTG	ATA	TTT.	ACA	ACG	GTC	TCA	CTT	TTG	ACG	TCG	TTT	
G	ı	М	М	L	i	٧	Ţ	T	٠٧	L	Ļ	٧	L	Ε	N	N	N	N	Y	K	¢	С	Q	S	ε	N	С	s	K	
ΔΔ	TAT	GTG	ACA	CTG	CTG	TCA	AATI	ATO	: 11	тст	TCC	стс	GGA	ATT	GCT	TTT	TCT	GGA	TAC	TGC	CTG	GTC	ĄTC	TCT	GCC	TŢĠ	GGT	стт	GTÇ	
																<del></del>		_			- 1		<del></del>					GAA		450
		.,				c		ī	_	c	ç	ı	G	ı	Δ	F	S	G	Υ	c	L	٧	i	S	А	L	G	L	٧	
K	T	٧	'	L	Ĺ	3	•	ı	r	3	3	_	G	•	^	•	Ū	J			_									
			1			:_				<b></b>																		TGG		54
371	ccc	:GG1	ATA	ACC	GCC	TG	GGAA	ACTA	4000	GACC	стс	ATA	CGA	AAA	CTT	CCG	iTGA	CGA	CCT	GCA	AAG	GAA	TGI	CIA	AAGA	1166	HAI	TACC	IAA	-
٥	G	Ρ	Y	C	R	T	L	0	G	W	Ε	Y	Α	F	Ε	G	T	Α	G	R	F	L	T	0	S	S	Į	W	i	
CAC	TGC	сто	GAA	ACC.	rgc.A	ACA'	TGT	TGT	GGA	GTGC	SAAC	ATO	ATI	TTA	<b>.</b> T T T	TCC	ATT	стс	ATA	ACC	стс	AGT	GGG	CT	ГСАА	kg T G	ATC	CATC	TGÇ	63/
													_					<del></del>										GTAG		Ų.J.
•	_	,	_		٨	ы	v	v	_	W	N	1		ı	F	S	i	E.	į	T	L	S	G	L	a	٧	1	I	С	
u	L	L	E	٢	А	П	٧	٧	-		.,	•		_	•	_	•	_	•		_					= .				
٠.			4 .							4									_				-					GACA		رے ر
GAI	STA	GTC.	TCA	TCA	GTA	GT	TGA	TAG	GTT	CTA	TGAC	CAC	ACC.	TTC	SATA	AGI	CAC	CTAG	AA(	GTC	:GG/	ACCI	TA	G I A				CTGT		
L	ī	R	٧	٧	М	Q	L	s	ĸ	1	ŗ	С	G	S	Y	S	٧	I	F	۵	ρ	G	i	i	. (	<u> S</u>	20	IJ	) N	vO.
TG	TTT	TCC.	ΑΤΤ.	ATC	AAG	ACA	TGG	CCA	TCT	ATC	TAA	ATA'	TTA	TAT	CAAC	TG.	TGT	TAGA	CT.	rgad	GGG	CAAl	ΓΑΤ	TGA.	AAA	TGA:	rgg:	TGC	гттс	:
																												ACGA		
.,•					•	,												•												
TG	CAT	TTG	GTG	TTT	ATT	TGT	AAA	AAA	TTT	GCA	GTC	CTC.	A C T	GCA	CAT	GCA.	AGT.	ATAC	CAI	cc.	TTC	CAT	TTA	GTA	TGT	TTŢ	TTA	AGT	AATA	- 90
																												TCA		

HINPRSO

AGAAACCTTTATGCAATATGTATATTGCAACATTATTTAATATTCTGGAAAATTGGAAACACCCCAAAATTCTAACTCAAA

TCTTTGGAAATACGTTATACATATAACGTTGTAATAAAATTATAAGACCTTTTAACCTTTGTGGGGTTTTAAGATTGAGTTT

1071

																			_ `									
TTGTGGAG	GGC/	AGCA	GAG	AG	FACC	CA	GCT	GGA	ATO	CT	TTC	CTG	CTG.	A T G	AGC	CCA	GGC	TG	GAG	GTG	CC	TGC	TCA	CAT	GCT	CTT	ccc	90
AACACCTO	CCG	rcgr	CTO	:TC/	ATGO	3G T	CGA	CCT	CTAC	GGA.	AAG	GAC	GAC	TAC	TCG	GGGT	ccc	GAC	CTC	CAC	GGG.	ACG.	AGT	GTA	CGA	GAAC	GGG	30
														M	s	ρ	R	L	Ε	٧	P	С	S	н	Α	L	ρ	
													<b>.</b>	<b></b>	040			- A T	. T. T.	A C T	ctc	۸۰۲	CIC	۸۵۲	- C A C	CACI	r c t	
AGGGTCTC							<del></del>				<del></del>						<del>-</del>				+	$\overline{}$						180
3TCCCAGA0	AGC	GGAC	CCC	اکاند	_AG	IAG	IAI	LAII	المالما	J ( )	GAL	LAG	AAL	ull	CIC	666	100	31 A	MAA	IUA	LAC	100	GAC	166		310	CGA	
Q G L	S	P	G	Q	٧	Ī	I	٧	R	G	L	٧	L	a	Ε	Ρ	Ķ	Н	F	Ţ	٧	S	L	R	O	a	Α	
GCCCATGCT	CCT	GTGA	CAC	CTC.	AGG	GCC	TCC	TTC	GCA	GAC	AGA	ACT	CTG	GCC	TGG	ATC1	rcç	cgc	TGG	GGG	CAG	AAG	AAA	CTO	SATO	TCA	GCC	270
GGGTACGA	GGA	CACT	GTO	GAG	TCC	CGG	AGG	AAG	CGT	CTG	TCT	TGA	GAC	CGG	ACC	TAGA	4GG	GCG	ACC	CCC	GTC	TTC	TTT	GAC	CTAG	AGT	cgg	210
а н а	P	v	т	l_	R	Α	s	F	Α	D	R	T	L	Α	W	í	s	R	W	G.	Q	K	K	L	I	s	Α	
	·																								C C A C		CTC	
сссттсст											<del></del>	+				+		_			<del></del>							360
GGGAAGGA	SAAA	ATGG	GG																		CGC	GAG	114		J G 1 C		GAC	
PFL	F	Y	Ρ	Q	R	F	F	Ε	٧	L	L	L	F	Q	Ε	G	G	L	K	L	Α	L	N	G	a	G	L	
GGGGCCAC	CAGC	ATGA	AC	CAG	CAG	GCC	ст	GAG	CAG	ста	CGG	GAG	сто	CGG	SATC	AGT	GGA	AGT	GTC	CAG	ÇTC	TAC	TGT	GT	CCAC	TCC	TGA	450
CCCCGGTG	STCG	TACT	TG	GTC	GTC	CGG	GAC	стс	GTC	GAC	GCC	CTC	GAG	GCC	TAG	TCA	CCT	TCA	CAG	GTC	GAG	ATG	ACA	LC A	GGT	AGG		
G A T	s	М	N	a	a	Α	L	Ε	Q	L	R	Ε	L	R	ı	s	G	s	٧	Q	L	Y	С	٧	н	s	٠.	NO: -
AGGATGGT		224		• • •		C 4 /			ACT	C A C		\ C T C	יררו	٠ ۷ ٢ ٢	ירר	. C C V	רדר	trr	Trr	ררז		ΤΔΔ	ACC	` A T	CCAI	ссте	SAAC	
TCCTACCA						+								-+		<del></del>	+							-+				540
TUUTALLA.	AGG	CCI	IIA	166	CGI	C 1 1		1170	., .,			un.	Juu				٥٨٥											
ACCAGCAC	ATCA	.GGG(	CCT	GGT	TCA	CC.	тсто	GGG	TCA	CGA	AGAC	TGA	AGTO	TAC	CAGO	AGC	TTT	GGG	CCT	GAC	GG/	AGG	CAC	CAA	GAG	TGCA	AAAG	
TGGTCGTG														-		<del></del>								$\overline{}$				030
																•												
GTTCCTCG	AACT	CTG	CAC	CTT	сст	CC	ACCA	AGGA	GCC	TG	GGA	TATO	GGC.	rcç	ATC 1	rgcc	TTC	AGG	GCC	TG	GAC	rgc/	ACTO	CAC	AGA	GGC#	AAGT	720
CAAGGAGC	TTGA	GAC	GTG	GAA	GGA	GG.	TGG	CCI	CGG	AC	CCTA	ATA	CCG	٩GG	TAGA	CGG	AAC	TCC	CGC	ACC	TG	ACG1	ΓGΑ	G T G	TCT	CCGT	TTCA	
GTTGTAGA	CTAA	CAA	AGA	TAC	:TCC	:AA	AAT	ACAA	TGG	CT	TAA	AGA	ATG	TGG	TCA	ГТТА	TTC	TT	TAT 1	TAT	TTA	111	ATT	TGT	GGT	CAAA	4 T A A	810
CAACATCT	GATI	GIT	TCT	ATO	AGG	TT	TTA	TGT	TACC	GA	ATT	TCT	TAC.	ACC.	AGT	AAAT	AAC	GAAA	ATA	ATA,	AAT.	AAA'	TAA	ACA	CCA	GTT	TATT	
																									_			
ATAAATAA	GGTT	TATT	TAT	TTA	AAA	AAA.	AAA.	AAA	AAA	AAA	AAA.	AAA	AAA	AAA	AAA	A <i>R</i>	165	/<	GP.	D .	CD	<u>N</u>	0:	1/	4.	)		
TATTTATT	CCA	ATAA	ATA	AAA	rtti	гтт	TTT	TTT	FTT	TTT	TTT	TTT	TTT	TTŤ	TTT	TT		<i>ـ</i> ــ	~~	4 0		ų J				-		

HROAD63

Päge 1

						-+		,					<del> </del>			-+-	CTG		<del></del>				<del>†                                      </del>	+						90
CGT	GCT	СТС	TGC	TGT	FAGT	CTC	TAC	TCC	TGT	CGT	AAC	GAC	GAG	GAA	CGT	CGG	GAC	CGA	CAC	CGA	TGT	CCC	GGT	CGG	GAA	TGG	GAC	GCG	ACG	
							м	R	Т	Α	L	L	L	L	Α	A	L	A	٧	Α	T	G	Ρ	Α	L	T	L	R	С	
			-			-+-							+	+			CAGC						+			<del>- ; -</del>				18
GTO	CAC	ACG	TGG	TC	GAGG	TCG	TTG	ACG	TTC	GTA	AGA	CAC	ĊAG	ACG	GGC	CGC	STCG	AGA	GCG	AAG	ACG	TTC	TGG	TGC	TTG	TGT	'C A C	CTC	GGA	
н	٧	С	T	S	S	s	N	С	K	Н	S	٧	٧	С	. <b>P</b>	Α	s	\$	R	F	С	K	T	T	N	T	٧	Ε	P	
ст	AG	GCT	TCC	CC	GAAA	GŢC	TGG	GAC	CAG	GTO	CAG	GTG	GGC	ATO	GAA	TG	CTGA	TGA	CTT	GGA	GCA	GGC	ccc	ACA	GAC	ccc	:ACA	GAG	GAT	27
GAG	TC	CCGA	AGG	GG	CTTT	CAG	ACC	CTG	GTO	CAC	GTC	CAC	CCG	TAC	CTT	AC	GACT	ACŤ	GAA	CÇT	CGT	CCG	GGG	TGI	CTG	GGG	STGT	CTC	CTA	
L	R	A	S	P	K	٧	W	D	Q	٧	Q	٧	G	М	ε	С	•	(3	50	$\bigcirc$	IĮ	1 (	40	);	3C		)			
GA	4GC	CAC	:cc/	CA	GAGG	ATG	CAG	ccc	CCA	AGC.	rgc A	TGG	AAC	GTO	GAG	GA	CAGA	AGC	CCT	GTO	GAT	ccc	CGC	ATI	TTCA	CAC	:TC	: T T C	TGŢ	3
CT.	rcg	GTG	GG	rgr	стсс	TAC	GTC	GGG	GG	rc G/	ACGT	TACC	TTC	CAC	CTC	CT	GTCT	TCG	igg.	ACAC	CTA	GGG	GCC	TAA	AAGT	GTO	SAGO	GAAG	SACA	اد
TT	TGT	TGC	:GT	ГТА	TTTT	TGI	ACT	CAA	ATO	CTC.	TACA	ATGG	AGA	ATA/	AATG	ΑŢ	TTAA	ACC	AG	ΓΑΑΑ	AAA	AAA	AAA.	AAA	AAAA	AAA	Δ (	<u>.</u> S	Q	J
			-		+	+		~	- T A (	240	A T C I		70	T A T	T A C	- <del>-  </del> -	AATT	TCC	TC	\ T T '	TTT		77	TTT	TTTI	TT.		41	NO	٦,

HMW6546

CONTINUENCE CONTIN	TGTTC ACAAC L L TCGG1 AGCCA	GAGGGC1	GAG GAG GAG	GAT CTA GGG CCC G	CAG GTC CCG GGC A	TGT GCG GG	M GCCC	GGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGG	GAGA GTCT GCCT GGGA A	N FACE	L GGT(	AAGO K GTGG CACO	D CGCG R	AACC	GCG(	GGAC CCTC G	CGGCC GCCC R	TGC GACG L	GGC P	GGC A	GGC G G	GGG GGC	GGG CCC R	GCA CGT G	TGG ACC M	GCA CG1 G	TTC1	GGG A	TG/ AC	AAG TTC K
HARDON L K D L A G R L P A G P R G M G T A L K  TGTTGCTGGGGGCCGGGGGCCTACGGGTCTGCGGGAATCTGTTTACCGGGGAATCTGTTCACATCGG ACAACGACCCCCGGCCGGGCACCGGATGCCACACGGCGTTAGACCACCGGGAATCTGTTTCACTGGGAAGGGGGGCAAGAGCCATCTTCTTCAATCGG ACAACGACCCCCGGCCGGGCACCGGATGCCACACGGCCTTAGACACAAGTGGCACCTTCCGCCGGTGTTCTGGTAGAAGAAGAAGTAGAC L L L G A G A V A Y G V R E S V F T V E G G H R A I F F N R  TCGGTGGAGTGCAGCAGGACACTATCCTGGCCGAGGGCCTTCACTTCACTTCAGGTTCCAGTTCCAGTACCCCATTATCTATGACATTCGG AGCCACCTCACGTCGTCCTGTGATAGGACCGGCCTCCCGGAAGTGAAGACCTTCAGGTTCCAGGTTCCAGTTCCAGTTCCAGTTCCAGTACCCCATTATCTATGACATTCGG AGCCACCTCACGTCGTCCTGTGATAGGACCGGCCCCCGGAAGTGAAGAGTCCCTTGGGTAACACAGGTCACAACAAGAACTTCTCTCGCCCGAAGAAAATCTCCCTCC	TGTTC ACAAC L L TCGG1 AGCCA	GAG GC1 CG/ L TGC ACC	reger ACC	GGG CCC G	GTC CCG GGC A	GCG GG	M GCCG GGG	A GTGC	GTCT GCCT GGGA A	TACO Y	L GGT( CCA)	TTCC K GTGC CACC	D CGCC GCGC	L GAA1	A CCT	G G GTG1	R TTC	SACG L ACCG	GGC P	GGC A	ccc G	GGG P GGC	CCC R	CGT G GAG	ACC M CCA	G G	T	GGG A TCA	AC	K CGG
H A Q N L K D L A G R L P A G P R G M G T A L K  TGTTGCTGGGGGCCGGGGCCCTGGGCCTACGGTGTGCGCGAATCTGTTTACCGGGGAAGGCGGGCAAGAGCCATCTTCTTCAATCGG ACAACGACCCCCGGGCGGCCCGGGCCCCGGATGCCACACGGGCTTAGACACAAGTGGCACCTTCCGCCCGTGTCTCGGTAGAAGAAGAAGTTAGCC  L L G A G A V A Y G V R E S V F T V E G G H R A I F F N R  TCGGTGGAGTGCAAGAAGACACTATCCTGGGCCGAAGGCCTTCACTTCAGGATCCCTTGGTTCCAGTACCCCATTATCTATGACATTCGG AGCCACCTCACGTCGTCCTGTGATAGGACCGGCTCCCGGAAGTGAAGTCCTAGGGAACCAAGGTCATGGGGTAATAGATACTTAGACATTCGG AGCCACCTCACGTCGTCCTGTGATAGGACCGGCTCCCGAAGAGTCCAAGAGTCCTAGGGAACCAAGGTCATGGGGTAATAGATACTTATGAAGAC  I G G V Q Q Q D T I L A E G L H F R I P W F Q Y P I I Y D I R  TCCAGACCTCCGAAAAATCTCCTCCCCTACAGGCTCCCAAAGACCTACAGATGGTGAATATCTCCCCTGCGGAGTGTTGCTCTGACCCAATGCT  A R P R K I S S P T G S K D L Q H V N I S L R V L S R P N A  TAGGAGCTTCCTAGCATGACAAGGGGGATCCCGAAGTTACGAGGAACCAAGGGTGTTGCCCTCCAACAAGAGCTGGGTTACGA  TCCCTCGAAGGATCGTACATGGTCGGGGATCCCGAACTGATGATGCTCCTTGCTCACAACGAGGTGTTGCCCAACAAGAGCTGAGAGTGTG TCCCTCGAAGGATCGTACATGGTCGGGATCCCGACCTGATGCTCTTGCCAACAGGGGAGCAACAAGAGGTGCTCAACAAGAGGTGCTCAACAGGGAGCTCAAAGAGTGTGCCCCAAGAGTGTTCTCACCA  G E L P S H Y Q R L G L Q Y E E R V L P S I V N E V L K S V  TCGGCCAAGTTCAAGTTCACAGCGCGATCCCCAACGAGGTCCCAAGAGGCCCAAGGAGAGCCCAAAGAGGCCCAAAGAGGCCCAAGGAGG	TGTTC ACAAC L L TCGG1 AGCCA	GC1 CGA L TG0 AC0	IGGG	GGG CCC G	GGC A	GCG CGC G	M GCCC GGC A	A GTGC CACC V	GCCT	N TACO Y Y	G G G	K GTG( CAC(	D CGC( GCG( R	L GAA1	A	G GTG1	R TTC	L	P TGG	A A A G	G GCG	P GGC	R ACA	G GAG	M CCA	G	T TTC1	TCA	L AT	K CGG
TGTTGCTGGGGGCCGGGGCCCTGGCCTACGGTGTGCGCGAATCTGTGTTCACCGTGGAAGGCGGGCACAGAGCCATCTTCTTCAATCGG ACAACGACCCCCGGGCGGGCACCGGATGCCACAGGGCTTAGACACAAGTGGCACCTTCCGCCCGTGTCTCGGTAGAAAAAATTAGCC L L G A G A V A Y G V R E S V F T V E G G H R A I F F N R TCGGTGGAGTGCAGCAGGACACTATCCTGGCCGAGGGCCTTCACTTCAGGATCCCTTGGTTCCAGTACCCCATTATCTATGACATTCGG AGCCACCTCACGTCGTCGTGATAGGACCGGCTCCCGGAAGTGAAGTCCTTAGGGAACCAAAGGTCATGGGGTAATAGATACTGTAAGCC I G G V Q Q Q D T I L A E G L H F R I P W F Q Y P I I Y D I R ICCAGACCTCCGAAAAATCTCCTCCCCCTACAGGCTCCCAAAAGACCTACAGATGGTGAATATCTCCCTGGGAGGTGTTGTCTCGACCCAATGCT GGTTCTGGAGCTTTTTAGAGGAAGGGGGATGTCCGAGGTTTCTGGATGTCTACCACTTATAGAGGGACGCTCACAACAGAGGTGTTGCCGACCCAATGCT A R P R K I S S P T G S K D L Q H V N I S L R V L S R P N A ACAGGAGCTTCCTAGCATGTACCAAGGCCTAGGGCTGGACTACGAAGGGAACGAGGTTGCCGTCCATTGTCAACGAGGTGCTCAACAAGGGTGTTCACAC GG E L P S H Y Q R L G L D Y E E R V L P S I V N E V L K S V ATGGCCCAAGTTCAAAGTCCCCAAGGGGGGGCCCAGGGGCCCCAGGGTACCAACGGCAGGAACAATGGGCCCCAAGGAC ACCCGGTTCAAGTTACGGAGGTGCCCAAGGGGGCCCAGGGGCCCCAGGGTACCAACGGCAGGCCCTCAACCAAGGGCCCAAGGAC ACCCGGTTCAAAGTCCCCAAGCTGATCACCCAGCGGGCCCCAGGGGCCCAAGGAACCAAGGCGCCCCTGAACCAAGGGCCCAAGGACCAAGGCCCAAGGACC ACCCGGTTCAAGTTACGGAGTGTCGACCAACGAGGGGCCCAAGGGGCCCAAGGGACCAACCA	ACAAC L L TCGG1 AGCCA	CGA L TG0 AC0	GAG	CCC G TGC	GGC A	G G	A GACA	V ACT/	GCCT GGGA A	Y TG(	G G	GTGG CACG	GCG(			GTG						+	+				+			cgg
ACACAGGACCICCAGGAGGACACTATCCTGGCCGAAGGGCCTTAGACACAGGTGGCACCTTCCGCCCGTGTCTCGGTAGAAGAAGATTAGCC  L L G A G A V A Y G V R E S V F T V E G G H R A I F F N R  TCGGTGGAGTGGAAGAACATATCCTGGCCGAAGGGCCTTCACTTCAGGATCCCTTGGTTCCAGTACCCCATTATCTATGACATTCGG AGCCACCTCACGTCGTCCTGGTAGAGACCGGCTCCCGGAAGTGAAGTCCTTAGGGAACCAAGGTCATGGGGTAATAGATACTGTAAGCC  I G G V Q Q D T I L A E G L H F R I P W F Q Y P I I Y D I R  CCCAGACCTCGAAAAAATCTCCTCCCCCTACAGGCTCCAAAGACCTACAGATGGTGAATATCTCCCTGGGAGTGTTGTCTCGACCCAATGCT  GGTTCTGGAACATTTTTAGAGGAGGGGATGTCCGAAGGTTTCTGGATGTTCACCACTTATAGAGGGACCTCAAACAGAGCTGGGTTACGA  A R P R K I S S P T G S K D L Q H V N I S L R V L S R P N A  CAGGAGCTTCCTAGCATGTACCAGCGCCTAGGGCTGACTACGAAGAACCAGGTGTTGCCCGTCCATTGTCAACGAGGTGCTCAAGAGTTGT  GTCCTCGAAGGATCGTACATGGCCCCAAGGCCCTAGGGCCTGACTGTCCCTTGCTCACAACGGCAGGTAACAGTTGCTCCACAAGAGTTCTCACAC  O E L P S M Y Q R L G L D Y E E R V L P S I V N E V L K S V  CTGGCCCAAGTTCAATGCCTCACAGCTGATCACCCCAGCGGCCCCAGGTATCCCTTGTTGATCCGCCGGGGAGCTGACAGAGAGGGCCCAAGGAC  AACTGGGCCAAGTTCAATGCCTCACAGCTGATCACCCAGCGGCCCCAGGTTATCCCTGTTGATCCGCCGGGGAGCTGACAGAGAGGGGCCAAGGAC  AACTGGGCCAAGTTCAATGCCTCACAGCTGATCACCCAGCGGCCCCAGGTCCATAGGGACAACCAAGGGGGCCCAAGGAC  AACTGGGCCAAGTTCAAGTTACCGGATGATCACCCAACAGAGCTGACCCTGACTGTTGCCCGGGGGCCCTCGACTGTCCTCCCGGTTCCTG  AAGTCGGGATGAGACCTACTACACCGGATCCTCACAGAGCTGAAACCAAGGGGCCCTCGACTGTCTCCCCGGTTCCTC  F S L I L D D V A I T E L S F S R E Y T A A V E A K Q V A Q  CAGGGAGGCCCCAGCGGCCCCAATTCTTGGTAGAAAACCAAAGAGAACAAGCAAAACAAAGCAAAAAAACCAAAAGCAAAAACAAAGCAAAAACAAAGCAAAAACAAAGCAAAAACAAAGCAAAAACAAAGCAAAAACAAAGCAAAAACAAAGCAAAAACAAAGCAAAAACAAAGCAAAAACAAAAGCAAAAACCAAAACCAAGGGCCCCAAGCCCCGGCTCTTTTTAACACCGGCCCGGGTCCCACTCCCGGCTCCCACTCCCGGCTCCCACTCCCGGCTCCCACTCCCGGCTCCCACTCCCGGCTCCCACTCCCGGCTCCCCACTCCCGGCTCCCACTCCCGCCCCGGCTCCCAC	ACAAC L L TCGG1 AGCCA	CGA L TG0 AC0	GAG	CCC G TGC	GGC A	G G	A GACA	V ACT/	A A	Y TG(	G G	CAC(	GCG(									+	+				+			+
ACACGACCCCGGCCGGGCCCGGGTCCCCGGATGCCCACACGGGTTAGACACAAGTGGCACCTTCCGCCCGGTGTCTCGGTAGAAAAATTTGGCCCCACTCACACGGGTCCCCGGTTCCCGGTTCCCGGTTCCCGGTTCCCCGGTTCCCCGGTTCCCCGGTTCCCCGGTTCCCCGGTTCCCCGGTTCCCCGGTTCCCCGGTTCCCCGGTTCCCCGGTTCCCCGGTTCCCCGGTTCCCCGGTTCCCCGGTTCCCCGGTTCCCCGGTTCCCCCGGTTCCCCCACTTCCGGTTCCCCCACTTCCGGTTCCCCCCACTTCCGGTTCCCCCCACTTCCGGTTCCCCCCACTTCCGGTTCCCCCCACTTCCGGTTCCCCCCCACACACA	TCGG1 AGCCA	TG(	GAG CTC	G TGC	A	G AG	A GAC	V ACTA	A	Y TG(	G	٧	R	E	S S	CACA	AAG	rggc	ACC	TTC	CGC	CCG	TGT	стс	GGT	AGA	AAGA	AGT	TA	GCC
TEGGTGGAGTGCAGCAGGACACTATCCTGGCCGAGGGCCTTCACTTCAGGATCCCTTGGTTCCAGTACCCCATTATCTATGACATTCGG AGCCACCTCACGTCGTCCTGTGATAGGACCGGCTCCCGGAAGTGAAGTCCTTAGGGAACCAAGGTCATGGGGTAATAGATACTGTAAGCC  I G G V Q Q D T I L A E G L H F R I P W F Q Y P I I Y D I R  CCAGACCTCGAAAAATCTCCTCCCCTACAGGCTCCAAAGACCTACAGATGGTGAATATCTCCCTGCGAGTTGTTCTCGACCCAATGCT  GGTCTGGAGCTTTTTAGAGGAGGGGGATGTCCGAAGGACCTACAGATGGTGAATATCTCCCTGCGAGTTGTTCTCGACCCAATGCT  A R P R K I S S P T G S K D L Q M V N I S L R V L S R P N A  CAGGAGCTTCCTAGCATGTACCAGGCCTAGGGCTGGACTACAGAGACGAGGTGTTGCCGTCCATTGTCAAACGAGGTGCTCAAGAGTTGG  GTCCTCGAAGGATCGTACAATGGTCGGGGATCCCGACCTGATGCTCCTTGCTCACAACGGCAGGTAACAGTTGCTCCACGAGTTCTCACAC  Q E L P S M Y Q R L G L D Y E E R V L P S I V N E V L K S V  CTGGCCAAGTTCAATGCCTCACAGCTGATCACCCAGCGGGCCCAGGGTATCCCTGTTGATCCGCCGGGAGCTGACAGAGGGGCCAAGGAC  CACCGGTTCAATGCCTCACAGCTGATCACCCAGCGGGCCCAGGGTCCATAGGGACACAACAAGTGCCCCAGGTCCTCC  V A K F N A S Q L I T Q R A Q V S L L I R R E L T E R A K D  TTCCAGCCTCATCCTGGATGATGTGGCCATCACAGAGCTGAGCTTTAGCCGAGAGTACAACAAGTGGCCCAAG  AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCCGACTTAGGCGGAGCACAACAACAAGTGGCCCAAG  AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCGACTGAAACAAGTGGCCCAAG  F S L I L D D V A I T E L S F S R E Y T A A V E A K Q V A Q  CAAGGAGGCCCCAGGGGCCCCAATTCTTGGTAGAAAAAAGCAAAAGCAAAAGCAAAAGCAAAGCAGAACAAGGGGCCGAGGGTGAGGCCCAAGCACCACCAGGCCCAGGCTCTCCTGGCTCCCACCCCGGCTCCCACCCCGGCTCCCACCCCGGCTCCCACCCCGGCTCCCCACCCCGGCTCCCACCCCGGCTCCCCACCCCGGCTCCCCACCCCGGCTCCCCACCCCGGCTCCCCCCCC	AGCC/	ACC	CTC								G GCC	۷	R	Ε	S	٧	F													
AGCCACCTCACGTCGTCCTGTGATAGGACCGGCTCCCGGAAGTGAAGTCCTAGGGAACCAAGGTCATGGGGTAATAGATACTGTAAGCC  I G G V Q Q D T I L A E G L H F R I P W F Q Y P I I Y D I R  CCCAGACCTCGAAAAATCTCCTCCCCTACAGGCTCCAAAGACCTACAGATGGTGAATATCTCCCTGCGAGTGTTGTCTCGACCCAATGCT  CGGTCTGGAGCTTTTTAGAGGAGGGGGATGTCCGAGGTTTCTGGATGTTCTACCACTTATAGAGGGACGTCCAAACAGAGCTGGGTTACGA  A R P R K I S S P T G S K D L Q M V N I S L R V L S R P N A  CAGGAGCTTCCTAGCATGTACCAGCGCCTAGGGCTGACCAGAGAGCAGGAACGAGGTGTTGCCGTCCATTGTCAACGAGGTGCTCAAGAGTGTG  CTCCTCGAAGGATCGTACATGGTCGGCGGATCCCGACCTGATGCTCCTTGCTCACAACGGCAGGTAACAGTTGCTCCACAGAGTTGTC  CTCCTCGAAGGTTCAATGCCTCACAGCTGATCACCCAGCGGGCCCAGGTTCCCTTGCTCACAACGGCAGGTAACAGTTGCTCCACAGAGTTCC  CACCGGTTCAAGTTCAATGCCTCACAGCTGATCACCCAGCGGGGCCCAGGGTACCAGGACGAGCTGACAGAGAGGGCCAAGGAC  CACCGGTTCAAGTTACAGGAGTGTCGACTAGTGGGTCGCCCGGGGTCCATAGGGACAACTAGGCGGCCCTCGACTGTCTCTCCCGGTTCCTG  V A K F N A S Q L I T Q R A Q V S L L I R R E L T E R A K D  CTCCAGCCTCATCCTGGATGATGTGGCCATCACAGAGCTGACAGAGCTTAGCCGAGAGTACACAGCTGCTGTAGAAGCCAAACAAGTGGCCCAG  F S L I L D D V A I T E L S F S R E Y T A A V E A K Q V A Q  CAGGAGGCCCAGCGGGCCCAATTCTTGGTAGAAAAAGCAAAAAGCAAAAGCAAAAGCAGAAGCAGGAACAAGCGGCCGAGGGTGAGGCCCAGGCTCCCACTCCCGGCTCCCAC  TAGGAGGCCCCAGCGGGCCCAATTCTTGGTAGAAAAAGCAAAAAGCAAAAGCAAAAGCAAAAGCAAAAGCAAAAAGCAAAAGCAAAAAGCAAAAGCAAAAGCAAAAAGCAAAAAGCAAAAAA	AGCC/	ACC	CTC								GCC	כ א רי					•	I	٧	ε	G	G	н	R	Α	I	F	F	N	R
AGCCACCTCACGTCGTCGTGGTGATAGGACCGGCTCCCGGAAGTGAAGTCCTAGGGAACCAAGGTCATGGGGTAATAGATACTGTAAGGC  I G G V Q Q D T I L A E G L H F R I P W F Q Y P I I Y D I R  CCCAGACCTCGAAAAATCTCCTCCCCCTACAGGCTCCAAAGACCTACAGATGGTGAATATCTCCCTGGGAGTGTTGTCTCGACCCAATGGT  GGTCTGGAGCTTTTTAGAGGAGGGGGATGTCCGAGGTTTCTGGATGTCTACCACTTATAGAGGGACGCTCACAACAGAGCTGGGTTACGA  A R P R K I S S P T G S K D L Q H V N I S L R V L S R P N A  CAGGAGCTTCCTAGCATGTACCAGCGCCTAGGGCTGGACTACGAGGAACGAGTGTTGCCGTCCATTGTCAACGAGGTGCTCAAGAGTGTG  GTCCTCGAAGGATCGTACATGGTCGCGGATCCCGACCTGATGTCCACCAACGGCAGGTAACAGTTGCTCCACGAGTTCTCACAC  Q E L P S H Y Q R L G L D Y E E R V L P S I V N E V L K S V  GTGGCCAAGTTCAATGCCTCACAGCTGATCACCCCAGCGGGCCCAAGGACCAACTAGGCGGCCCTCGACTGTCTCTCCCGGTTCCTG  V A K F N A S Q L I T Q R A Q V S L L I R R E L T E R A K D  GTCCAGCCTCATCCTGGATGATGTGGCCATCACACGAGGTGCCCTGGAACCAACAGCTGCTGTAGAAGCCAAACAAGTGGCCCAG  AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCCTCGGAAATCGGCTCTCATGTGTCGACGACCAACAACAAGTGGCCCAG  F S L I L D D V A I T E L S F S R E Y T A A V E A K Q V A Q  CAGGAGGCCCAAGCGGGCCCAATTCTTGGTAGAAAAAAGCAAAAAGCAGAAAGAA	I G	AC	3	ACG V	a a	TCC	TG	rga1	TAGO	SACI		GAGI	GGC	CTTC	CAC	TTC	AGG.	ATCC	CTI	rgĢī	тсс	AGT	ACÇ	CCA	TTA	TC	TATO	ACA	TT	CGG
CCAGACCTCGAAAAATCTCCTCCCCTACAGGCTCCAAAGACCTACAGTGGTGAATATCTCCCTGCGAGTGTTGTCTCGACCCAATGCT-42  GGTCTGGAGCTTTTTAGAGGAGGGGGATGTCCGAGGTTTCTGGATGTCTACCACTTATAGAGGGACGCTCACAACAGAGCTGGGTTACGA  A R P R K I S S P T G S K D L Q M V N I S L R V L S R P N A  AGGAGCTTCCTAGCATGTACCAGCGCCTAGGGCTGGACTACGAGGAACGAGGTGTTGCCGTCCATTGTCAACGAGGTGCTCAAGAGTGTG  GTCCTCGAAGGATCGTACCAGCGCCTAGGGCTGGACTACGAGGAACGAGGTGTTGCCGTCCATTGTCAACGAGGTGCTCAAGAGTGTG  GTCCTCGAAGGATCGTACCAGGGGATCCCGACCTGATGCTCCTTGCTCACAACGGCAGGTAACAGTTGCTCCACGAGGTTCTCACAC  Q E L P S M Y Q R L G L D Y E E R V L P S I V N E V L K S V  GTGGCCAAGTTCAATGCCTCACAGCTGATCACCCAGCGGGCCCAGGTACCCTGTTGATCCGCCGGGAGCTGACAGAGAGGGCCAAGGAC  CACCGGTTCAAGTTACGGAGTGTCGACTAGTGGGTCGCCCGGGTCCATAGGGACAACTAGGCGGCCCTCGACTGTCTCTCCCGGTTCCTG  V A K F N A S Q L I T Q R A Q V S L L I R R E L T E R A K D  GTCAGCCTCATCCTGGATGATGTGGCCATCACAGAGGCTGACTTAGGCCGAAACACGGCTGCTGTAGAAGCCAAACAAA	CCAG	AC	G - TC	٧	a	0					CGG	CTC	CCG	GAA	GTG	AAG	TCC	TAGG	GAA	ACCA	AGG	TCA	TGG	GGT	AAT	AG	ATA(	TGT	AA	GCC
A R P R K I S S P T G S K D L Q M V N I S L R V L S R P N A  **AGGAGCTTCCTAGCATGTACCAGCGCCTAGGGCTGGACTACGAGGAACGAGGTGTTGCCGTCCATTGTCAACGAGGTGCTCAAAAGGTGTGG **CCCTCGAAGGATCGTACCAGCGCCTAGGGCTGGACTACGAGGAACGAGGTGTTGCCGTCCATTGTCAACGAGGTGCTCAAAAGGTGTGG **CCCTCGAAGGATCGTACCATGTCACCCGACCTGATGCTCCTTGCTCACAACGGCAGGTAACAGTTGCTCCACGAGTTCTCACAC  **Q E L P S M Y Q R L G L D Y E E R V L P S I V N E V L K S V  **CTGGCCAAGTTCAATGCCTCACAGCTGATCACCCAGCGGGCCCAGGTATCCCTGTTGATCCGCCGGGAGCTGACAGAGAGGGCCAAGGAC **CACCGGTTCAAGTTACGGAGTGTCGACCTAGTGGGTCGCCCGGGTCCATAGGGACAACTAGGCGGCCCTCGACTGTCTCTCCCGGTTCCTC **V A K F N A S Q L I T Q R A Q V S L L I R R E L T E R A K D  **CTCAGCCTCATCCTGGATGATGTGGCCCATCACAGAGCTGAGCTTTAGCCGAGAGTACACAGCTGCTGTAGAAGCCAAACAAGTGGCCCAG **AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCGACTGAAATCAGGCTCTCATGTGTCGACGACAACTATCTTCGTTTCACCGGGTC **F S L I L D D V A I T E L S F S R E Y T A A V E A K Q V A Q  **CAGGAGGCCCAGCGGGCCCAATTCTTGGTAGAAAAAAGCAAAGCAAAGCAAGC			ר דר			u	0	T	ı	L	Α	Ε	G	L	н	F	R	ı	P	w	F	Q	Y	Р	i	I	Υ	D	I	R
A R P R K I S S P T G S K D L Q M V N I S L R V L S R P N A  **AGGAGCTTCCTAGCATGTACCAGCGCCTAGGGCTGGACTACGAGGAACGAGGTGTTGCCGTCCATTGTCAACGAGGTGCTCAAAAGGTGTGG **CCCTCGAAGGATCGTACCAGCGCCTAGGGCTGGACTACGAGGAACGAGGTGTTGCCGTCCATTGTCAACGAGGTGCTCAAAAGGTGTGG **CCCTCGAAGGATCGTACCATGTCACCCGACCTGATGCTCCTTGCTCACAACGGCAGGTAACAGTTGCTCCACGAGTTCTCACAC  **Q E L P S M Y Q R L G L D Y E E R V L P S I V N E V L K S V  **CTGGCCAAGTTCAATGCCTCACAGCTGATCACCCAGCGGGCCCAGGTATCCCTGTTGATCCGCCGGGAGCTGACAGAGAGGGCCAAGGAC **CACCGGTTCAAGTTACGGAGTGTCGACCTAGTGGGTCGCCCGGGTCCATAGGGACAACTAGGCGGCCCTCGACTGTCTCTCCCGGTTCCTC **V A K F N A S Q L I T Q R A Q V S L L I R R E L T E R A K D  **CTCAGCCTCATCCTGGATGATGTGGCCCATCACAGAGCTGAGCTTTAGCCGAGAGTACACAGCTGCTGTAGAAGCCAAACAAGTGGCCCAG **AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCGACTGAAATCAGGCTCTCATGTGTCGACGACAACTATCTTCGTTTCACCGGGTC **F S L I L D D V A I T E L S F S R E Y T A A V E A K Q V A Q  **CAGGAGGCCCAGCGGGCCCAATTCTTGGTAGAAAAAAGCAAAGCAAAGCAAGC				GAA	AAA	TÇI	rcc'	TCC	CCTA	ACA(	GGC	TCC.	ΑΑΑ	GAC	CTA	CAG	ATG	GTGA	ATA	ATCI	rccc	TGC	GAG	TGT	TGT	CT	CGA	CCA	AT	GCŢ
AGGAGCTTCCTAGCATGTACCAGCGCCTAGGGCTGGACTACGAGGAACGAGTGTTGCCGTCCATTGTCAACGAGGTGCTCAAGAGTGTG  STCCTCGAAGGATCGTACATGGTCGCGGATCCCGACCTGATGCTCCTTGCTCACAACGGCAGGTAACAGTTGCTCCACGAGTTCTCACAC  Q E L P S M Y Q R L G L D Y E E R V L P S I V N E V L K S V  STGGCCAAGTTCAATGCCTCACAGCTGATCACCCAGCGGGCCCAGGTATCCCTGTTGATCCGCCGGGAGCTGACAGAAGAGGGCCAAGGAC  CACCGGTTCAAGTTACGGAGTGCGACTAGTGGGTCGCCCGGGTCCATAGGGACAACTAGGCGGCCCCTCGACTGTCTCTCCCCGGTTCCTG  V A K F N A S Q L I T Q R A Q V S L L I R R E L T E R A K D  STCAGCCTCATCCTGGATGATGTGGCCATCACAGAGCTGAGCTTTAGCCGAGAGTACACAGCTGCTGTAGAAGCCAAACAAGTGGCCCAG  AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCGACTCGAAATCGGCTCTCATGTGTCGACCGAC	GGTC.													<del>,</del>									$\overline{}$				- 1			
CTCCTCGAAGGATCGTACATGGTCGCGGATCCCGACCTGATGCTCCTTGCTCACAACGGCAGGTAACAGTTGCTCCACGAGTTCTCACAC  O E L P S M Y O R L G L D Y E E R V L P S I V N E V L K S V  STGGCCAAGTTCAATGCCTCACAGCTGATCACCCCAGCGGGCCCAGGGTATCCCTGTTGATCCGCCGGGAGCTGACAGAGAGGGCCAAGGAC CACCGGTTCAAGTTACGGAGTGTCGACTAGTGGGTCGCCCGGGTCCATAGGGACAACTAGGCGGCCCTCGACTGTCTCCCCGGTTCCTG  V A K F N A S O L I T O R A O V S L L I R R E L T E R A K D  STCCAGCCTCATCCTGGATGATGTGGCCATCACAGAGCTGAGCTTTAGCCGAGAGTACACAGCTGCTGTAGAAGCCAAACAAGTGGCCCAG AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCGACTCGAAATCGGCTCTCATGTGTCGACGACCATCTTCGGTTTGTTCACCGGGTC  F S L I L D D V A I T E L S F S R E Y T A A V E A K O V A O  CAGGAGGCCCAGCGGGCCCCAATTCTTGGTAGAAAAAGCAAAAGCAGAGAGAACAGCGGCAGAAAAATTGTGCAGGCCGAGGGTGAGGCCGAGGCT  GTCCTCCGGGTCGCCCGGGTTAAGAACCATCTTTTTCGTTTCGTCTCTCCTTGTCGCCGTCTTTTAACACGTCCGGCTCCCACTCCGGCTCCGA	ΔR		P	R	ĸ	1	s	s	ρ	Т	G	s	ĸ	D	L	a	м	٧	N	i	S	L	R	<b>v</b> .	L	s	R	ρ	N	Α
CTCCTCGAAGGATCGTACATGGTCGCGGATCCCGACCTGATGCTCCTTGCTCACAACGGCAGGTAACAGTTGCTCCACGAGTTCTCACAC  O E L P S M Y O R L G L D Y E E R V L P S I V N E V L K S V  STGGCCAAGTTCAATGCCTCACAGCTGATCACCCCAGCGGGCCCAGGGTATCCCTGTTGATCCGCCGGGAGCTGACAGAGAGGGCCAAGGAC CACCGGTTCAAGTTACGGAGTGTCGACTAGTGGGTCGCCCGGGTCCATAGGGACAACTAGGCGGCCCTCGACTGTCTCCCCGGTTCCTG  V A K F N A S O L I T O R A O V S L L I R R E L T E R A K D  STCCAGCCTCATCCTGGATGATGTGGCCATCACAGAGCTGAGCTTTAGCCGAGAGTACACAGCTGCTGTAGAAGCCAAACAAGTGGCCCAG AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCGACTCGAAATCGGCTCTCATGTGTCGACGACCATCTTCGGTTTGTTCACCGGGTC  F S L I L D D V A I T E L S F S R E Y T A A V E A K O V A O  CAGGAGGCCCAGCGGGCCCCAATTCTTGGTAGAAAAAGCAAAAGCAGAGAGAACAGCGGCAGAAAAATTGTGCAGGCCGAGGGTGAGGCCGAGGCT  GTCCTCCGGGTCGCCCGGGTTAAGAACCATCTTTTTCGTTTCGTCTCTCCTTGTCGCCGTCTTTTAACACGTCCGGCTCCCACTCCGGCTCCGA										C T 4				T A C	~ 4 ~		CC 4	C T C 1				. TTC	TC /	. A C C	: A G f	e T G	רדר	ΔΑΓΑ	LG T	GTG
Q E L P S M Y Q R L G L D Y E E R V L P S I V N E V L K S V  GTGGCCAAGTTCAATGCCTCACAGCTGATCACCCAGCGGGCCCAGGTATCCCTGTTGATCCGCCGGGAGCTGACAGAGAGGGCCAAGGAC CACCGGTTCAAGTTACGGAGTGTCGACTAGTGGGTCGCCCGGGTCCATAGGGACAACTAGGCGGCCCTCGACTGTCTCTCCCGGTTCCTG  V A K F N A S Q L I T Q R A Q V S L L I R R E L T E R A K D  GTCAGCCTCATCCTGGATGATGTGGCCATCACAGAGCTGAGCTTAGCCGAGAGTACACAGCTGCTGTAGAAGCCAAACAAGTGGCCCAG AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCGACTCGAAATCGGCTCTCATGTGTCGACGACATCTTCGGTTTGTTCACCGGGTC  F S L I L D D V A I T E L S F S R E Y T A A V E A K Q V A Q  CAGGAGGCCCAGCGGGCCCAATTCTTGGTAGAAAAAGCAAAGCAAAGCAGGGAACAGCGGCAGAAAATTGTGCAGGCCGAGGGTGAGGCCGAGGCT  GTCCTCCGGGTCGCCCGGGTTAAGAACCATCTTTTTCGTTTCGTCCTTGTCGCCGTCTTTTAACACGTCCGGCTCCCACTCCGGCTCCGA										-+-				+			-+-		—			<del></del>								
STEGECCAAGTTCAATGCCTCACAGCTGATCACCCAGCGGGCCCAGGTATCCCTGTTGATCCGCCGGGAGCTGACAGAGAGGGCCAAGGAC CACCGGTTCAAGTTACGGAGTGTCGACTAGTGGGTCGCCCGGGTCCATAGGGACAACTAGGCGGCCCTCGACTGTCTCTCCCGGTTCCTG  V A K F N A S Q L I T Q R A Q V S L L I R R E L T E R A K D  TTCAGCCTCATCCTGGATGATGTGGCCATCACAGAGCTGAGCTTTAGCCGAGAGTACACAGCTGCTGTAGAAGCCAAACAAGTGGCCCAG  AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCGACTCGAAATCGGCTCTCATGTGTCGACGACATCTTCGGTTTGTTCACCGGGTC  F S L I L D D V A I T E L S F S R E Y T A A V E A K Q V A Q  CAGGAGGCCCAGCGGGCCCAATTCTTGGTAGAAAAAGCAAAGCAGAGCAGGAACAGCGGCAGAAAATTGTGCAGGCCGAGGGTGAGGCCGAGGCT  GTCCTCCGGGTCGCCCGGGTTAAGAACCATCTTTTTCGTTTCGTCCTTGTCGCCGTCTTTTAACACGTCCGGCTCCCACTCCGGCTCCGA	116611	L G	ААЬ	IGA I	LUI	AL	A I G	G 1 C	GCG	GA 1		dac								_					_	.,		· ·		
CACCGGTTCAAGTTACGGAGTGTCGACTAGTGGGTCGCCCGGGTCCATAGGGACAACTAGGCGGCCCTCGACTGTCTCCCCGGTTCCTG  V A K F N A S Q L I T Q R A Q V S L L I R R E L T E R A K D  TICAGCCTCATCCTGGATGATGTGGCCATCACAGAGCTGAGACTTTAGCCGAGAGTACACAGCTGCTGTAGAAGCCAAACAAGTGGCCCAG  AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCGACTCGAAATCGGCTCTCATGTGTCGACGACATCTTCGGTTTGTTCACCGGGTC  F S L I L D D V A I T E L S F S R E Y T A A V E A K Q V A Q  CAGGAGGCCCAGCGGGCCCAATTCTTGGTAGAAAAAAGCAAAGCAGAGGAACAGCGGCAGAAAATTGTGCAGGCCGAGGGTGAGGCCGAGGCT  GTCCTCCGGGTCGCCCGGGTTAAGAACCATCTTTTTCGTTTCGTCCTTGTCGCCGTCTTTTAACACGTCCGGCTCCCACTCCGGCTCCGA	Q E		L	Р	S		Y	•		L	_	L	_		_			٧	L	Р	-	1	٧	N	t.	٧	L	K	· .	•
V A K F N A S Q L I T Q R A Q V S L L I R R E L T E R A K D  TTCAGCCTCATCCTGGATGATGTGGCCATCACAGAGCTGAGCTTTAGCCGAGAGTACACAGCTGCTGTAGAAGCCAAACAAGTGGCCCAG  AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCGACTCGAAATCGGCTCTCATGTGTCGACGACATCTTCGGTTTGTTCACCGGGTC  F S L I L D D V A I T E L S F S R E Y T A A V E A K Q V A Q  CAGGAGGCCCAGCGGGCCCAATTCTTGGTAGAAAAAGCAAAGCAGAGGAACAGCGGCAGAAAATTGTGCAGGCCGAGGGTGAGGCCGAGGCT  GTCCTCCGGGTCGCCCGGGTTAAGAACCATCTTTTTCGTTTCGTCCTTGTCGCCGTCTTTTAACACGTCCGGCTCCCACTCCGGCTCCGA																				<del></del> +										
TTCAGCCTCATCCTGGATGATGTGGCCATCACAGAGCTGAGCTTTAGCCGAGAGTACACAGCTGCTGTAGAAGCCAAACAAGTGGCCCAG AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCGACTCGAAATCGGCTCTCATGTGTCGACGACATCTTCGGTTTGTTCACCGGGTC  F S L I L D D V A I T E L S F S R E Y T A A V E A K Q V A Q CAGGAGGCCCAGCGGGCCCAATTCTTGGTAGAAAAAGCAAAGCAGAGCAGGAACAGCGGCAGAAAATTGTGCAGGCCGAGGGTGAGGCCGAGGCT  GTCCTCCGGGTCGCCCGGGTTAAGAACCATCTTTTTCGTTTCGTCCTTGTCGCCGTCTTTTAACACGTCCGGCTCCCACTCCGGCTCCGA	ACCG	GT	TCA	AGI	ΓΤΑΟ	:GG	AGT	GTC	GAC	TAG	TGG	GTC	GCC	CGG	GTC	CAT	AGG	GAC	AAC	TAG	GCG	3000	TCC	GAC1	rgto	СТС	TCC	CGGT	ГТС	CTG
AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCGACTCGAAATCGGCTCTCATGTGTCGACGACATCTTCGGTTTGTTCACCGGGTC  F S L I L D D V A I T E L S F S R E Y T A A V E A K Q V A Q  CAGGAGGCCCAGCGGGCCCAATTCTTGGTAGAAAAAGCAAAGCAGGGAACAGCGGCAGAAAATTGTGCAGGCCGAGGGTGAGGCCGAGGCT  GTCCTCCGGGTCGCCCGGGTTAAGAACCATCTTTTTCGTTTCGTCCTTGTCGCCGTCTTTTAACACGTCCGGCTCCCACTCCGGCTCCGA	V A		ĸ	F	N	Α	s	Q	L	ı	Ŧ	a	R	A	Q	٧	s	L	L	i	R	R	ε	L	T	Ε	R	A	K	0
AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCGACTCGAAATCGGCTCTCATGTGTCGACGACATCTTCGGTTTGTTCACCGGGTC  F S L I L D D V A I T E L S F S R E Y T A A V E A K Q V A Q  CAGGAGGCCCAGCGGGCCCAATTCTTGGTAGAAAAAGCAAAGCAGGGAACAGCGGCAGAAAATTGTGCAGGCCGAGGGTGAGGCCGAGGCT  GTCCTCCGGGTCGCCCGGGTTAAGAACCATCTTTTTCGTTTCGTCCTTGTCGCCGTCTTTTAACACGTCCGGCTCCCACTCCGGCTCCGA	TCAG	CC	TCA	TC	CTG	GAT	GAT	GTG	GCC.	ATC	ACA	GAG	CTG	AGC	TTT	AGC	CGA	GAG	TAC.	ACA	GCT	GCTO	TAC	GAAC	GCC/	AAA	CAA	GTG	gcc	CAĢ
CAGGAGGCCCAGCGGGCCCAATTCTTGGTAGAAAAAGCAAAGCAGGGAACAGCGGCAGAAAATTGTGCAGGCCGAGGGTGAGGCCGAGGCT GTCCTCCGGGTCGCCCGGGTTAAGAACCATCTTTTTCGTTTCGTCCTTGTCGCCGTCTTTTAACACGTCCGGCTCCCACTCCGGCTCCGA	AGTC	GG	AGT	r A G (	GAC	TA	<del>C T A</del>	CAC	CGG	TAG	TGT	CTC	GAC	TCG	AAA	TCG	GCT	CTC	ATG	TGT	CGA	CGA	CAT	TTC	GG.	111	GTT	CAC	CGG	GTC
GTCCTCCGGGTCGCCCGGGTTAAGAACCATCTTTTTCGTTTCGTCCTTGTCGCCGTCTTTTAACACGTCCGGCTCCCACTCCGGCTCCGA	F S	3	L	i.	L	D	0	٧	Α	Į	Ţ	ε	L	s	F	S	R	E	Υ	т	Α	A	٧	Ε	A	K	Q	٧	A	a
GTCCTCCGGGTCGCCCGGGTTAAGAACCATCTTTTTCGTTTCGTCCTTGTCGCCGTCTTTTAACACGTCCGGCTCCCACTCCGGCTCCGA	TAGGA	166	ccc	AGI	CGGC	SCC	CAA	TTC	TTG	GTA	GAA	AAA	GCA	AAG	CAG	GAA	CAG	CGG	CAG	AAA	ATT:	GTG	CAG	GCC	GAG	GGT	GAG	GCC	GAG	GCŢ
										_							<del></del>		+					<del></del>					,	•
Q E A Q R A Q F L V, E K A K Q E Q R Q K I V Q A E G E A E A																														
CCCAACATCCTTGGAGAAGCACTGAGCAAGAACCCTGGCTACATCAAACTTCGCAAGATTCGAGCAGCCCAGAATATCTCCAAGACGATC																														SATO
CGGTTCTACGAACCTCTTCGTGACTCGTTCTTGGGACCGATGJAGTTTGAAGCGTTCTAAGCTCGTCGGGTCTTATAGAGGTTCTGCTAG																	${ o}$			$\rightarrow$				+						
AKMLGEALSKNPGYIKLRKIRAAQNISKTI								-																						

HMWG546

Page 2

GCCACATCACAGAATCGTATCTCACAGCTGACAACCTTGTGCTGAACCTACAGGATGAAAGTTTCACCAGGGGAAGTGACAGCCTC
CGGTGTAGTGTCTTAGCATAGATAGAGTGTCGACTGTTGGAACACGACTTGGATGTCCTACTTTCAAAGTGGTCCCCTTCACTGTCGGAG

ATSONRIYLTADNL"VLNLODESFTRGSOSL

ATCAAGGGTAAGAAATGAGCCTAGTCACCAAGAACTCCACCCCCACAAGAAGTGGATCTGCTTCTCCAGTTTTTGA
TAGTTCCCATTCTTTACTCGGATCAGTGGTTCTTGAGGTGGGGGGGTGTTCTTCACCTAGACGAAGAGGTCAAAAACT

(SCQIDN

IKGKK (SCQIDNO: 31)

BNSDOCID: <WO\_\_\_9831799A2\_I\_>

HIMF. WOG

CGTGCTCTACTGTAGTGATTCACCGGCTAGACGTGTCTCGTCCGGTCCTCGTTGGTGTGTCCGAAGGACGTGTACCTGACGCTCTAGTT  M D C E I K  GGCCGCCCCTGCTGCATCGGCACCAAGGGCAGCTGTGAGATCACCACCCGGGAATACTGTGAGTTCATGCACGGCTATTTCCATGAGGA CCGGCGGGGGACGACGTAGCCGTGGTTCCCGTCGACACTCTAGTGGTGGGGCCCTTATGACACTCAAGTACGTGCCGATAAAGGTACTCCT  G R P C C I G T K G S C E I T T R E Y C E F M H G Y F H E E  GCAACACTCTGCTCCCAGGTGAGGCGAGGCCTGGAGTAGTGGAGGAGGAGGACGCTGGGGCATGGCAGCCTGGGGCCCGAG  A T L C S Q V R R G R P G V V E E R T L G M A A C W G R G S  CCGCACTCCCTCCCATGTCGGAGCCTCAGACTCAGGCTGCTTCTGGGGCCCCATATGCCCATTCCCAGGTGCACTGTTTTGGA GCGTGAGGGGAGGG
GGCCGCCCTGCTGCATCGGCACCAAGGGCAGCTGTGAGATCACCACCCGGGAATACTGTGAGTTCATGCACGGCTATTTCCATGAGGA CCGGCGGGGACGACGTAGCCGTGGTTCCCGTCGACACTCTAGTGGTGGGGCCCTTATGACACTCAAGTACGTGCCGATAAAGGTACTCCT  G R P C C I G T K G S C E I T T R E Y C E F M H G Y F H E E  GCAACACTCTGCTCCCAGGTGAGGCGAGGCCTGGAGTAGTGGAGGAGGAGGACGCTGGGCATGGCAGCCTGCTGGGGCCCGGGCCCC CGTTGTGAGACGAGGGTCCACTCCGCTCCG
CCGGCGGGGACGACGTAGCCGTGGTTCCCGTCGACACTCTAGTGGTGGGCCCTTATGACACTCAAGTACGTGCCGATAAAGGTACTCCT  G R P C C I G T K G S C E I T T R E Y C E F M H G Y F H E E  GCAACACTCTGCTCCCAGGTGAGGCGAGGCCTGGAGTAGTGGAGGAGGACGCTGGGGCATGGCAGCCTGCTGGGGCCGGGCTC  CGTTGTGAGACGAGGGTCCACTCCGCTCCG
CCGGCGGGGACGACGTAGCCGTGGTTCCCGTCGACACTCTAGTGGTGGGCCCTTATGACACTCAAGTACGTGCCGATAAAGGTACTCCT  G R P C C I G T K G S C E I T T R E Y C E F M H G Y F H E E  GCAACACTCTGCTCCCAGGTGAGGCGAGGCAGGCCTGGAGTAGTGGAGGAGGAGGACGCTGGGCAGGCA
GCAACACTCTGCTCCCAGGTGAGGCAGGCCTGGAGTAGTGGAGGAGGAGGACGCTGGGCAGGCTGCTGGGGGCCGGGGCTC  CGTTGTGAGACGAGGGTCCACTCCGCTCCG
CGTTGTGAGACGAGGGTCCACTCCGCTCCGGACCTCATCACCTCCTCCTGCGACCGTACCGTCGGACGACCCCGGCCCCGAG  A T L C S Q V R R G R P G V V E E R T L G M A A C W G R G S  CGCACTCCCTCCCATGTCGGAGCCTCAGACTCAGGCTGCTTCTGGGGCGCTGAGCACCCATATGCCCATTCCCAGGTGCACTGTTTTGGA 3
CGTTGTGAGACGAGGGTCCACTCCGCTCCGGACCTCATCACCTCCTCCTGCGACCGTACCGTCGGACGACCCCGGCCCCGAG  A T L C S Q V R R G R P G V V E E R T L G M A A C W G R G S  CGCACTCCCTCCCATGTCGGAGCCTCAGACTCAGGCTGCTTCTGGGGCGCTGAGCACCCATATGCCCATTCCCAGGTGCACTGTTTTGGA 3
CGCACTCCCTCCCATGTCGGAGCCTCAGACTCAGGCTGCTTCTGGGGGGGCGCTGAGCACCATATGCCCATTCCCAGGTGCACTGTTTTGGA
R.TPSHVGASDSGCFWGAEHHMPIPRCTVLD
AAGGTGTGTTGGGCTGCTGCCTTCCTCAACCCTGAGGTCCCAGATCAGTTTTACAGGTCTGGCTGTCTCTTTTCCTACATGTTGGGTAA
TTCCACACACCCGACGGAGGAGGAGGTTGGGACTCCAGGGTCTAGTCAAAATGTCCAGACCGACAGAGAAAAGGATGTACAACCCATT
K V C W A A F L N P E V P D Q F Y R S G C L F S Y M L G K
CAGGTCCTCAATGCCCCCGAACCCGACCCCTGTGATGGACACCCAGGCGGACCCCTGGGGAAAGGTTCCTGGGCCAGGGTATGGTCGGTC
TCCAGGAGTTACGGGGGGCTTGGGCTGGGGACACTACCTGTGGGTCCGCCTGGGGACCCCTTTCCAAGGACCCGGTCCCATACCAGCCAG
R S S M P P N P T P V M D T Q A D P W G K V P G P G Y G R S
CAACCTGCCGAAGACTACTGCTCCTGAAGTGTCTGGATGAAGGCCGCTGCCTGGTGTGTCCCTCCC
TTGGACGGCTTCTGATGACGAGGACTTCACAGACCTACTTCCGGCGACGGACCACACAGGGAGGG
NLPKTTAPEVSG. (SCOID NO:32)
(COSTO IV
TGTCTTGTGGGTCTTTTCAAATGACATCCCTGAAGGGGACCTGGAGGAAGGTGGTCCGGCTGGCACCCTATCGC  TOTAL T

# INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred on page 13 . Hine	ed to in the description Table 1
B. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an additional sheet
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country 12301 Parklawn Drive Rockville, Maryland 20852 United States of America	77)
Date of deposit May 16, 1997	Accession Number 209053
C. ADDITIONAL INDICATIONS (leave blank if not applicable	This information is continued on an additional sheet
	·
D. DESIGNATED STATES FOR WHICH INDICATIONS AI	RE MADE (if the indications are not for all designated States)
E. SEPARATE FURNISHING OF INDICATIONS (leave blan	ak if not applicable)
The indications listed below will be submitted to the International E Number of Deposit")	Bureau later (specify the general nature of the indications e.g., "Accession
For receiving Office use only	For International Bureau use only
Authorized officer Yvetta Simma  Paralagal Specialis: IAPD-PCT Operation	This sheet was received by the International Bureau on:  Authorized officer

REC'D

2 3 FER 1998

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

	WIFO FOT
A. The indications made below relate to the microorganism referre	ed to in the description
on page <u>13</u> . li <del>ne</del> <u>Ta</u>	able l
B. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an additional sheet
Name of depositary institution	
American Type Culture Collection	
Address of depositary institution (including postal code and country 12301 Parklawn Drive Rockville, Maryland 20852 United States of America	יי
Date of deposit	Accession Number
May 16, 1997	209053
C. ADDITIONAL INDICATIONS (leave blank if not applicable	This information is continued on an additional sheet
EUROPE	
the mention of the grant of the European application has been refused of withdrawn the issue of such a sample to an expert sample (Rule 28(4) EPC).  D. DESIGNATED STATES FOR WHICH INDICATIONS AND ADDRESS AND AD	n or is deemed to be withdrawn, only by nominated by the person requesting the
D. DESIGNATED STATES FOR WINCH MADICAL CO.	The manual state of the manual states of the
E. SEPARATE FURNISHING OF INDICATIONS (leave blan	nk if not applicable)
The indications listed below will be submitted to the International In Number of Deposit")	Bureau later ispecify the general nature of the indications e.g., "Accession
For receiving Office use only	For International Bureau use only
This sheet was received with the international application	This sheet was received by the International Bureau on:
Authorized officer very simple in Specialist	Authorized officer

Form PCT/RO/134 (July 1992)

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism reference on page 13 ./././_Tal	
B. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an additional sheet
Name of depositary institution  American Type Culture Collection	
Address of depositary institution (including postal code and cour 12301 Parklawn Drive Rockville, Maryland 20852 United States of America	utry)
Date of deposit May 16, 1997	Accession Number 209054
C. ADDITIONAL INDICATIONS (leave blank if not applicate	ble) This information is continued on an additional sheet
D. DESIGNATED STATES FOR WHICH INDICATIONS	ARE MADE (if the indications are not for all designated States)
E. SEPARATE FURNISHING OF INDICATIONS (leave bla	ank if not applicable)
The indications listed below will be submitted to the International Number of Deposit")	Bureau later (specify the general nature of the indications e.g., "Accession
For receiving Office use only	For International Bureau use only
This sheet was received with the international application	This sheet was received by the International Bureau on:
Authorized officer Cardinal Specialist APD-PCT Operations	Authorized officer

# INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

REC'D	2 3	FEB	1998
WiPO	PC	OT.	

A. The indications made below relate to the microorganism refer on page	· · · · · · · · · · · · · · · · · · ·
B. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an additional sheet
Name of depositary institution  American Type Culture Collection  Address of depositary institution (including postal code and coun	(קינו)
12301 Parklawn Drive Rockville, Maryland 20852 United States of America	
Date of deposit May 16, 1997	Accession Number 209054
C. ADDITIONAL INDICATIONS (leave blank if not applicab	le) This information is continued on an additional sheet
EUROPE In respect of those designations in whi of the deposited microorganism will be the mention of the grant of the Europea application has been refused or withdra by the issue of such a sample to an exp the sample (Rule 28(4) EPC).  D. DESIGNATED STATES FOR WHICH INDICATIONS A	made available until the publication n patent or until the date on which wn or is deemed to be withdrawn, only ert nominated by the person requesting
E. SEPARATE FURNISHING OF INDICATIONS (leave bla	
The indications listed below will be submitted to the International Number of Deposit") .	Bureau later (specify the general nature of the indications e.g., "Accession
For receiving Office use only	For International Bureau use only
Authorized officer  Authorized officer  Wisse Smins  Ratalegal Specialist  IAPD-PCT Operations	This sheet was received by the International Bureau on:  Authorized officer

15

20

25

35

What is claimed is:

- 1. An isolated polynucleotide comprising a nucleotide sequence that has at least 80% identity over its entire length to a nucleotide sequence encoding the receptor polypeptide of SEQ ID NO:Y; or a nucleotide sequence complementary to said isolated polynucleotide.
- The polynucleotide of claim 1 wherein said polynucleotide comprises the nucleotide sequence contained in SEQ ID NO:X encoding the receptor polypeptide of
   SEQ ID NO:Y.
  - 3. The polynucleotide of claim 1 wherein said polynucleotide comprises a nucleotide sequence that is at least 80% identical to that of SEQ ID NO:X over its entire length.

4. The polynucleotide of claim 3 which is polynucleotide of SEQ ID NO: X.

- 5. The polynucleotide of claim 1 which is DNA or RNA.
- 6. A DNA or RNA molecule comprising an expression system, wherein said expression system is capable of producing a receptor polypeptide comprising an amino acid sequence, which has at least 80% identity with the polypeptide of SEQ ID NO:Y when said expression system is present in a compatible host cell.
  - 7. A host cell comprising the expression system of claim 6.
- 8. A process for producing a receptor polypeptide comprising culturing a host of claim 7 under conditions sufficient for the production of said polypeptide and recovering the polypeptide from the culture.
  - 9. A process for producing a cell which produces a receptor polypeptide thereof comprising transforming or transfecting a host cell with the expression system of claim 6 such that the host cell, under appropriate culture conditions, produces a receptor polypeptide.
  - 10. A receptor polypeptide comprising an amino acid sequence which is at least 80% identical to the amino acid sequence of SEQ ID NO:Y over its entire length.

20

25

- 11. The polypeptide of claim 10 which comprises the amino acid sequence of SEQ ID NO:Y.
- 5 12. An antibody immunospecific for the receptor polypeptide of claim 10.
  - 13. A method for the treatment of a subject in need of enhanced activity or expression of receptor polypeptide of claim 10 comprising:
  - (a) administering to the subject a therapeutically effective amount of an agonist to said receptor; and/or
  - (b) providing to the subject polynucleotide of claim 1 in a form so as to effect production of said receptor activity in vivo.
- 14. A method for the treatment of a subject having need to inhibit activity or expression of the receptor polypeptide of claim 10 comprising:
  - (a) administering to the subject a therapeutically effective amount of an antagonist to said receptor; and/or
  - (b) administering to the subject a nucleic acid molecule that inhibits the expression of the nucleotide sequence encoding said receptor; and/or
  - (c) administering to the subject a therapeutically effective amount of a polypeptide that competes with said receptor for its ligand.
  - 15. A process for diagnosing a disease or a susceptibility to a disease in a subject related to expression or activity of the receptor polypeptide of claim 10 in a subject comprising:
  - (a) determining the presence or absence of a mutation in the nucleotide sequence encoding said receptor polypeptide in the genome of said subject; and/or
- (b) analyzing for the presence or amount of the receptor polypeptide
   30 expression in a sample derived from said subject.
  - 16. A method for identifying agonists to the receptor polypeptide of claim 10 comprising:
    - (a) contacting cells produced by claim 9 with a candidate compound;
- 35 and
  - (b) determining whether the candidate compound effects a signal generated by activation of the receptor polypeptide.

- 17. An agonist identified by the method of claim 16.
- 18. The method for identifying antagonists to the receptor polypeptide of claim 10 comprising:

- (a) contacting said cell produced by claim 9 with an agonist; and
- (b) determining whether the signal generated by said agonist is diminished in the presence of a candidate compound.
  - 19. An antagonist identified by the method of claim 18.

10

15

5

- 20. An isolated receptor polynucleotide comprising a nucleotide sequence selected from the group consisting of:
- (a) a nucleotide sequence having at least 80% identity to a nucleotide sequence encoding the receptor polypeptide expressed by the cDNA insert deposited at the ATCC; and
- (b) a nucleotide sequence complementary to the nucleotide sequence of (a).
- 21. A recombinant host cell produced by a method of Claim 9 or a membrane thereof expressing a receptor polypeptide.

FIGURE 1

				T
1	MLLLULLELE LUMGRERV MLPULLUPLUMG	20 7 E G Q K S N R K D	30 YSLTMOSSYTV	40 L / g E G M HAPCR70.AA / g E G L OB-1.aa
1			70	80
41	50 CMHVRCSESXFVDSQ	FDSDPVHGY	A STAR NOT SWK	ALP MAIT HMACR70.AA
36	CWHVRCSFSYPWRSW CWLYPCSFSYPWRSW 	1 2 度 1 度 1 x (米)	110	120
81.	90 如解PAWAUQEE型RDES	HEIGDPOTK	HA made to make M	S TI A G R HMACR70.AA
76	MARDANA VQEETRORF MARDANA VKPETRORF	Kibinininin vina vina	150	160
121	130	140 KYDQLSV		HMACR70.AA
116	Y E E R M E K G - N I K W N Y Y E E R V E R G R D V K Y S Y	QON KREET - V	190	200
142	170	180		HMACR70.AA
156			230	240
142	210	220		N V T V P HMACR70.AA
196	N		270	280
14'	250 7 PQNLEVTVTQGEGT	260 STALGNSSS	# c## # ed s 1 8 1	V C A V D HMACR70.AA
23	7 PONL EVEVE VE QGEQT P 5 PO FIRITR - NO I	ELECTOR LINE	SHE A DESIGNATION OF THE STREET	320
18	290 7 SNPPARLSWTWRS LI	300 PSNPPPPPPP	310 LVLELQ - VHLC	DEGER HMACR70.AA
27	7 SNPPARISWTWRS LI 3 SNPPAHLSWFQGSPI	A EN A TI PLIMENT	·	
22	330 6 1 C H A Q N S 1 G S Q H V S	340 LNLS	350	HMACR70.AA
31	6 TCRAONSIGSQHVS 3 TCRAOHPLGFLQIF	一 一 一 一 一 一 一 一 一 一 一 一 一 一 一 一 一 一 一		400
24	370	380	390 	 HMACR70.AA
35	G C S F R A R P A P S L C W R	LEEKPLEGNS		440
2	410 41 QQEYTGKM	420 P V	430	HMACR70.AA
	NSSLI HGGLSSDL	VSCKAWNI	YGSQSGSVLLL	V C K 2 M D D 1.000




FIGURE 2

			× 12 × 12 × 12 × 12 × 12 × 12 × 12 × 12	1	
		10	20	30	40
1	MAE				SUKKL HIELK48.aa
1	м G M				SUKLL MRC-OX44.aa GUAVM PETA-3.aa
1	GEFNEKE	TTCGTVCLE	YLLFTYNC	CrwbsA	<u>G: 84 v (13</u> 1121 3.00
			60	70	<del></del>
		50 	- <del></del>	1	V S G I I  HIEDK48.aa
15			L : I L G F G I	1000   1   2   2   2	TYGIL MRC-OX44.aa
10 36	KYVLFFF	1 F L F W V C G C C	A V G	IWTLALKS	DYISL PETA-3.aa
20					<del></del>
		90	100	110	120
29	TVGLGTGG	KCGGIAISLTI	VIGLSSA	LLHVGNLCLV	M G C I T HTEDK48.aa
44 44	FRNLPF -	LIT L G   F	10歳のけんほうっ	1 1 11	V V MRC-OX44.aa V T PETA-3.aa
52	LASGTY -	LLA TAL	ZIIVVAGT	/ <u>[V M]</u>	<u>V: 1 FEIA-</u> 3.00
			140	150	160
		130	140	1	
69	VLLGCAG	WYGATRESR	GTULFCLL	SMVIVLIMEVT	LAILL MRC-OX44.aa
67 75	A F L G C M G	TFRERR	NLIRLYFI	LLLILLARVT	AGILA PETA-3.aa
,,	O(43)#ii.8ii.#ii. 4 71				<del></del>
		170	180	190	200 L
109	LLFFPIV	GDVALEHTF	VT BRKNY -	RGYNEP DDY	STQWN HIELK48.aa
104				QHYHSDNST KRYHQPGHEAV	
112	YAYYQQL	MTELIKE	- N葉 K DI M T	K K HH M P G H L A V	10
		210	220	230	240
		210			MTTGH HTEDK48.aa
146	LVMEKLK	C C G V N N Y T D C C G V N G S S D	FSGSS WISGP		MRC-OX44.aa
137 148	FIQSQLQ QLQQEFH	CCGSMNSO	WRDSEWIR	SQEAGGRVV	PETA-3.aa
2.0	2 E E E - C				
		250	260	270	280
174	TYPRSCC	KSIGSVSCD	GRDVSPNV	IHQKG-CFHKI	LKITK HIELK48.aa
158			PSG	ADVQG - QYKKQ	GQAWFH MRC-OX44.aa LETFIQ PETA-3.aa
181	P D S C C	KTVVAL - CG	QRDHASMI	I K V Egg G gg I I iss -	
		200	300	310	320
		290	981 -1 28	2  0 :	AGLELL HIELK48.aa
213	TQSFTLS	GSSLGAAVI	VILSMSFA	LTLNCQIDI	KTSQAL MRC-OX44.aa
218 78(	SEHLRVIG	AVGIGIACV	VEGMIFT	CC	RSLKLE PETA-3.aa
24!	5 A .				HTEDK48.aa MRC-OX44.aa
21	B G L				PETA-3.aa
25	2 HY				<del></del>

FIGURE 3


					- 47		eost	-	-	_	T 10		_									-			; 20				-										30														40											
1	М	- A			C	F	_	. ! _			<u>-</u> -			- :		7.3	• •	•	20	- 7	•	•	ιτ	71	3.1	т		7	-	17		1 .	٠.		1.4	٠.		1 7	ì		ш		<b>7</b> : •					12)		- 6	٠,		DTT		•	•	_	_	-					
	M	-	-	E	T		Ē	7	<u>,                                     </u>	I	Т	C	: :	L	ĸ	1	<u>'</u>		L	_]		I	<u> </u>	ر ا -	s	F	_	<u>V</u>	F	V	V.	I	1	<u> </u>	ې	7	7	I	L	<u>.</u>		P	\ <u> </u>	¥.	G		<u> </u>	W)	G	K	(L	<u>L</u> ) —	T	-	T	A	LI	A	-1		<b>a</b> .a	ł,		
											50	)													60								_						7					_							_		80											
36 38	GO	A G				K				G S	P	I	-	S P	EG 10	5	3 .	A. -	M	ς.																																	Ċ											
39	Ĺ	G	T	Y	I	s	I	_] :	I	A	Ε	-	-	N	S	-	-	-	_		-	_		Γ	N		<u></u>	P	Y		7	1			G	_	Г	G	_	r	T	U	_	V —	<u>\</u>		<u>_</u>	G	_ <u>_</u>		Ξ:	Ģ	¢	_	Ί.	<u>'</u> A	لىك	_A		١.	az	<b>a</b>		
											9(														10		_									_				LO L						-		_	_	_	-1		120			<u>.</u> .			~ .	_	_	_		
76 73	Y L	G G	A	K	T	E		5	K K	C	P	. 1																																																				
73	F	A	Т	С	R		;	5	P	M	ľ	[	اند	K	L	, '	ΥĹ	A	M	1_	F	£	ļ	S	Ĭ	L	V	F	<u></u> I		A			L.	V	' L	<u>A</u>	G	; <u> </u>	IJ	S	_	3 i	F	Ĭ	<u></u>	F	R	Н		E (	1	К		.1	ئىل	ХĽ	-يما	<u>.</u>	L.	a	a		
											13														14															50 50			_	_		_				_	_		16	_					_	_				
116 113		F			· I			<b>→ I</b>	$\sim$	$\sim$	. 7	`	₹ .	77	~~	- 1	_	τ.	- 1	4	Ι.			1 -		ι.	1 1			v	\/	٠,		L	1 1		LV		٠.	**1	_		_	-	٠,	~	•	_	1 -	1 .	• •					N	₹W	-2	2	aa	ı			
113		T		-			-[	F	Ĺ	R		r	Y	Т	ב		A	M	(	2	Т			N	[_(	3_	N	ב	) ]	Ξ	-		-	R	5	<u>:</u> ]	R	P	_	V		) ]	H 	V	. (	2	R _	s	I	؛ <u>ل</u> ه	s _	Ç	¢	  -	7	П	AI.	1.2	<i>ī</i>	1.	.a	a		
							_		_	_			_			_	_	_					_											_					~	,	`												20	10								•		
											1														18													_		90			_			_			_			_		ــــ						_		-		
151		F	· 1		1 1	,	C	D	-	F			D V	s	E	 -	Y -	F	· ]	К	E		N T					1	1	P D	F		7-367	C	1	1	D	1		•		<del>-</del> -	N -	T -	•	-	S	Ē		5	¢	T	K		!	N	AC	-	2.	a	3			
151 149 147		F V	7 5	: h			r r	D D N	- W	F			D V T	s -	F	?	Y - Y	F Y F	1	K N	E A E		T H					14	1	P D P	F 9		T.) - (-7 - (-1	CCC		I L M	DEN	1 :		•		<del>-</del>	- -	7		-	N S T	Ē	:   5	5	¢	T	K		!	N	PW AC AI	-	2.	a	3			
151 149 147		F V	7 5			X		D D	- W -	F	v		_	s s	F	0.		_					_	SRO	2	A V I	F -		_	_			_		_		_		N F E	V +30	ר ר כ	-	-	_	-	-	S	Ē	:   5	5	¢	r 1	24	L < 1 P 1 40	- ! ! -	N T	AC AI	L	2 . A-	a 1	a .a	ıa	-	
190	-	) F		A F	H 1	D ·	Q	K	V	· E	2	E E S 10	_				_	_		  -	G	;		SRO	2	A V II - CC IN E	F   Q	1 2		L	7		D	I		- R	TE		ZEE	V T3(A L	T C	- - - - -		7 7	· - - /	G	S T	<u> </u>	)	-	-	r 2	24 24	1 C 1 C 1 G F	- 1	N T	AC AI PV AC	i-i L	2. A- E2	1 1 25	a .a	aa aa		
	-			A F	H 1	D ·	Q	K	V	· E	2	E E S 10	_				_	_		  -	G	;		SRO	2	A V II - CC IN E	F   Q	1 2		L	7		D	I		- R	TE		ZEE	V T3(A L	T C	- - - - -		7 7	· - - /	G	S T	<u> </u>	)	-	-	r 2	24 24	1 C 1 G	- 1	N T	AC AI PV AC	i-i L	2. A- E2	1 1 25	a .a	ıa		
190 180	-	) F		A F	H 1	D ·	Q	K	V	· E	2 = 4	E E S 10	- T				_	_		  -	G	;		SFG	2	LAVI CONED -6	F - CTL	I	7 7	L K T	7 7 2		D W F	I I M		2 E	TET		A A A A A A A A A	-V +3(-A L M -77)	0 7 7	- - - - - - - - - - - - -	TAI	777	7 7 5	G G A	ST	E			- L	1 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	24 24 25 28	1 G F G F 8		TINH	AC AI PV PAC TAI	L A L	2. A 2. A	1 25 a	a .a .a	aa aa		
190 180 179	- C C C C C C C C C C C C C C C C C C C	) k		A (	H	D T	Q W L	K W T	VKV	F 2	2 2 4 4 2	E E S T IO A T SI E	- T	K	7	- - - -	- N	-		 K	GFG			SFG	2	LAVI CONED -6	F - CTL	I	7 7	L K T	7 7 2		D W F	I I M		2 E	TET		A A A A A A A A A	-V +3(-A L M -77)	0 7 7	- - - - - - - - - - - - -	TAI	777	7 7 5	G G A	ST	E	)		- L	1 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	24 24	1 G F G F 8		NT HNT		I A A A A A A A A A A A A A A A A A A A	2. A- E2 A- E2	1 5 a -1	a	aa aa		
190 180 179 218 213	- C C C C C C C C C C C C C C C C C C C	1 Q I		A (	H 1	T T D	Q W L G L	K W T	VKV	F 1	2 2 2 1	EES TO A TOET	T	K	7		- и А	-	1	 K	O F O			SFO	2	LAVI CONED -6	F QTI	I	7 7	L K T	7 7 2		D W F	I I M		2 E	TET		A A A A A A A A A	-V +3(-A L M -77)	0 7 7	- - - - - - - - - - - - -	TAI	777	7 7 5	G G A	ST	E			- L	1 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	24 24 25 28	1 G F G F 8		NT HNT		I A A A A A A A A A A A A A A A A A A A	2. A- E2 A- E2	1 5 a -1	a	aa aa		
190 180 179 218 213	- C C C C C C C C C C C C C C C C C C C	1 Q I		A (	H 1	T T D	Q W L G L	K W T	VKV	F 1	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	EES TO A TOET	T	K	7		- и А	-	1	- K	O F O			SFO	2	LAVI CONED FOI	F O CTI	I	7 7	L K T	7 7 2		D W F	I I M		2 E	TET		A A A A A A A A A	-V +3(-A L M -77)	0 7 7	- - - - - - - - - - - - -	TAI	777	7 7 5	G G A	ST	E			- L	1 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	24 24 25 28	1 G F G F 8		TO HAT HAT	PAN AN A	A SI	2. A- E2. A-	1 25 a-1 25	a	aa aa		
190 180 179 218 213		1 Q I		A ()	H I	TI	QWLGLA	KWT F	VKV	F 1	2 E A A 2 L L Q	LEES LOL - A LOLET L LOL	T	K	. T	V - M M	ATL		1	- K NYA	O F O	(1)		SFO	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	LAVI COLNED FOIL S FOIR	F - CTI	7	H	LKT V	7 5		D W F	I I M		2 E	TET		A A A A A A A A A	-V +3(-A L M -77)	0 7 7	- - - - - - - - - - - - -	TAI	777	7 7 5	G G A	ST	E			- L	1 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	24 24 25 28	1 G F G F 8		NT HAT HAT		A SI WG	2.A- E2.A E.2	1 25 -1 25 -1 25	a	aa aa	ı	


## FIGURE 4

-	10	20	30	40
1 1	M N S M T S A V P V A M S V L M T T P R N S V N G T E	V V A P H N G Y P (	TPGIMSHVPI IA	YPNSQ HTPEF86.aa
± ;		60		<del></del>
41	FQVHLVPGNPPSLVS 	NVNGQPVQK		GAIQI HTPEF86.aa
23	MQSGKPEFF	RMSSLVGPT (	OSFF <u>MRESEE</u>	——————————————————————————————————————
	90	100	110	120
78 58	IIGU ABIGIGSIMAN MNGUFBIA UGGLLM	P-AGIYAPI	CVTVWYRLWG	G I M Y I I Bl.aa
	130	140	150	160
118	SCSLSV AAENQPYSY SCSLLAATEKNSRK-	CLLSGSLGL CLVKGKMIM	N IVSAICSAV N SLSLFAAIS	GVILFI HTPEF86.aa G <u>M</u> ILSI B1.aa
97		<del></del>	190	200
158	TELSIPH MEILNIKISHFLKM	180 		•
136	MOTLNIKISHFLKM	ESLNFIRAH <u>T</u>	PRINIR NCE-	PA報题SE BL.aa
	210	220	230	240 F
182 175	KNSPSTQYCYSIQS	MAISGVLU LFLGILSVMU	VEC LE IHAFFQE VI	AGIVEN Bl.aa
	250	260	270	280
199	ACASSHFGCQ	V C C Q S S N V S L L S A E E K K E Q	VIYPNIYAAN TIEIKE	P V I T P E HIPEF86.aa E V V G L T Bl.aa
215	EWKRT SRPKSMIV	ELSAFE K K ELO		<del></del> _
	290	300	310	320 HIPEF86.aa
23 <b>5</b> 251	- 1 1939 1939 - 1 - 2 - 2   - 3/29	IPIQEEEEEE	TETNFPEPPQ	DQESSP Bl.aa
			•	
248 291	1 1			HIPEF86.aa Bl.aa


# FIGURE 5

	10	20	30	40
1	MASPSRRLQT MET	SPVITCFKSVL SPVITCLKTLL	LIYTGI PWITGV LIXSEVEWITGV	III AVCI HSBBF02.aa III AVC <u>V</u> TALLA-1.aa
-		60	70	<del></del>
41	50 WGKVSEENXF	STUNEKATNVE	FVLIATGTVLIL	LGTFGCF HSBBF02.aa
34	WGKLTEGTYI	SLIAENSENAE	YVLLGTGTTLVV	FÖLEGGE TALIA-1.aa
	90	100	110	120
81. 74	ATCRASAGML ATCRGSPWML	ELYAMFLTLVF ELYAMFLSLVF	LVELVAAIVGFV L <u>A</u> BLVAGISGFV	方 R H E I K N HSHRF02.aa I R H E I K D TALLA-1.aa
	130	140	150	160 ·
121		volue e melny se	HAVEKTONTTHE	C G V T D Y R HSBBF02.aa
114	TILRITEM	QTTHGNDE-RE	RAVDHVORSISC	ONET TALLA-1.aa
	170		190	200
161 153	DMTDTNYYSE NWSTSPYFLE	K C F P K S C C K L E H C I P P S C C M N E	- D C Т P Q R Т D C N P Q D L H N L Т	- DADKVII HSBHF02.aa VAATKVII TALLA-1.aa
	210		230	<del></del>
194			ISFGVACFQLIG VARGIAFSQLIG	IFIAYCI HSPBF02.aa ML ACCI TALLA-1.aa
193	OKE YDL NTS	FMTTNKSII ———	<u>△</u> V V 製 製 T 製 L 2 M H H H K	M Liming Cissing Prints 1.00
	250			HSBBF02.aa
234 233				TALLA-1.aa





	10	20	30	40
1	M G R F R C C L R C L K Y L L C F M P V - K C G T K C L K Y L L E G E	M L L P N L A	GSAVIAFGLW GIAVLAIGLW	FREGIGI HALLOS9.aa LREDS CD9 antigen.aa
1	脚 P V - [K]與 T K   與   與   與   與   與   與   與   與   與	Hart - Harrison		<del></del>
	50	60	70 	80
41	AIKELSSEDKSPEYEY	VCLYVLV	GAGALMMAVG	F F C C C HAIDQ59.aa F L C C C CD9 antigen.aa
40	Q TX SIFEQET N N N N S S F S	T M. VIII. LIEF L		1011-1011111111111111111111111111111111
	90	100	110	120
79	GAMRZSQCVLGSFFTCLI	UIFAABV	TTGVFAFIGK	G V A I R HAIDQ59.aa
80	GAMRESQCVIGSFFICLI GAVQBSQCMIGLEFGFUI	制数制制数字 T 验 T	A A A I W G I S II A	
	130	140	150	160
119	HIZTMEEAYDDYLKDRO	KGN-GTL	ITFHSTFQCC	G K E S S HAIDQ59.aa G L A G G CD9 antigen.aa
120		DEPORE	K A(LEX X) A(D N) Sign	
	170	180	190	200
158		LGHKNCI	EIETIISVKI	QLIGIHAIDQ59.aa HIIGACD9 antigen.aa
159	VEQFISDISENKDVLET	FTVKSCPP	ANKEVEDNE	In Line A Coy and general
	210	220	230	
192	SECTION SECRET SERVICES	VLCCAIRI	ISRDVI.	HAIDQ59.aa CD9 antigen.aa
199		LLCCAIRF	RIN REMV	CD9 andgen.aa






FIGURE 11




FIGURE 12

		20	30	40
1 1 1	10 1 S P R L E - 1 S S F S T Q T P Y P N L A - M M L S L N N	V V V LQNIIYNPV		G G Q V HJACE54.aa D S K S rGalectin-5.aa D G T L hGalectin-8.aa
-	50	. 60	70	80
23 31 33	I I V R G L V L Q E P K H F T I V I S G V V L S D A K R E Q I V I R G H V P S D A D R F Q	V S L R D Q A A I N L R C - G G V D L Q N - G S S I	MKPRADVAFHFN	HJACE54.aa rGalectin-5.aa IPRFK hGalectin-8.aa
-	90	100	110	120
46 53 72	RAGCIVCNTLINEKW	GREEITYDT	PFQKEKKSFEI	HJACE54.aa rGalectin-5.aa / I M V L hGalectin-8.aa
	130	140	150	160
46 53 112	KAKFQVAVNGKHTLL	 	IDTLGIYGKVN	HJACE54.aa rGalectin-5.aa I H S I G hGalectin-8.aa
	170	180	190	200
46 53 152	FSFSSDLQSTQASSI	ELTEISREN	VPKSGTPQLRL	HJACE54.aa rGalectin-5.aa P F A A R hGalectin-8.aa
	210	220	230	240
46 53 192	LNTPMGPGRTVVVK	G E V N A N A K S F		A P V T L HJACE54.aa I A F H L rGalectin-5.aa I A L H L hGalectin-8.aa
	250	260	270	280
52 59 232	NPRFDENAVVRNTQ	19211221	LISAPFLEYPO SLPGSMPESRG RNITSFPESPG	III Riiki SIVI EGALECCAL Jian
	290	300	310	320
87 99 271	LLLFQEGGLELALN WILCEGHCFEVAVD IIYCDVREFEVAVN	G Q G L G A T S M N G Q H I C E Y S H F G V H S L E Y K H F	QQALEQLRELR LMNLPDINTLE RFKELSSIDTLE	IS S S V HJACE54.aa V A S D I rGalectin-5.aa IN S D I hGalectin-8.aa
139	330 Q L Y C V H S . Q L T H V E T H L L E V R S W			HJACE54.aa rGalectin-5.aa hGalectin-8.aa

FIGURE 13






FIGURE 14





## PCT

#### VORLD INTELLECTUAL PROPERTY ORGANIZ. International Bureau



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6: WO 98/31799 (11) International Publication Number: A3C12N 15/12, 15/85, 5/10, C07K 14/705, (43) International Publication Date: 23 July 1998 (23.07.98) 16/28, C12Q 1/68, A61K 38/17, 48/00 (21) International Application Number: PCT/US98/00959 (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW. HU, ID, IL, IS. JP, KE, KG, KP, KR, KZ, (22) International Filing Date: 21 January 1998 (21.01.98) LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, (30) Priority Data: TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO 60/034,204 21 January 1997 (21.01.97) US patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian 60/034,205 21 January 1997 (21.01.97) patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, (71) Applicant (for all designated States except US): HUMAN CM, GA, GN, ML, MR, NE, SN, TD, TG). GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US). Published (72) Inventors; and With international search report. (75) Inventors/Applicants (for US only): NI, Jian [CN/US]; 5502 Before the expiration of the time limit for amending the claims Manorfield Road, Rockville, MD 20853 (US). GENTZ,

Rolling Hill Road, Laytonsville, MD 20882 (US). (74) Agents: BROOKES, A., Anders et al.; Human Genome Sciences, Inc., 9410 Key West Avenue, Rockville, MD

Reiner, L. [DE/US]; 13404 Fairland Park Drive, Silver Spring, MD 20904 (US). ROSEN, Craig, A. [US/US]; 22400

and to be republished in the event of the receipt of amendments.

(88) Date of publication of the international search report: 5 November 1998 (05.11.98)

## (54) Title: POLYNUCLEOTIDES AND POLYPEPTIDES ENCODING RECEPTORS

#### (57) Abstract

20850 (US).

Receptor polypeptides and polynucleotides and methods for producing such polypeptides by recombinant techniques are disclosed. Also disclosed are methods for utilizing receptor polypeptides and polynucleotides in the design of protocols for the treatment of diseases and diagnostic assays for such conditions.

## FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Annenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
ΑU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
ΑZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Тодо
BB	Barbados	GH	Ghana	MG	Madagascar	ТJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
ВJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	<b>Iceland</b>	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		_
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denniark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		
							•


## INTERNATIONAL SEARCH REPORT

International Application No

A. CLASSI IPC 6	FICATION OF SUBJECT MATTER C12N15/12 C12N15/85 C12N5/1 C12Q1/68 A61K38/17 A61K48/		C07K16/28			
According to	o International Patent Classification (IPC) or to both national classific	ation and IPC				
B. FIELDS	SEARCHED					
Minimum as IPC 6	ocumentation searched (classification system followed by classificati C12N C07K C12Q A61K	on symbols)				
Documenta	tion searched other than minimum documentation to the extent that s	uch documents are included in the	fields searcned			
Electronic d	iata base consuited during the international search (name of data ba	se and, where practical, search ter	ms used)			
C. DOCUM	ENTS CONSIDERED TO BE RELEVANT					
Category 3	Citation of document, with indication, where appropriate, of the relation	evant passages	Relevant to claim No.			
X	DATABASE GENBANK Accession No. T48852, 8 February "EST70319 from Stratagene place XP002066898 see the whole document	1995 nta"	1-12,20			
X	DATABASE GENBANK Accession No. G14442, 4 January "STS SHGC-10055" XP000206689 see the whole document	1-3,20				
		-/				
X Furth	ner documents are listed in the continuation of box C.	Patent family members a	re listed in annex.			
**X° document defining the general state of the art which is not considered to be of particular relevance  "E" earlier document but published on or after the international filing date  "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)  "O" document referring to an oral disclosure, use, exhibition or other means  "P" document published prior to the international filing date but later than the priority date claimed  "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention  "X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered no						
Date of the a	actual completion of the international search	Date of mailing of the internati	•			
4	June 1998	1 6. 0	9. 1998			
Name and n	nailing address of the ISA  European Patent Office, P.B. 5818 Patentiaan 2  NL - 2280 HV Rijswijk  Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Authorized officer  Galli, I				

Form PCT/ISA/210 (second sheet) (July 1992)

## INTERNATIONAL SEARCH REPORT




C (Continu	ation) DOCUMENTS CONSIDERED TO BE RELEVANT	P 35 98/00959
Category °		Relevant to claim No.
A	ZANNETTINO A.C.W. ET AL.: "A powerful technique for isolating genes encoding surface antigens using retroviral expression cloning" J. IMMUNOL., vol. 156, no. 2, 15 January 1996, pages 611-620, XP002066893 see abstract	1-21
A	ADAMS M D ET AL: "COMPEMENTARY DNA SEQUENCING: EXPRESSED SEQUENCE TAGS AND HUMAN GENOME PROJECT" SCIENCE, vol. 252, no. 5013, 21 June 1991, pages 1651-1656, XP000645049 cited in the application see abstract	1-21
A	SIMMONS D. & SEED B.: "Isolation of a cDNA encoding CD33, a differentiation antigen of myeloid progenitor cells" J. IMMUNOLOGY, vol. 141, no. 8, 15 October 1988, pages 2797-2800, XP002066894 see abstract see figure 4	1-21
<b>A</b>	CROCKER P.R. ET AL.: "Sialoadhesin, a macrophage sialic acid binding receptor for haemopoietic cells with 17 immunoglobulin-like domains" EMBO J., vol. 13, no. 19, 3 October 1994, pages 4490-4503, XP002066895 see abstract see figure 2	1-21
Α	KELM S. ET AL.: "The Sialoadhesins: a family of sialic acid-dependent cellular recognition molecules within the immunoglobulin superfamily" GLYCOCONJ. J., vol. 13, no. 6, December 1996, pages 913-926, XPOO2066896 see the whole document	1-21
A	DATABASE GENBANK Accession No. U71382, 12 November 1996 PATEL N. ET AL.: "OB binding protein - 1" XP002066901 cited in the application see the whole document	1-21
	-/	

## INTERNATIONAL SEARCH REPORT

T/US 98/00959

		T/05 98/00959							
	Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT								
ategory °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.							
<b>,</b> X	DATABASE GENBANK Accession No. AA344713, 21 April 1997 "EST50650, H. sapiens cDNA similar to CD33 antigen" XP002066900 see the whole document	1-12,20							
,Α	TAKEI Y. ET AL.: "Molecular cloning of a novel gene similar to myeloid antigen CD33 and its specific expression in placenta" CYTOGENET. CELL GENET., vol. 78, 1997, pages 295-300, XP002066897 see abstract see figure 1	1-21							
	-								
		j 							
		·							
	•	·							
		·							

Form PCT/ISA/210 (continuation of second sneet) (July 1992)



Box I	Observati ns where certain claims were found unsearchable (Continuation of item 1 of first sheet)
This Inte	rnational Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:
1. <b>X</b>	Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely:  Remark: Although claims 13 and 14 are directed to a method of treatment of the human/animal body, and claim 15 (as far as in vivo methods are envisaged) is directed to a diagnostic method, the search has been carried out and based on the alleged effects of the compound/composition.
2.	Claims Nos.: because they relate to parts of the international Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3.	Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).
D= 41	
BOX II	Observations where unity of invention is lacking (Continuation of item 2 of first sheet)
	ernational Searching Authority found multiple inventions in this international application, as follows:  e further information sheet .
· **;	
1.	As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2.	As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3.	As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. X	No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:  see further information sheet, subject 1.
Remark	The additional search fees were accompanied by the applicant's protest.  No protest accompanied the payment of additional search fees.

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: (1-21) - partial

An isolated polynucleotide encoding the HMACR70 cell surface receptor (Seq. IDs 1,18), polynucleotides complementary thereto, fragments thereof, homologs thereof.

An expression system comprising said polynucleotides, a process to produce a cell harbouring said expression system. Said cell or membrane thereof expressing said receptor.

A process for the production of the HMACR70 polypeptide comprising culturing said cell.

Said HMACR70 polypeptide or homologs. An antibody directed against said polypeptide.

Therapeutic and diagnostic applications.

A method to identify agonists or antagonists to said receptors. Said agonists and antagonists.

2. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 2,3,19

3. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 4,20

4. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 5,21

5. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 6,22

6. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 7,23

7. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 8,24

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

- 8. Claims: (1-21) partial

  Idem as subject matter 1 but limited to Seq. IDs 9,10,25
- 9. Claims: (1-21) partial

  Idem as subject matter 1 but limited to Seq. IDs 11,26

- 13. Claims: (1-21) partial
   Idem as subject matter 1 but limited to Seq. IDs 15,30

## CORRECTED **VERSION\***

## PCT

#### WORLD INTELLECTUAL PROPERTY ORGAN International Bureau



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6:

C12N 15/12, 15/85, 5/10, C07K 14/705, 16/28, C12Q 1/68, A61K 38/17, 48/00

(11) International Publication Number:

WO 98/31799

(43) International Publication Date:

23 July 1998 (23.07.98)

(21) International Application Number:

PCT/US98/00959

A3

(22) International Filing Date:

21 January 1998 (21.01.98)

(30) Priority Data:

21 January 1997 (21.01.97) 60/034,204 60/034,205 21 January 1997 (21.01.97) US US

(71) Applicant (for all designated States except US): HUMAN GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): NI, Jian [CN/US]; 5502 Manorfield Road, Rockville, MD 20853 (US), GENTZ, Reiner, L. [DE/US]; 13404 Fairland Park Drive, Silver Spring, MD 20904 (US). ROSEN, Craig, A. [US/US]; 22400 Rolling Hill Road, Laytonsville, MD 20882 (US).

(74) Agents: BROOKES, A., Anders et al.; Human Genome Sciences, Inc., 9410 Key West Avenue, Rockville, MD 20850 (US).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL. TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

#### Published

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(88) Date of publication of the international search report: 5 November 1998 (05.11.98)

(54) Title: POLYNUCLEOTIDES AND POLYPEPTIDES ENCODING RECEPTORS

(57) Abstract

Receptor polypeptides and polynucleotides and methods for producing such polypeptides by recombinant techniques are disclosed. Also disclosed are methods for utilizing receptor polypeptides and polynucleotides in the design of protocols for the treatment of diseases and diagnostic assays for such conditions.

## FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

			-				
AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
ΑT	Austria	FR	France	LU	Luxembourg	SN	Senegal
ΑU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	ТJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	ΙE	Ircland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL.	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Vict Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		•
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	· LR	Liberia	SG	Singapore		

WO 98/31799 PCT/US98/00959

## POLYNUCLEOTIDES AND POLYPEPTIDES ENCODING RECEPTORS

## FIELD OF INVENTION

5

10

15

20

25

30

35

This invention relates to newly identified polynucleotides and the polypeptides encoded by them, the use of such polynucleotides and polypeptides, and their production. More particularly, the polynucleotides and polypeptides of the present invention relate to specific receptor families described in the specification and known in the art. The invention also relates to inhibiting or activating the action of such polynucleotides and polypeptides.

## BACKGROUND OF THE INVENTION

Receptor proteins are found on the membrane of the cells and are generally involved in signal transduction. There are many types of receptor proteins, and for convenience, these proteins are grouped in families based on similarity in structure and function.

For example, the TM4SF superfamily of cell surface proteins, also known as the tetraspan receptor superfamily, is comprised of at least seventeen individual gene products (these include CD9, CD20, CD37, CD53, CD63, CD81, CD82, A15, CO-029, Sm23, RDS, Uro B, Uro A, SAS, Rom-1, PETA3, and YKK8). The TM4SF superfamily is the second largest group in the CD antigen superfamily. Each member of the TM4SF superfamily can be characterized by several putative physical features including four highly conserved transmembrane domains, two divergent extracellular loops, and two short and highly divergent cytoplasmic tails. Expression patterns for members of the TM4SF superfamily tend to be rather broad and can vary widely between members. The functional roles of TM4SF superfamily members are primarily associated with signal transduction events and pathways, but also include cell adhesion in platelets and other lymphocytic and non-lymphocytic cell lines, as well as cell motility, proliferation, and metastasis. In addition, recent evidence suggests that a subset of the members of the TM4SF superfamily may function as potassium channel molecules.

One member of the TM4SF family, CD20, is a four membrane spanning domain cell surface phosphoprotein expressed exclusively on B lymphocytes. Although the precise functional role of CD20 has yet to be determined, it is thought to function primarily as a receptor during B-cell activation. Furthermore, a large number of experimental observations suggest several additional speculative roles for the CD20 molecule. For example, CD20-specific immunoprecipitation of biochemically cross-linked plasma membrane proteins suggests that CD20 assumes a multimeric structural

WO 98/31799 PCT/US98/00959

conformation characteristic of other previously described membrane channel proteins. Further experimentation has revealed that expression of exogenous CD20 on the cell surface specifically increases Ca<sup>2+</sup> conductance across the plasma membrane. Together, these results suggest that CD20 complexes may function as B-cell specific Ca<sup>2+</sup> ion channels. In addition, monoclonal antibodies raised against CD20 have been used to stimulate resting B-cells to transition out of the G0/G1 segment of the cell cycle. It has also been demonstrated that CD20 is associated with both serine and tyrosine kinases and, more specifically, that CD20 is associated, although not directly, with the Src family of tyrosine kinases including p56/53lyn, p56lck, and p59fyn.

A second example of a receptor subfamily, called sialoadhesin molecules, belongs to the Ig superfamily of receptor-like molecules. The more than 100 members of the Ig superfamily are generally considered to engage in specific cell-cell interactions through which intercellular communication may occur. In addition to classical protein-protein interactions, intercellular communication may also be mediated through protein-carbohydrate interactions. In fact, all members of the sialoadhesin family of the Ig superfamily are capable of mediating protein-sialic acid binding interactions. To date, only a small number of proteins have been assigned to the sialoadhesin family including sialoadhesin, CD33, CD22, the myelin-associated glycoprotein (MAG), and the Schwann cell myelin protein (SMP). Each of these proteins is expressed in a restricted subset of cell types. For example, CD22 and CD33 are expressed exclusively by B-lymphocytes and cells of the myelomonocytic lineage, respectively.

Similarly, galectins are a family of the lectin superfamily of carbohydrate-binding proteins which have a high affinity for b-galactoside sugars. Although a large number of glycoproteins containing b-galactoside sugars are produced by the cell, only a few will bind to known galectins *in vitro*. Such apparent binding specificity suggests a highly specific functional role for the galectins. Galectin 1 (conventionally termed *LGALS1* for lectin, galactoside-binding, soluble  $\underline{-1}$ ) is thought to specifically bind laminin, a highly polylactosaminated cellular glycoprotein, as well as the highly polylactosaminated lysosome-associated membrane proteins (LAMPs). Galectin 1 has also been shown to bind specifically to a lactosamine-containing glycolipid found on olfactory neurons and to integrin  $a_7b_1$  on skeletal muscle cells. Galectin 3 has also been observed to bind specifically to laminin, immunoglobulin E and its receptor, and bacterial lipopolysaccharides.

Various galectins have been shown to function in the mechanisms of intercellular communication. For example, depending on cell type, galectin 1 has been observed to modulate cell adhesion either positively or negatively. More specifically, galectin 1 appears to inhibit cell adhesion of skeletal muscle presumably by galectin 1 mediated disruption of laminin-integrin  $a_7b_1$  interactions. Alternatively, galectin 1 appears to promote cell adhesion in several non-skeletal muscle cell types examined

5

10

15

20

25

30

WO 98/31799 3 PCT/US98/00959

presumably by a glycoconjugate cross-linking mechanism. Galectin 3 has also been observed to function in modulating cell-adhesion, as well as in the activation of certain immune cells by cross-linking IgE and IgE receptors. In addition, galectins have been observed to be involved in the regulation of immune cell activity, as well as in such diverse processes as cell adhesion, proliferation, inflammation, autoimmunity, and metastasis of tumor cells. Furthermore, a galectin-like antigen designated HOM-HD-21 was recently found to be highly expressed in a Hodgkin's Disease cDNA library. Very recently, a novel galectin, termed PCTA-1, was identified as a specific cell surface marker on human prostate cancer cell lines and patient-derived carcinomas. Galectins have also been found to function intracellularly as a component of ribonucleoprotein complexes. Finally, galectins 1 and 3 have each been found to modulate T-cell growth and apoptosis by interaction with CD45 and possibly Bc12, respectively.

A relatively new family of cell-surface proteins has been identified and termed the Ly6 superfamily. The members of this family include murine and human SCA-2, rat Ly-6 (also termed ThB), human CD59 [also known as protectin or membrane attack complex inhibition factor (MACIF)], and E48 antigen. The determination of an initial functional role for SCA-2 may lie in an analysis of its expression profile with regard to the complex process of hematopoiesis. SCA-2 is highly expressed in early thymic precusor cells. In turn, progeny of the intrathymic precusor population continue to express SCA-2, but only until the point of transition occurs from blast cell to small cell. Further experimental evidence demonstrates that mature thymocytes and peripheral T-cells do not express SCA-2. As a result, it seems very likely that SCA-2 plays an important role in thymocyte maturation and differentiation. A plausible explanation for this functional hypothesis is that SCA-2 may act as a receptor for a unknown cytokine which regulates thymocyte maturation and differentiation.

In addition, CD59 is a recently identified integral membrane protein which appears to be involved in the regulation of complement. Recent studies show that the CD59 antigen may prevent damage from complement C5b-9 and protect astrocytes during inflammatory and infectious disorders of the nervous system. Expression of recombinant human CD59 on porcine donor organs have been shown to prevent complement-mediated lysis and activation of endothelial cells that leads to hyperacute rejection. Recently, researchers at Alexion Pharmaceuticals (New Haven, CT) reported on the production of transgenic pigs which expressed human CD59. In these animals, xenogeneic organs were resistant to hyperacute rejection. (Fodor, et al., "Expression of a functional human complement inhibitor in a transgenic pig as a model for the prevention of xenogeneic hyperacute organ rejection," Proc. Natl. Acad. Sci., 91:1153-11157 (1994).) The same company also reported that expression of recombinant transmembrane CD59 in paroxysmal nocturnal hemoglobinuria (PNH) B-

5

10

15

20

25

30

WO 98/31799 4 PCT/US98/00959

cells confers resistance to human complement. (Rother et al., "Expression of recombinant transmembrane CD59 in paroxysmal nocturnal hemoglobinuria B-cells confers resistance to human complement," Blood, 84:2604-2611 (1994).) PNH is an acquired hematopoietic disorder characterized by complement-mediated hemolytic anemia, pancytopenia, and venous thrombosis. It is thought that retroviral gene therapy with this molecule could provide a treatment for PNH patients.

A final Ly6 superfamily member, the E48 antigen, is involved in intercellular adhesion between keratinocyte cells of the squamous epithelium. Such keratinocytes are attached to adjoining cells by large numbers of desmosomes, which are thought to play a role in the transition of transformed keratinocytes to metastatic tumor cells. Treatment with a monoclonal antibody raised against the E48 antigen has been successful in the eradication of residual, postoperative squamous cell carcinoma cells of the upper aerodigestive tract in several *in vivo* models and, to some degree, in humans. (van Dongen, et al., "Progress in radioimmunotherapy of head and neck cancer," Oncol. Rep. 1:259-264 (1994).) The gene encoding the E48 antigen has been mapped to the q24-qter region of human chromosome 8. Interestingly, a number of human diseases have been mapped to this region of chromosome 8 including Langer-Giedion syndrome, brachio-otorhinolaryngeal syndrome, trichorhinolaryngeal syndrome, and epidermolysis bullosa simplex.

A further example of a receptor family includes the prohibitin receptors. The prohibitin gene product is expressed in a wide variety of tissues and has been implicated as a component of a number of anti-proliferative mechanisms. The prohibitin gene encodes a 30 kD postsynthetically modified polypeptide located primarily in the mitochondria, but also may be associated with the IgM receptor on the B-cell plasma membrane. The protein functionally inhibits DNA synthesis and entry into S phase of the cell cycle by an unknown mechanism. Interestingly, although the prohibitin gene product is hypothesized to be involved in the maintenance of senescence and the prevention of cancer, one study found that, although somatic mutations in the prohibitin gene were present in a small number of breast cancers, no mutations were identified in any other breast, ovary, liver, and lung cancers examined. (Sato et al., Genomics 17:762-764 (1993).) However, the prohibitin gene has been mapped to human chromosome 17q12-21, the same region thought to contain the gene involved in sporadic breast cancer. Furthermore, DNA sequence analysis of the prohibitin gene identified somatic mutation in 4 of 23 cases of sporadic breast cancer examined. Thus, prohibitin family members may be involved in the development of cancer.

Moreover, the EGFR family of plasma membrane proteins are an integral component of normal cellular proliferation and in the pathogenesis of the cancerous state. The family is relatively small and includes the EGFR, c-erbB-2, c-erbB-3, and others. Various cancers are correlated with aberrant expression of one or more of these

10

15

20

25

30

5

genes. A number of ligands have been identified which bind to the EGFR-like receptors listed above including TGF-a, heparin-binding EGF, amphiregulin, criptoregulin, heregulin, and others. A large fraction of adenocarcinomas examined to date, especially those of the breast, colon, and pancreas, are typified by the amplification or overexpression of the c-erbB-2 gene. EGF, or an analogous ligand, initiates the cellular growth factor response by binding to the EGFR, or EGFR-related, receptor. Following the binding event, the receptor molecule dimerizes activating its intracellular tyrosine kinase domain. This event results in the phosphorylation of specific tyrosine residues near the carboxy terminus of the receptor. The diversity of signals able to be transduced through the relatively small number of EGFR-related receptor molecules is amplified considerably by the recent finding that EGFR-like receptor molecules can function when dimerized with other EGFR family members forming heterodimers.

Members of the EGFR-related family of integral membrane proteins have been implicated in the pathogenesis of a number of human disease-states. For example, a mutation in the EGFR itself appears to play an important role in the development of glioblastomas. (Sang et al., J. Neurosurg 82:841-846 (1995).) The EGFR gene is amplified or overexpressed in the majority of primary human glioblastomas. Although not conferring a distinct advantage on cell growth, an increase in EGFR expression was found to confer an increase in the ability of glioma cells to maintain anchorage-independent growth in soft agar especially in response to EGF and retinoic acid.

Anchorage-independent growth *in vitro* correlates highly with tumorigenicity *in vivo*, therefore, it is likely that cells which express abnormally high levels of EGFR in human glioblastoma cells may be involved in the high potential for these cells to cause tumors *in vivo*.

Moreover, overexpression or amplification of c-erbB-2 has been reported to be involved in a high number adenocarcinomas, particularly of the breast, colon, and pancreas, and in a small proportion of ovarian carcinomas.

Thus, there is a clear need for identifying and exploiting novel members of the receptor families, such as those described above. Although structurally related, these receptors will likely possess diverse and multifaceted functions in a variety of cell and tissue types. Receptor type molecules should prove useful in target based screens for small molecules and other such pharmacologically valuable factors. Monoclonal antibodies raised against such receptors may prove useful as therapeutics in an antitumor, diagnostic, or other capacity. Furthermore, receptors described here may prove useful in an active or passive immunotherapeutical role in patients with cancer or other immunocompromised disease states.

5

10

15

20

25

30

WO 98/31799 6 PCT/US98/00959

## SUMMARY OF THE INVENTION

5

10

15

20

25

30

35

In one aspect, the invention relates to receptor polypeptides and polynucleotides, as well as the methods for their production. Another aspect of the invention relates to methods for using such receptor polypeptides and polynucleotides. Such uses include the treatment of the specified diseases, among others. In still another aspect, the invention relates to methods to identify agonists and antagonists using the materials provided by the invention, and treating conditions associated with receptor imbalance with the identified compounds. Yet another aspect of the invention relates to diagnostic assays for detecting diseases associated with inappropriate receptor activity or levels.

# DESCRIPTION OF THE INVENTION Definitions

The following definitions are provided to facilitate understanding of certain terms used frequently herein.

"Receptor" refers, among others, to a polypeptide comprising the amino acid sequence set forth in SEQ ID NO:Y, or an allelic variant thereof.

"Receptor Activity" or "Biological Activity of the Receptor" refers to the metabolic or physiologic function of said receptor including similar activities or improved activities or these activities with decreased undesirable side-effects. Also included are antigenic and immunogenic activities of said receptor.

"Receptor gene" refers to a polynucleotide comprising the nucleotide sequence set forth in SEQ ID NO:X or allelic variants thereof and/or their complements.

"SEQ ID NO:X" comprises all or a substantial portion of the polynucleotide encoding each receptor of the invention. The value X for the nucleotide sequence is an integer specified in Table 1. This nucleotide sequence was translated into the receptor polypeptide identified in Table 1 as "SEQ ID NO:Y," where the value of Y for each receptor polypeptide is an integer defined in Table 1.

The invention further provides a composition of matter comprising a nucleic acid molecule which comprises a human cDNA clone identified by a cDNA Clone ID (Identifier) in Table 1, which DNA molecule is contained in the material deposited with the American Type Culture Collection ("ATCC") and given the ATCC Deposit Number shown in Table 1 for that cDNA clone. The ATCC is located at American Type Culture Collection (ATCC), 12301 Park Lawn Drive, Rockville, Maryland 20852, USA. The deposit has been made under the terms of the Budapest Treaty on the international recognition of the deposit of micro-organisms for purposes of patent procedure. The strain will be irrevocably and without restriction or condition released to the public upon the issuance of a patent. The deposit is provided merely as convenience to those of skill in the art and is not an admission that a deposit is required for enablement, such as that

WO 98/31799 7 PCT/US98/00959

required under 35 U.S.C. §112. The nucleotide sequence of the polynucleotides contained in the deposited material, as well as the amino acid sequence of the polypeptide encoded thereby, are controlling in the event of any conflict with any description of sequences herein.

"Antibodies" as used herein includes polyclonal and monoclonal antibodies, chimeric, single chain, and humanized antibodies, as well as Fab fragments, including the products of an Fab or other immunoglobulin expression library.

"Isolated" means altered "by the hand of man" from the natural state. If an "isolated" composition or substance occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living animal is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein.

"Polynucleotide" generally refers to any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotides" include, without limitation single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, "polynucleotide" refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term polynucleotide also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications has been made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. "Polynucleotide" also embraces relatively short polynucleotides, often referred to as oligonucleotides.

"Polypeptide" refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres. "Polypeptide" refers to both short chains, commonly referred to as peptides, oligopeptides or oligomers, and to longer chains, generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene-encoded amino acids. "Polypeptides" include amino acid sequences modified either by natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications can

5

10

15

20

25

30

WO 98/31799 8 PCT/US98/00959

occur anywhere in a polypeptide, including the peptide backbone, the amino acid sidechains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched and branched cyclic polypeptides may result from posttranslation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphotidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. (See, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York, 1993 and Wold, F., Posttranslational Protein Modifications: Perspectives and Prospects, pgs. 1-12 in POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic Press, New York, 1983; Seifter et al., "Analysis for protein modifications and nonprotein cofactors", Meth Enzymol (1990) 182:626-646 and Rattan et al., "Protein Synthesis: Posttranslational Modifications and Aging", Ann NY Acad Sci (1992) 663:48-62.)

"Variant" as the term is used herein, is a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide respectively, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to

5

10

15

20

25

30

5

10

15

20

occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis.

"Identity" is a measure of the identity of nucleotide sequences or amino acid sequences. In general, the sequences are aligned so that the highest order match is obtained. "Identity" per se has an art-recognized meaning and can be calculated using published techniques. (See, e.g.: COMPUTATIONAL MOLECULAR BIOLOGY, Lesk, A.M., ed., Oxford University Press, New York, 1988; BIOCOMPUTING: INFORMATICS AND GENOME PROJECTS, Smith, D.W., ed., Academic Press, New York, 1993; COMPUTER ANALYSIS OF SEQUENCE DATA, PART I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; SEQUENCE ANALYSIS IN MOLECULAR BIOLOGY, von Heinje, G., Academic Press, 1987; and SEQUENCE ANALYSIS PRIMER, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991.) While there exist a number of methods to measure identity between two polynucleotide or polypeptide sequences, the term "identity" is well known to skilled artisans. (Carillo, H., and Lipton, D., SIAM J Applied Math (1988) 48:1073.) Methods commonly employed to determine identity or similarity between two sequences include, but are not limited to, those disclosed in Guide to Huge Computers, Martin J. Bishop, ed., Academic Press, San Diego, 1994, and Carillo, H., and Lipton, D., SIAM J Applied Math (1988) 48:1073. Methods to determine identity and similarity are codified in computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, GCS program package (Devereux, J., et al., Nucleic \_\_\_ Acids Research (1984) 12(1):387), BLASTP, BLASTN, FASTA (Atschul, S.F. et al.,

25

30

35

J Molec Biol (1990) 215:403.)

As an illustration, by a polynucleotide having a nucleotide sequence having at least, for example, 95% "identity" to a reference nucleotide sequence of SEQ ID NO:X is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence of SEQ ID NO: X. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5 ' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence.

Similarly, by a polypeptide having an amino acid sequence having at least, for example, 95% "identity" to a reference amino acid sequence of SEQ ID NO:Y is intended that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid of SEQ ID NO:Y. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

15

20

25

10

5

## Polypeptides of the Invention

In one aspect, the present invention relates to receptor polypeptides (or receptor proteins). The receptor polypeptides include the polypeptide of SEQ ID NO:Y; as well as polypeptides comprising the amino acid sequence of SEQ ID NO:Y; and polypeptides comprising the amino acid sequence which have at least 80% identity to that of SEQ ID NO:Y over its entire length, and still more preferably at least 90% identity, and even still more preferably at least 95% identity to SEQ ID NO:Y. Furthermore, those with at least 97-99% identity to SEQ ID NO:Y are highly preferred. Also included within receptor polypeptides are polypeptides having the amino acid sequence which have at least 80% identity to the polypeptide having the amino acid sequence of SEQ ID NO:Y over its entire length, and still more preferably at least 90% identity, and even still more preferably at least 95% identity to SEQ ID NO:Y. Furthermore, those with at least 97-99% are highly preferred. Preferably receptor polypeptides exhibit at least one biological activity of the receptor.

30

The receptor polypeptides may be in the form of the "mature" protein or may be a part of a larger protein such as a fusion protein. It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, prosequences, sequences which aid in purification such as multiple histidine residues, or an additional sequence for stability during recombinant production.

35

Fragments of the receptor polypeptides are also included in the invention. A "fragment" is a polypeptide having an amino acid sequence that entirely is the same as part, but not all, of the amino acid sequence of the aforementioned receptor polypeptides. As with receptor polypeptides, fragments may be "free-standing," or comprised within a larger polypeptide of which they form a part or region, most

WO 98/31799 PCT/US98/00959

П

preferably as a single continuous region. Representative examples of polypeptide fragments of the invention, include, for example, fragments from about amino acid number 1-20, 21-40, 41-60, 61-80, 81-100, and 101 to the end of receptor polypeptide. In this context "about" includes the particularly recited ranges larger or smaller by several, 5, 4, 3, 2 or 1 amino acid at either extreme or at both extremes.

Preferred fragments include, for example, truncation polypeptides having the amino acid sequence of receptor polypeptides, except for deletion of a continuous series of residues that includes the amino terminus, or a continuous series of residues that includes the carboxyl terminus or deletion of two continuous series of residues. one including the amino terminus and one including the carboxyl terminus.

Also preferred are fragments characterized by structural or functional domains, such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions. The "domains" of each receptor polypeptide are illustrated in the Figures. The Figures compare SEQ ID NO:Y to the closest know homologue. Identical amino acids shared between the two polypeptides are shaded, while conservative amino acid changes are boxed. By examining the regions or amino acids shaded and/or boxed, the skilled artisan can readily identify conserved domains between the two polypeptides. The amino acids sequences of SEQ ID NO:Y falling within these conserved domains are "fragments" and are specifically contemplated by the present invention. Especially preferred is the extracellular domains of a receptor of the invention. Soluble extracellular domains have antagonist activity mediated by competition with a receptor ligand.

Other preferred fragments are biologically active fragments. Biologically active fragments are those that mediate receptor activity, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also included are those that are antigenic or immunogenic in an animal, especially in a human.

Preferably, all of these polypeptide fragments retain a biological activity of the receptor, including antigenic activity. Variants of the defined sequence and fragments also form part of the present invention. Preferred variants are those that vary from the referents by conservative amino acid substitutions -- i.e., those that substitute a residue with another of like characteristics. Typical such substitutions are among Ala, Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and Glu; among Asn and Gln; and among the basic residues Lys and Arg; or aromatic residues Phe and Tyr. Particularly preferred are variants in which several, 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination.

5

10

15

20

25

30

WO 98/31799 12 PCT/US98/00959

The receptor polypeptides of the invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

## Polynucleotides of the Invention

Another aspect of the invention relates to receptor polynucleotides. Receptor polynucleotides include isolated polynucleotides which encode the receptor polypeptides and fragments, and polynucleotides closely related thereto. More specifically, a receptor polynucleotide of the invention includes a polynucleotide comprising the nucleotide sequence contained in SEQ ID NO:X encoding a receptor polypeptide of SEQ ID NO:Y, and polynucleotide having the particular sequence of SEQ ID NO:X.

15

20

25

30

10

5

Receptor polynucleotides further include a polynucleotide comprising a nucleotide sequence that has at least 80% identity over its entire length to a nucleotide sequence encoding the receptor polypeptide of SEQ ID NO:Y, and a polynucleotide comprising a nucleotide sequence that is at least 80% identical to that of SEO ID NO:X over its entire length. In this regard, polynucleotides at least 90% identical are particularly preferred, and those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred and those with at least 98-99% are most highly preferred, with at least 99% being the most preferred. Also included under receptor polynucleotides are a nucleotide sequence which has sufficient identity to a nucleotide sequence contained in SEQ ID NO:X, or contained in the cDNA insert in the plasmid deposited with ATCC, to hybridize under conditions useable for amplification or for use as a probe or marker. Moreover, the receptor polynucleotide includes a nucleotide sequence having at least 80% identity to a nucleotide sequence encoding the receptor polypeptide expressed by the cDNA insert deposited at the ATCC, and a nucleotide sequence comprising at least 15 contiguous nucleotides of such cDNA insert. In this regard, polynucleotides at least 90% identical are particularly preferred, and those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred and those with at least 98-99% are most highly preferred, with at least 99% being the most preferred. The invention also provides polynucleotides which are complementary to all the above receptor polynucleotides.

35

The receptors of the invention are structurally related to other proteins of specified receptor families, as shown by the results in the Figures. The cDNA sequence of SEQ ID NO:X encodes a polypeptide as described in Table 1 as SEQ ID NO:Y. Because the receptor polypeptides contain domains similar in structure to other receptor family members, the receptors of the present invention are expected to have,

inter alia, similar biological functions/properties to their homologous polypeptides and polynucleotides, and their utility is obvious to anyone skilled in the art.

Table 1

5

Clone ID Name	SEQ ID NO:X	SEQ ID NO:Y	ATCC Deposit No.	ATCC Deposit Date	Receptor Family	Homology
HMACR70			209054	05/16/97	Ig	Sialoadhesin
	1	18	#####	01/21/98	_	OB-1
HTEDK48			209054	05/16/97	TM4SF	MRC-OX44 PETA-3
1-1849 bp	2					
160-900 bp	3	19				
HTPED39			209054	05/16/97	TM4SF	NAG-2
HPWAE25	4	20	#####	1/21/98		TALLA-1
HTPEF86	5	21	209053	05/16/97	TM4SF	. CD20
						B1 Antigen
HSBBF02	6	22	209054	05/16/97	TM4SF	TALLA-1
HLTAH80	7	23	97242	08/02/95	TM4SF	TALLA-I
			209054	05/16/97		
HTPBA27	8	24	97242	08/02/95	TM4SF	NAG-2
			209054	05/16/97		
HAIDQ59			209054	05/16/97	TM4SF	CD9
						Antigen
5' Sequence	9	25		<u></u>	-	
3' Sequence	10					
HHFEK40	11	26	209054	05/16/97	TM4SF	PETA-3
HGBGV89	12	27	209125	06/09/97	TM4SF	L6H
			209054	05/16/97		
HUVBB80	13	28	209054	05/16/97	TM4SF	L6
HJACE54	14	29	209053	05/16/97	Lectin	Galectin-3 Galectin-5
			<u> </u>			Galectin-8
HROAD63	15	30	209053	05/16/97	Ly6	E48 splice
						variant
HMWGS46	16	31	209053	05/16/97	Prohibitin	BAP-37
HNFGW06	17	32	209053	05/16/97	EGFR	EGFR

The novel full-length cDNA clone designated HMACR70 may be a member of the sialoadhesin family of the Ig superfamily of receptor-like molecules and a CD33 homologue. HMACR70 contains a 1497 nucleotide cDNA insert encoding a 315 amino acid ORF and was cloned from a GM-CSF-treated human macrophage cDNA library. The only additional cDNA libraries in the HGS database which include this clone are human eosinophils and possibly human gall bladder. A BLAST analysis of the amino acid sequence of HMACR70 demonstrates that this clone exhibits approximately 50% identity and 69% similarity over a 300 amino acids stretch of a gene termed human

10

- 15

WO 98/31799 PCT/US98/00959

differentiation antigen, and 38% identity and 62% similarity of the human myelin-associated glycoprotein precursor CD33 gene.

A more recent BLAST analysis confirms HMACR70's designation as a sialoadhesin family member. HMACR70 is homologous to two recently identified sialoadhesin family members, human OB binding protein (OB) 1 and 2. (See, Genbank Accession No. U71382; see Figure 1.) It is thought that OB-1 and OB-2 may bind leptin. Thus, HMACR70, as a sialoadhesin family member, may act to attenuate or even amplify intercellular routes of communication, including binding to leptin or modulating the activity of immune cells, such as macrophages. Clearly, any diseases affected by these processes could be treated by the polypeptide or fragment of HMACR70.

The full-length nucleotide sequences of ten novel human cDNA clones which potentially belong to the TM4SF superfamily are disclosed in the table above and will be addressed sequentially.

The cDNA clone **HTEDK48** contains a 1849 nucleotide cDNA insert encoding a 245 amino acid ORF that was cloned from a human testes cDNA library. The coding sequence of HTEDK48 (SEQ ID NO: 3) may be fused to other human proteins, such as 3-hydroxyacyl-CoA dehydrogenase. BLAST analysis of the amino acid sequence of HTEDK48 demonstrates that this clone exhibits approximately 30% identity and 51% similarity over a 245 amino acid stretch of the CD82 molecule. Recent studies have shown that CD82 can associate with CD4 or CD8 and deliver costimulatory signals for the TCR/CD3 pathway. CD82 has also been found to be involved in syncytium formation in HTLV-I-infected T-cells. And finally, in a recently published study in which the expression of the CD82 gene by tumors of the lung was examined retrospectively, it was reported that CD82 may be linked to the suppression of tumor metastasis of prostate cancer. The study also reported that decreased CD82 expression may be involved in malignant progression of such cancers. Thus, HTEDK48 may also be involved in the development of cancer.

A more recent BLAST analysis shows that HTEDK48 is homologous the rat leukocyte antigen, MRC OX-44, and the platelet endothelial tetraspan antigen -3 (PETA-3). (See Figure 2X.) MRC OX-44, a member of a new family of cell surface proteins, appears to be involved in growth regulation. (See, Bellacosa, A., et al., "The Rat Leukocyte antigen MRC OX-44 is a Member of a New Family of Cell Surface Proteins which Appear to be Involved in Growth Regulation," Mol. Cell. Bio. 11: 2864-2872 (1991).) Similarly, PETA-3 has been located to platelet endothelial cells, and an anti-PETA-3 antigen monoclonal antibody can stimulate platelet aggregation and mediator release. (See, Fitter, S., "Molecular Cloning of cDNA Encoding a Novel Platelet-Endothelial Cell Tetra-Span Antigen, PETA-3," Blood; 86(4):1348-1355 (1995).) Thus, HTEDK48 may function similar to MRC OX-44 or PETA-3 to affect

10

15

20

25

30

WO 98/31799 15 PCT/US98/00959

growth of blood cells. Administering polypeptides or fragments of HTEDK48 may be an effective treatment of blood disorders.

The cDNA clone **HPWAE25** contains a 1288 nucleotide cDNA insert encoding a 273 amino acid ORF that was cloned from a human pancreas tumor cDNA library, while clone **HTPED39** represents a truncated cDNA sequence. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including keratinocytes, ulcerative colitis, striatum depression, lymph node breast cancer, ovarian cancer, stage B2 prostate cancer, kidney medulla, and others. Northern blot analysis of HLTAH80 also shows expression in a variety of human cell lines including U937, MM96, WM115, and MDAMB231. A BLAST analysis of the amino acid sequence of HTPED39 demonstrates that this clone exhibits approximately 35% identity and 50% similarity over the entire length of the CD37 molecule. The CD37 antigen is expressed on B cells and on a subpopulation of T cells, but not on pre-B or plasma cells. It has been reported that CD37 expression is downregulated in conjunction with B-cell activation, suggesting that CD37 may be involved in the processes which dictate the activation state of the B-cell.

Moreover, HPWAE25 is also homologous to recently identified TM4SF members, NAG-2 and TALLA-1. (See Figure 3.) NAG-2 is thought to complex with integrins and other TM4SF proteins, while TALLA-1 is a highly specific marker of T-cell acute lymphoblastic leukemia and neuroblastoma. (See, Tachibana, I., et al., "NAG-2, A Novel Transmembrane-4 Superfamily (TM4SF) Protein that Complexs with Integrins and Other TM4SF Proteins," J. Biol. Chem., 272:29181-29189 (1997); Takagi, S., "Identification of a Higly Specific Surface Marker of T-cell Acute Lymphoblastic Leukemia and Neuroblastoma as a New Member of the Transmembrane 4 Superfamily," Int. J. Cancer 61(5):706-715 (1995).) Thus, HPWAE25 may be involved the development of cancer, particularly leukemia, lymphoma, and neuroblastoma. HPWAE25 may be used as an effective treatment of these cancers, as well as a diagnostic marker.

A subfamily of TM4SF receptors include CD20 proteins. A CD20-like cDNA clone was obtained from a human pancreas tumor cDNA library and contains a 1236 nucleotide insert which encodes a 250 amino acid ORF. A BLAST analysis of the deduced amino acid sequence of HTPEF86 exhibits approximately 41% identity and 61% similarity to the CD20 gene, also known as B1 antigen. (See Figure 4.) Expression of this gene is detected in only two additional HGS human cDNA libraries; amygdala depression and 9 week early stage human. Although the precise functional role of CD20 has yet to be determined, it is clear that CD20 plays a key role in the regulation of B-cell activation. Based primarily on sequence identity, the novel CD20-like molecule presented herein may also be involved in cell cycle activation. Potential therapeutic and/or diagnostic applications for HTPEF86 may include such clinical

5

10

15

20

25

30

WO 98/31799 16 PCT/US98/00959

presentations as juvenile rheumatoid arthritis, Graves' Disease, and a number of B-cell lymphomas or other lymphoid tumors.

The clone **HSBBF02** contains a 1115 nucleotide cDNA insert encoding a 245 amino acid ORF and was cloned from an HSC 172 cell line cDNA library. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including brain amygdala depression, endothelial cells, fetal liver and heart, osteoblasts, testes, and others. A BLAST analysis of the amino acid sequence of HSBBF02 demonstrates that this clone exhibits approximately 64% identity and 80% similarity with the A15 molecule over a 131 amino acid stretch (A15 is composed of 244 amino acids). A more recent BLAST search shows that HSBBF02 is similar to the TALLA-1 protein and may in fact be a closely related family member. (See Figure 5.)

In addition, a second cDNA clone, designated **HLTAH80**, exhibits sequence similarity to the A15 molecule and TALLA-1. (See Figure 6.) This clone contains a 1662 nucleotide cDNA insert encoding a 253 amino acid ORF and was cloned from a human T-cell lymphoma cDNA library. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including B-cell lymphoma, corpus collosum, endometrial tumor, osteosarcoma, testes, and others. Northern blot analysis of HLTAH80 also shows expression in a variety of human tissues including spleen, lymph node, thymus, PBLs, heart, and a particularly strong signal in skeletal muscle and pancreas. A BLAST analysis of the amino acid sequence of HLTAH80 demonstrates that this clone exhibits approximately 35% identity and 55% similarity over the entire length of the A15 molecule.

Since expression of A15 drops to undetectable levels when comparing immature T-cells to peripheral blood lymphocytes, it is thought that A15 may play a role in the development of T-cells. Furthermore, the MXS1(CCG-B7) gene which codes for A15 contains a number of triplet nucleotide repeats which have been associated with neuropsychiatric diseases such as Huntington's chorea, fragile X syndrome, and myotonic dystrophy. In addition, A15 appears to be expressed exclusively on T-cell acute lymphoblastic leukemia cell lines, including several derived from adult T-cell leukemia and those established by immortalization with human T-cell leukemia virus type 1 or Herpesvirus saimiri. Thus, clones HLTAH80 and/or HSBBF02 may also be involved in diseases caused by the expansion of repeats or chromosomal instability.

The cDNA clone **HTPBA27** contains a 1345 nucleotide cDNA insert encoding a 238 amino acid ORF and was cloned from a human tumor pancreas cDNA library. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including cerebellum, breast lymph node, osteosarcoma, adult testes, RS4;11 bone marrow cell line, microvascular endothelial cells, and others. A BLAST analysis of the amino acid sequence of HTPBA27 demonstrates that this

5

10

15

20

25

30

WO 98/31799 PCT/US98/00959

clone exhibits approximately 40% identity and 64% similarity with a glycoprotein termed CD53 over its entire length. CD53 is thought to be involved in thymopoiesis. since rat CD53 can be detected on immature CD4-8-thymocytes and the functionally mature single-positive subset, but cannot be detected on the intermediate CD4+8+ thymocytic subset of cells. The CD53 molecule has also been implicated as a component of signal transduction pathways in B cells, monocytes and granulocytes, rat macrophages, NK, and T cells. Moreover, as illustrated in Figure 7, HTPBA27 was recently confirmed as a TM4SF receptor. (See, Tachibana, I., et al., "NAG-2, A Novel Transmembrane-4 Superfamily (TM4SF) Protein that with Integrins and Other TM4SF Proteins," J. Biol. Chem., 272:29181-29189 (1997).) Calling the HTPBA27 polypeptide NAG-2, this group confirmed HTPBA27's status as a TM4SF receptor by showing that NAG-2 complexes with integrin and other TM4SF receptors. Thus, diseases caused by the failure of HTPBA27 to complex with integrin and other TM4SF receptors can be treated by administering HTPBA27. HTPBA27 can also be used to diagnose these diseases.

The cDNA clone **HAIDQ59** contains cDNA insert encoding a 221 amino acid ORF that was cloned from a human epithelial cell induced with TNFa and INF cDNA library. The 5' end of HAIDQ59 is represented by the SEQ ID NO: 9, while the 3' end is represented by SEQ ID NO: 10. This clone appears in only two additional cDNA libraries in the HGS database. These two libraries were constructed from the human Jurkat T-cell line and human microvascular endothelial cells. A BLAST analysis of the amino acid sequence of HAIDQ59 demonstrates that this clone exhibits approximately 53% identity and 69% similarity over 226 amino acids of the CD9 TM4SF molecule. (See Figure 8.) It has been demonstrated that the CD9 molecule is involved in signal transduction pathways in platelets, as well as in cell adhesion in both platelets and pre-B-cell lines. Intriguingly, a monoclonal antibody (vpg15), which recognizes the feline homologue of CD9, has been shown to block infection by feline immunodeficiency virus (FIV). Furthermore, a recent study shows that cells expressing high levels of CD9 exhibited suppressed cell motility. Thus, HAIDQ59 may also be involved in signal transduction of blood cells.

The cDNA clone **HHFEK40** contains a 936 nucleotide cDNA insert encoding a 252 amino acid ORF and was cloned from a human fetal heart cDNA library. This clone appears once in the human fetal heart cDNA library and possibly in a hemangiopericytoma cDNA library. A BLAST analysis of the amino acid sequence of HHFEK40 demonstrated that this clone exhibits approximately 60% identity and 75% similarity over the entire length of a molecule designated PETA-3. (See Figure 9.) PETA-3 was originally identified as a novel human platelet surface glycoprotein termed gp27. Although PETA-3 is present in low abundance on the platelet surface, an anti-PETA-3 monoclonal antibody can stimulate platelet aggregation and mediator release.

5

10

15

20

25

30

WO 98/31799 18 PCT/US98/00959

Thus, HHFEK40 may function similar to PETA-3 to affect growth of blood cells. Administering polypeptides or fragments of HHFEK40 may be an effective treatment of blood disorders.

The cDNA clone HGBGV89 contains a 738 nucleotide cDNA insert encoding a 197 amino acid ORF and was cloned from a human gall bladder cDNA library. The only two additional appearances of this clone in the HGS database are in a normalized fetal liver cDNA library and in a fetal liver/spleen cDNA library. The cDNA clone HUVBB80 contains a 1071 nucleotide cDNA insert encoding a 201 amino acid ORF and was cloned from a human umbilical vein cDNA library. This clone appears in several additional cDNA libraries in the HGS database including prostate BPH, thyroid, and fetal liver/spleen. BLAST analyses of the amino acid sequences of HGBGV89 and HUVBB80 demonstrate that these clones exhibit approximately 49% identity and 65% similarity and 47% identity and 68% similarity, respectively, over the entire length of a molecule designated L6 surface protein or human tumor-associated antigen L6. (See Figures 10 & 11.) Moreover, another group has confirmed the TM4SF receptor homology of HGBGV89 by describing the protein as a putative transmembrane protein L6H. (See Genbank Accession No 2587054; see Figure 10.) The L6 cell surface antigen is highly expressed on lung, breast, colon, and ovarian carcinomas. Promising results of phase 1 clinical studies have been reported with an anti-L6 monoclonal antibody, or its humanized counterpart, suggesting that the L6 antigen may be an attractive target for monoclonal antibody-based cancer therapy.

In summary, there is a clear need for identifying and exploiting novel members of the TM4SF superfamily such as those described herein. Although structurally related, these factors will likely possess diverse and multifaceted functions in a variety of cell and tissue types. Receptor type molecules, such as the novel potential members of the TM4SF superfamily detailed here, should prove useful in target based screens for small molecules and other such pharmacologically valuable factors. Monoclonal antibodies raised against such factors may prove useful as therapeutics in an anti-tumor, diagnostic, or other capacity. Furthermore, factors such as the nine novel TM4SF superfamily-like molecules described here may prove useful in an active or passive immunotherapeutical role in patients with cancer or other immunocompromised disease states.

Besides TM4SF receptors, receptors from other families are also described. For example, clone **HJACE54**, also called galectin 11, exhibits significant sequence identity to the rat galectin 5, the chicken galectin 3 gene, and the human galectin 8 genes. (See Figure 12.) The galectin 11 cDNA clone contains an 865 nucleotide insert which encodes a 133 amino acid ORF. The clone was obtained from a Jurkat T-cell G1 phase cDNA library. A BLAST analysis of the deduced amino acid sequence of HJACE54 demonstrates approximately 35% identity and 57% similarity to the amino

5

10

15

20

25

30

WO 98/31799 - PCT/US98/00959

acid sequence of the rat galectin 5 gene. Expression of galectin 11 is quite limited in the HGS database. In fact, the only two additional ESTs in the HGS database which contain the HJACE54 sequence were found in human neutrophil and human infant adrenal gland cDNA libraries. Northern blot analyses have not been performed to examine expression patterns of the galectin 11 gene.

Various galectins have been shown to function in the mechanisms of intercellular communication. For example, depending on cell type, galectin 1 has been observed to modulate cell adhesion either positively or negatively. More specifically, galectin 1 appears to inhibit cell adhesion of skeletal muscle presumably by galectin 1mediated disruption of laminin-integrin a<sub>7</sub>b<sub>1</sub> interactions. Alternatively, galectin 1 appears to promote cell adhesion in several non-skeletal muscle cell types examined presumably by a glycoconjugate cross-linking mechanism. Galectin 3 has also been observed to function in modulating cell-adhesion, as well as in the activation of certain immune cells by cross-linking IgE and IgE receptors. In addition, galectins have been observed to be involved in the regulation of immune cell activity, as well as in such diverse processes as cell adhesion, proliferation, inflammation, autoimmunity, and metastasis of tumor cells. Furthermore, a galectin-like antigen designated HOM-HD-21 was recently found to be highly expressed in a Hodgkin's Disease cDNA library. Very \_ recently, a novel galectin, termed PCTA-1, was identified as a specific cell surface marker on human prostate cancer cell lines and patient-derived carcinomas. Galectins: have also been found to function intracellularly as a component of ribonucleoprotein complexes. Finally, galectins 1 and 3 have each been found to modulate T-cell growth. and apoptosis by interaction with CD45 and possibly Bcl2, respectively. As a result, the discovery of a novel galectin, such as that encoded by HJACE54, is likely to be a valuable asset both diagnostically and therapeutically.

Additionally, a full-length nucleotide sequence of a novel human cDNA clone which encodes an apparent splice variant of the previously described human E48 antigen has recently been determined. (See Figure 13.) Clone **HROAD63** contains a 441 nucleotide cDNA which encodes a 70 amino acid polypeptide. This novel clone exhibits significant sequence identity to several members of a relatively new family of cell-surface proteins termed the Ly6 superfamily. These members include murine and human SCA-2, rat Ly-6 (also termed ThB), and human CD59 [also known as protectin or membrane attack complex inhibition factor (MACIF)]. The novel E48 splice variant was obtained from the HGS human stomach cDNA library. The clone is present in only a limited number of other HGS cDNA libraries including kidney cancer, keratinocyte, and tongue. An alignment of the nucleotide sequences of the human E48 and HROAD63 cDNAs demonstrates that the initial 168 and 178 nucleotides of E48 and HROAD63, respectively, are identical, with the exception of an additional 10 nucleotides of sequence at the extreme 5' end of the HROAD63 sequence. The

5

10

15

20

25

30

WO 98/31799 20 PCT/US98/00959

sequence of the two clones is also identical for an additional 229 nucleotides including the 3' end of the coding sequences and the entire 3' untranslated regions. The only divergence of nucleotide sequence in this region of the clones is the deletion of a single thymidine residue in the 3' UTR of the E48 cDNA. The major difference between the two nucleotide sequences is a 329 nucleotide deletion from the HROAD63 sequence. This deletion causes a shift in the HROAD63 reading frame and encompasses the translational stop signal used in the E48 clone. As a result, the carboxy terminal sequence of HROAD63 is radically altered with regard to that of E48 (as illustrated in Figure 13 by the obvious differences between amino acids 56-128 of E48 and 56-70 of HROAD63 in the amino acid alignment). The clinical presentation of disorders, including abnormal skin and hair phenotypes, may be attributed, at least in part, to a non-functional Ly6 superfamily member such as E48 or HROAD63. HROAD63 may also be involved in blood disorders, as seen with its homologues SCA-2 and CD59.

A novel prohibitin cDNA clone presented herein was originally identified in a human bone marrow cell line (RS4;11) cDNA library. The clone contains a 1066 nucleotide insert which encodes a 299 amino acid polypeptide. BLAST and BestFit analyses of the predicted amino acid sequence of HMWGS46 demonstrate a highly significant sequence identity to a murine protein termed IgM B-cell receptor associated protein (BAP)-37 (Genbank accession number X78683). The HMWGS46 amino acid sequence exhibits nearly perfect identity and similarity over the entire length of the murine BAP-37 sequence. (See Figure 14.) In addition, the full-length nucleotide sequences of HMWGS46 and BAP-37 exhibit at least 87% identical. The HMWGS46 clone also exhibits approximately 49% sequence identity and 85% sequence similarity to a human gene designated prohibitin. Finally, the HMWGS46 cDNA appears in a substantial number of HGS human cDNA libraries in addition to the bone marrow cell line cDNA library from which it was cloned. Some of the cDNA libraries in which this clone appears include keratinocytes, induced endothelial cells, activated neutrophils, synovial sarcoma, colon carcinoma cell line, Jurkat cell line membrane bound polysomes, epileptic frontal cortex, primary dendritic cells, and a number of others. The novel gene related to prohibitin and BAP-37 may prove quite useful as a diagnostic for tumorigenesis, as well as a target for therapeutic intervention of such an event. Thus, although the precise functional role of the prohibitin family members are less than clear, it is quite likely that such homologues are involved in such complex processes as development, senescence, and tumor suppression. Therefore a novel gene, such as HMWGS46, may prove quite useful as a diagnostic for tumorigenesis, as well as a target for therapeutic intervention of such an event.

A human cDNA clone encoding a novel epidermal growth factor receptor (EGFR)-like molecule is also disclosed. The novel EGFR-like cDNA clone presented herein was originally identified in an activated human neutrophil cDNA library. The

5

10

15

20

25

30

clone contains a 704 nucleotide insert which encodes a 168 amino acid polypeptide. A BLAST analysis of the predicted amino acid sequence of **HNFGW06** demonstrates that this novel clone exhibits approximately 85% identity and 90% similarity to a protein designated epidermal growth factor receptor-related protein [Homo sapiens]. (See Figure 15.) The expression profile of the HNFGW06 clone in the HGS database indicates the existence of a fairly highly restricted expression pattern. In addition to the activated neutrophil library from which this clone was obtained, it also appears in the following HGS human cDNA libraries: synovial sarcoma, smooth muscle, placenta, and possibly primary dendritic cells.

The novel EGFR-like cDNA clone HNFGW06 may lead to a number of exciting possibilities for therapeutic and/or diagnostic treatments or reagents. For example, HNFGW06 may be involved in the onset of human breast cancers as well. In addition, due to the fact that TGF-a acts through binding to the EGFR, it is possible that HNFGW06 may also play a role in a variety of gastric processes including regulation of acid secretion, regulation of mucous cell growth, and protection against ethanol- and aspirin-induced injury to gastric tissues.

### GENERATING POLYNUCLEOTIDES

Polynucleotides of the present invention encoding a receptor may be obtained using standard cloning and screening, from a cDNA library derived from mRNA in cells specified in Table 1 using the expressed sequence tag (EST) analysis (Adams, M.D., et al. Science (1991) 252:1651-1656; Adams, M.D. et al., Nature, (1992) 355:632-634; Adams, M.D., et al., Nature (1995) 377 Supp:3-174.) Polynucleotides of the invention can also be obtained from natural sources such as genomic DNA libraries or can be synthesized using well known and commercially available techniques.

The nucleotide sequence encoding a receptor polypeptide of SEQ ID NO:Y may be identical to the polynucleotide encoding SEQ ID NO:Y, or it may be a sequence, which as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:Y.

When the polynucleotides of the invention are used for the recombinant production of a receptor polypeptide, the polynucleotide may include the coding sequence for the mature polypeptide or a fragment thereof, by itself; the coding sequence for the mature polypeptide or fragment in reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro- protein sequence, or other fusion peptide portions. For example, a marker sequence which facilitates purification of the fused polypeptide can be encoded. In certain preferred embodiments of this aspect of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in

5

10

15

20

25

30

WO 98/31799 PCT/US98/00959

22

Gentz et al., Proc Natl Acad Sci USA (1989) 86:821-824, or is an HA tag. The polynucleotide may also contain non-coding 5' and 3' sequences, such as transcribed, non-translated sequences, splicing and polyadenylation signals, ribosome binding sites and sequences that stabilize mRNA.

Further preferred embodiments are polynucleotides encoding receptor variants comprising the amino acid sequence of receptor polypeptide of Table 1 (SEQ ID NO:Y) in which several, 5-10, 1-5, 1-3, 1-2 or 1 amino acid residues are substituted, deleted or added, in any combination.

The present invention further relates to polynucleotides that hybridize to the herein above-described sequences. In this regard, the present invention especially relates to polynucleotides which hybridize under stringent conditions to the herein above-described polynucleotides. As herein used, the term "stringent conditions" means hybridization will occur only if there is at least 80%, and preferably at least 90%, and more preferably at least 95%, yet even more preferably 97-99% identity between the sequences.

Polynucleotides of the invention, which are identical or sufficiently identical to a nucleotide sequence contained in SEQ ID NO:X or a fragment thereof, or to the cDNA insert in the plasmid deposited at the ATCC, or a fragment thereof, may be used as hybridization probes for cDNA and genomic DNA, to isolate full-length cDNAs and genomic clones encoding the receptor and to isolate cDNA and genomic clones of other genes (including genes encoding homologs and orthologs) that have a high sequence similarity to the receptor gene. Such hybridization techniques are known to those of skill in the art. Typically these nucleotide sequences are 80% identical, preferably 90% identical, more preferably 95% identical to that of the referent. The probes generally will comprise at least 15 nucleotides. Preferably, such probes will have at least 30 nucleotides and may have at least 50 nucleotides. Particularly preferred probes will range between 30 and 50 nucleotides.

In one embodiment, to obtain a polynucleotide encoding the receptor polypeptide, including homologs and orthologs from other species, comprises the steps of screening an appropriate library under stringent hybridization conditions with a labeled probe having the SEQ ID NO:X or a fragment thereof; and isolating full-length cDNA and genomic clones containing said polynucleotide sequence. Such hybridization techniques are well known to those of skill in the art. Stringent hybridization conditions are as defined above or, alternatively, conditions under overnight incubation at 42°C in a solution comprising: 50% formamide, 5xSSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10 % dextran sulfate, and 20 microgram/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65°C.

5

10

15

20

25

30

WO 98/31799 23 PCT/US98/00959

The polynucleotides and polypeptides of the present invention may be employed as research reagents and materials for discovery of treatments and diagnostics to animal and human disease.

## 5 Vectors, Host Cells, Expression

The present invention also relates to vectors which comprise a polynucleotide or polynucleotides of the present invention, and host cells which are genetically engineered with vectors of the invention and to the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention.

For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof for polynucleotides of the present invention. Introduction of polynucleotides into host cells can be effected by methods described in many standard laboratory manuals, such as Davis et al., BASIC METHODS IN MOLECULAR BIOLOGY (1986) and Sambrook et al., MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989) such as calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction or infection.

Representative examples of appropriate hosts include bacterial cells, such as streptococci, staphylococci, E. coli, Streptomyces and Bacillus subtilis cells; fungal cells, such as yeast cells and Aspergillus cells; insect cells such as Drosophila S2 and Spodoptera Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, HEK 293 and Bowes melanoma cells; and plant cells.

A great variety of expression systems can be used. Such systems include, among others, chromosomal, episomal and virus-derived systems, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression systems may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides to produce a polypeptide in a host may be used. The appropriate nucleotide sequence may be inserted into an expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook et al., MOLECULAR CLONING, A LABORATORY MANUAL (supra).

10

15

20

25

30

WO 98/31799 24 PCT/US98/00959

For secretion of the translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the desired polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

If the receptor polypeptide is to be expressed for use in screening assays, generally, it is preferred that the polypeptide be produced at the surface of the cell. In this event, the cells may be harvested prior to use in the screening assay. If the receptor polypeptide is secreted into the medium, the medium can be recovered in order to recover and purify the polypeptide; if produced intracellularly, the cells must first be lysed before the polypeptide is recovered.

Receptor polypeptides can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during isolation and or purification.

# Diagnostic Assays

5

10

15

20

25

30

35

This invention also relates to the use of receptor polynucleotides or polypeptides for use as diagnostic reagents. Detection of a mutated form of the receptor gene associated with a dysfunction will provide a diagnostic tool that can add to or define a diagnosis of a disease or susceptibility to a disease which results from underexpression, over-expression or altered expression of the receptor. Individuals carrying mutations in the receptor gene may be detected at the DNA level by a variety of techniques.

Nucleic acids for diagnosis may be obtained from a subject's cells, such as from blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification techniques prior to analysis. RNA or cDNA may also be used in similar fashion. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to labeled receptor nucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in electrophoretic mobility of DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing. (See, e.g., Myers et al., Science (1985) 230:1242.) Sequence changes at specific locations may also be revealed by

WO 98/31799 25 PCT/US98/00959

nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method. (See Cotton et al., Proc Natl Acad Sci USA (1985) 85: 4397-4401.) In another embodiment, an array of oligonucleotides probes comprising receptor nucleotide sequence or fragments thereof can be constructed to conduct efficient screening of e.g., genetic mutations. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability. (See for example: M.Chee et al., Science, Vol 274, pp 610-613 (1996).)

The diagnostic assays offer a process for diagnosing or determining a susceptibility to specific diseases through detection of mutation in the receptor gene by the methods described.

In addition, specific diseases can be diagnosed by methods comprising determining from a sample derived from a subject an abnormally decreased or increased level of receptor polypeptide or receptor mRNA. Decreased or increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example, PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods. Assay techniques that can be used to determine levels of a protein in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

Thus in another aspect, the present invention relates to a diagnostic kit for a disease or susceptibility to a disease which comprises:

- (a) a receptor polynucleotide, preferably the nucleotide sequence of SEQ ID NO:X, or a fragment thereof;
  - (b) a nucleotide sequence complementary to that of (a);
- (c) a receptor polypeptide, preferably the polypeptide of SEQ ID NO:Y, or a fragment thereof; or
- (d) an antibody to a receptor polypeptide, preferably to the polypeptide of SEQ ID NO: Y.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

# Chromosome Assays

The nucleotide sequences of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome. The mapping of relevant sequences to chromosomes according to the present invention is an important first step in correlating those sequences with gene associated disease. Once a sequence has been mapped to a precise chromosomal location, the physical position of the

5

10

15

20

25

WO 98/31799 <sup>26</sup> PCT/US98/00959

sequence on the chromosome can be correlated with genetic map data. Such data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes).

The differences in the cDNA or genomic sequence between affected and unaffected individuals can also be determined. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

10

15

20

25

30

35

5

#### Antibodies

The polypeptides of the invention or their fragments or analogs thereof, or cells expressing them can also be used as immunogens to produce antibodies immunospecific for the receptor polypeptides. The term "immunospecific" means that the antibodies have substantially greater affinity for the polypeptides of the invention than their affinity for other related polypeptides in the prior art.

Antibodies generated against the receptor polypeptides can be obtained by administering the polypeptides or epitope-bearing fragments, analogs or cells to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler, G. and Milstein, C., Nature (1975) 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor et al., Immunology Today (1983) 4:72) and the EBV-hybridoma technique (Cole et al., MONOCLONAL ANTIBODIES AND CANCER THERAPY, pp. 77-96, Alan R. Liss, Inc., 1985).

Techniques for the production of single chain antibodies (U.S. Patent No. 4,946,778) can also be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms including other mammals, may be used to express humanized antibodies.

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or to purify the polypeptides by affinity chromatography. Antibodies against receptor polypeptides may also be employed to treat diseases.

## Vaccines

Another aspect of the invention relates to a method for inducing an immunological response in a mammal which comprises inoculating the mammal with a receptor polypeptide, or a fragment thereof, adequate to produce antibody and/or T cell immune response to protect said animal from a disease. Yet another aspect of the invention relates to a method of inducing immunological response in a mammal which

comprises, delivering a receptor polypeptide via a vector directing expression of the receptor polynucleotide in vivo in order to induce such an immunological response to produce antibody to protect said animal from diseases.

Further aspect of the invention relates to an immunological/vaccine formulation (composition) which, when introduced into a mammalian host, induces an immunological response in that mammal to a receptor polypeptide wherein the composition comprises a receptor polypeptide or receptor gene. The vaccine formulation may further comprise a suitable carrier. Since a receptor polypeptide may be broken down in the stomach, it is preferably administered parenterally (including subcutaneous, intramuscular, intravenous, intradermal etc. injection). Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation instonic with the blood of the recipient; and aqueous and nonaqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

#### Screening Assays

5

10

15

20

25

30

35

The receptor polypeptide of the present invention may be employed in a screening process for compounds which bind the receptor and which activate (agonists) or inhibit activation of (antagonists) the receptor polypeptide of the present invention. Thus, polypeptides of the invention may also be used to assess the binding of small molecule substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural substrates and ligands or may be structural or functional mimetics. See Coligan et al., Current Protocols in Immunology 1(2):Chapter 5 (1991).

The receptor polypeptides are responsible for many biological functions, including many pathologies. Accordingly, it is desirous to find compounds and drugs which stimulate the receptor on the one hand and which can inhibit the function of the receptor on the other hand. In general, agonists are employed for therapeutic and prophylactic purposes for such conditions and diseases. Antagonists may be employed for a variety of therapeutic and prophylactic purposes for such conditions and diseases.

In general, such screening procedures involve producing appropriate cells which express the receptor polypeptide of the present invention on the surface thereof.

10

15

20

25

30

35

Such cells include cells from mammals, yeast, Drosophila or E. coli. Cells expressing the receptor (or cell membrane containing the expressed receptor) are then contacted with a test compound to observe binding, or stimulation or inhibition of a functional response.

The assays may simply test binding of a candidate compound wherein adherence to the cells bearing the receptor is detected by means of a label directly or indirectly associated with the candidate compound or in an assay involving competition with a labeled competitor. Further, these assays may test whether the candidate compound results in a signal generated by activation of the receptor, using detection systems appropriate to the cells bearing the receptor at their surfaces. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed.

Further, the assays may simply comprise the steps of mixing a candidate compound with a solution containing a receptor polypeptide to form a mixture, measuring receptor activity in the mixture, and comparing the receptor activity of the mixture to a standard.

The receptor cDNA, protein and antibodies to the protein may also be used to configure assays for detecting the effect of added compounds on the production of receptor mRNA and protein in cells. For example, an ELISA may be constructed for measuring secreted or cell associated levels of receptor protein using monoclonal and polyclonal antibodies by standard methods known in the art, and this can be used to discover agents which may inhibit or enhance the production of the receptor (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues. Standard methods for conducting screening assays are well understood in the art.

Examples of potential receptor antagonists include antibodies or, in some cases, oligonucleotides or proteins which are closely related to the ligand of the receptor, e.g., a fragment of the ligand, or small molecules which bind to the receptor but do not elicit a response, so that the activity of the receptor is prevented.

Thus in another aspect, the present invention relates to a screening kit for identifying agonists, antagonists, ligands, receptors, substrates, enzymes, etc. for receptor polypeptides; or compounds which decrease or enhance the production of receptor, which comprises:

- (a) a receptor polypeptide, preferably that of SEQ ID NO:Y;
- (b) a recombinant cell expressing a receptor polypeptide, preferably that of SEQ ID NO:Y:
  - (c) a cell membrane expressing a receptor polypeptide; preferably that of SEQ ID NO: Y; or
    - (d) antibody to a receptor polypeptide, preferably that of SEQ ID NO: Y.

SUBSTITUTE SHEET (RULE 26)

WO 98/31799 29 PCT/US98/00959

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

# Prophylactic and Therapeutic Methods

5

10

15

20

25

30

35

This invention provides methods of treating an abnormal conditions related to both an excess of and insufficient amounts of receptor activity.

If the activity of the receptor is in excess, several approaches are available. One approach comprises administering to a subject an inhibitor compound (antagonist) as described along with a pharmaceutically acceptable carrier in an amount effective to inhibit activation by blocking the binding of ligands to the receptor or by inhibiting a second signal, and thereby alleviating the abnormal condition.

In another approach, soluble forms of the receptor polypeptides still capable of binding the ligand in competition with endogenous receptor may be administered. Typical embodiments of such competitors comprise fragments of the receptor polypeptide.

In still another approach, expression of the gene encoding endogenous receptor can be inhibited using expression blocking techniques. Known such techniques involve the use of antisense sequences, either internally generated or separately administered. (See, for example, O'Connor, J Neurochem (1991) 56:560 in Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988).) Alternatively, oligonucleotides which form triple helices with the gene can be supplied. (See, for example, Lee et al., Nucleic Acids Res (1979) 6:3073; Cooney et al., Science (1988) 241:456; Dervan et al., Science (1991) 251:1360.) These oligomers can be administered per se or the relevant oligomers can be expressed in vivo.

For treating abnormal conditions related to an under-expression of the receptor and its activity, several approaches are also available. One approach comprises administering to a subject a therapeutically effective amount of a compound which activates the receptor, i.e., an agonist as described above, in combination with a pharmaceutically acceptable carrier, to thereby alleviate the abnormal condition.

Alternatively, gene therapy may be employed to effect the endogenous production of the receptor by the relevant cells in the subject. For example, a polynucleotide of the invention may be engineered for expression in a replication defective retroviral vector, as discussed above. The retroviral expression construct may then be isolated and introduced into a packaging cell transduced with a retroviral plasmid vector containing RNA encoding a polypeptide of the present invention such that the packaging cell now produces infectious viral particles containing the gene of interest. These producer cells may be administered to a subject for engineering cells in vivo and expression of the polypeptide in vivo. For overview of gene therapy, see Chapter 20, Gene Therapy and

other Molecular Genetic-based Therapeutic Approaches, (and references cited therein) in Human Molecular Genetics, T Strachan and A P Read, BIOS Scientific Publishers Ltd (1996).

#### 5 Formulation and Administration

Peptides, such as the soluble form of receptor polypeptides, and agonists and antagonist peptides or small molecules, may be formulated in combination with a suitable pharmaceutical carrier. Such formulations comprise a therapeutically effective amount of the polypeptide or compound, and a pharmaceutically acceptable carrier or excipient. Such carriers include but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. Formulation should suit the mode of administration, and is well within the skill of the art. The invention further relates to pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

15

20

10

Polypeptides and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds. Preferred forms of systemic administration of the pharmaceutical compositions include injection, typically by intravenous injection. Other injection routes, such as subcutaneous, intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transmucosal and transdermal administration using penetrants such as bile salts or fusidic acids or other detergents. In addition, if properly formulated in enteric or encapsulated formulations, oral administration may also be possible. Administration of these compounds may also be topical and/or localized, in the form of salves, pastes, gels and the like.

25

30

The dosage range required depends on the choice of peptide, the route of administration, the nature of the formulation, the nature of the subject's condition, and the judgment of the attending practitioner. Suitable dosages, however, are in the range of 0.1-100 µg/kg of subject. Wide variations in the needed dosage, however, are to be expected in view of the variety of compounds available and the differing efficiencies of various routes of administration. For example, oral administration would be expected to require higher dosages than administration by intravenous injection. Variations in these dosage levels can be adjusted using standard empirical routines for optimization, as is well understood in the art.

35

Polypeptides used in treatment can also be generated endogenously in the subject, in treatment modalities often referred to as "gene therapy" as described above. Thus, for example, cells from a subject may be engineered with a polynucleotide, such as a DNA or RNA, to encode a polypeptide ex vivo, and for example, by the use of a retroviral plasmid vector. The cells are then introduced into the subject.

**SUBSTITUTE SHEET (RULE 26)** 

All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

																								11.	$\Pi \mathcal{F}$	$\Box \Box$	$\sim 13$	J_		
GCA	GT.	гсст	ĢAG	AGA	AGA	ACC	CTG	AGGA	AC	AGA	CGT1	rcc(	CTC	GCG	GCC	CTG	GCA	ССТ	CCA.	ACC	CA	GAT.	ATG	CTG	CTG	CTG	CTG	CTG	CTG	90
CGT	CA	AGGA	стс	TCT	TCT	TGG	GAC	TCC	гтĠ	TCT	GCA/	4GG	GAG	CGC	CGG	GAC	CGT	GGA	GGT	TGG	GGT	CTA	TAC	GAC	GAC	GAC	GAC	GAC	GAC	30
																							М	L	L	L	L	L	L	
																	<b></b>								T C C					
		GCTC	+			<del></del>			+				+					+	+		<del></del>		+			<del></del>			<del>+</del>	180
GGG	GAI	CGAG	ACC	CCC	TCC	CTC	TCC	CAC	STT	CCT	GTC	TTC	TÇA	TTG	GCC	TTC	CTA	AIG	AGC	GAC	i GC	IAC	النال	IÇA	AGG	CAU	i GG	CAC	GII	
Р	L	L	W	G	R	Ε	R	٧	ε	G	0	K	S	N	R	K	0	Υ	S	Ļ	T	M	۵	S	S	٧	٢	٧	a	
GAG	GGI	CATG	TGT	GTC	CAT	GTG	CGC	TGC	тсс	TTC	TCC	TAC	CCA	GTG	GAC.	AGC	CAG	ACT	GAC	TCT	GAC	CCA	GTT	CAT	GGC	TAC	TGG	ттс	CGG	
CTC	CC	GTAC	ACA	CAG	GTA	CAC	GC G	ACG/	<del>+</del> AGG	AAG	AGG	ATG	<del>:                                    </del>	CAC	CTG	TCG	GTC	TGA	C T G	AGA	CTG	GGT	CAA	GTA	CCG	ATG	ACC.	AAG	GCC	270
	•	м	_	V	ш	v		c	c	_	c	v	0	v		s		Т	D	s	n	o	V	ч	c	¥	W	£	p	
E.	G	•••	·	٧	п	٧	Α.		3		3	'	F	٧			_	•			Ü	•	•	"	G	٠.	"	'	•	
		GAAT	+-					+					+			<del></del>		<del></del>	$\overline{}$				+			<del></del> -			<del>+</del>	360
CGT	CC	CTTA	CTA	TAT	TCG	ACC	TTC	CGA	GGT	CAC	CGG	TGT	TTG	TTG	GGT	CGA	ACC	CGT	CAC	GTC	CTC	CTT	TGA	GCC	CTO	GCT	AAG	GTG	GAG	
A	G	N	D	i	\$	W	κ	Α	Ρ	٧	Α	T	N	N	Ρ	Α	W	Α	٧	a	Ε	٤	T	R	٥	R	F	Н	L	
CTT	GG	GGAC	CCA	CAG	ACC	AAA	AAT	TGC.	ACC	CTG	AGC.	ATC	AGA	GAT	GCC	AGA	ATG	AGT	GAT	GCG	GGG	AGA	TAC	TTC	TTT	rcgr	ATG	GAG	AAA	
	- +-	ССТО				-+-		+	+				+			-+-			+				+			<del>!</del>		<del></del>	+	450
	_			•	-															А		R	Y	_	_	R	м		K	
L	b	0	۲	u	•	ĸ	N	L	1	L	3	1	ĸ	U	А	ĸ	(1	3	Ü	^	G	K	1		•			-		
GGA	AA	TATA	AAA	TGG	AAT	TAT	AAA	TAT	GAC	CAG	CTC	TCT	GTG	AAC	GTG	ACA	TAC	CCT	CCT	CAG	AAC	TTG	ACT	GTG	AC1	rgto	TTC	CAA	GGA	540
CCT	TT	ATAI	ттт	ACC	TTA	ATA	TTT	ATA	CTG	GTC	GAG	AĠA	CAC	TTG	CAC	TGT	ATG	GGA	GGA	GTC	TTG	AAC	TGA	CAC	TGA	ACAG	AAG	GTT	CCT	
G	N	i	Κ	W	N	Y	K	Υ	0	Q	Ĺ	\$	٧	N	٧	T	Y	Ρ	Р	Q	N	L	T	٧	T	٧	F	Q	G	
GAA	GG	CACA	AGCA	TCC	ACA	AGCT	CTG	GGG	AAC	AGC	TCA	тст	стт	TCA	GTC	СТА	GAG	GGC	CAG	TCT	CTG	CGC	TTG	GTC	TGI	FGC1	GTT	GAC	AGC	
		GTGT	+			-+-		+					+			-+-		<del></del>	+				+					+		630
_				•	_			•		_																	v	0	c	
E	G	۲,	А	5	1	А	Ĺ	G	N	5	S	5	Ĺ	S	٧	L	Ε	G	u	S	L	ĸ	L	٧	L	А	٧	U	J	
AAT	CC	CCC	rgcc	AGG	CTO	SAGO	TGG	ACC	TGG	AGG	AGT	CTG	ACC	CTG	TAC	cçc	TCA	CAG	CCC	TCA	AAC	CCT	CTG	GTA	CTO	GGAG	CTG	CAA	GTG	720
TTA	GG	GGG	ACGO	TCC	GAC	TCG	ACC	TGG	ACC	тсс	TCA	GAC	TGC	GAC	ATG	GGG	AG1	rgro	GGG	AGT	TTG	GGA	GAC	CAT	GAG	CCTC	GAC	GTT	CAC	
N	ρ	Ρ	Α	R	L	s	W	Т	W	R	s	L	Ţ	Ĺ	Y	P	s	a	Р	s	N	Ρ	L	٧	L	Ε	L	a	٧	
C 4 C	. с т	GGG	- C A T		cci	2 C A A	TTC	۸۵۲	TC1	rcca	CCT	глс			רדם	cet	TC	- C A G	יר אר	CTT	TCC	·r T G	: A A C	רדר	TCI	осто	CAA	CAG	GAG	
	-+	CCC						+					<del></del>																	810
																													_	
u		C	0	두	G	F	E	Т	_	R	Δ	ก	N	S	1	G	S	Ω	н	V	S	L	N	L	S	L	a	a	ᆫ	

																							H	15	$\mathcal{Q}$	KL	18			
CGG	CAC	GAG	TGG	ACA.	ACC	ATC	AGG	GAG	CCA	GGA	CAC	AGA	GGG	GCA	GAG	CAA	GTC.	AGC#	ATT	GGC	GCC	CCT	ŢCC	TCA	GAT	ccc	TAT	CAT	CTT	
GCC	GTG	стс	ACC	TGT	TGG	TAG	тсс	CTC	GGT	CCT	GTG	TCT	CCC	CGT	CTC	GTT	CAG	TCG	TAA	CCG	CGG	GGA	AGG	AGT	СТА	GGG	ATA	GTA		90
						<del>;</del> -		+		TACA			+			<del></del>			<del></del>				<del></del>	-+					+	180
		0.0	.,,,				,,,,,					•		,,,,,	,,,,,,				. •				М	Α	E	1	н	T	Р	
TAT	тст	тсс	TTG	AAG	AAA	CTG	TTA	тст	TTA	ACTC	AAT	GGC	TTC	GTG	GCT	GŢG	TCT	GGC	ATC.	ATC	CTA	GTT	ggc	CTG	GGC	ATT	GGT	GGT.	AAA	
ATA	AGA	AGG	AAC	TTC	TTT	GAC.	AAT	AGA	AAT	GAG	TTA	CCG	AAG	CAC	CGA	CAC	AGA	CCG	TAG	TAG	GAT	CAA	ccs	GAC	ccg	TAA	CCA	CCA	111	270
Y	s	S	Ĺ	К	K	L	L	S	L	L	N	G	F	٧	А	٧	S	G	1	1	L	٧	S	L	G	į	G	G	Κ.	
TGT	GGA	GGG	gcc	TCT	CTG	ACG	AAT	GTC	СТС	GGG	CTG	тсс	TCC	GCA	TAC	стс	CTT	CAC	GTŢ	GGC	AAC	CTG	ŢGC	CTG	GTG	ΑŢG	GGA	TGC	ATÇ	200
ACA	CCT	ССС	CGG	AGA	GAC	TGC	TTA	CAG	GAG	CCC	GAC	AGG	AGG	CGT	ATG	GAG	GAA	GTG	CAA	ccG	TTG	GAC	ACG	GAC	CAC	TAC	CCT	ACG	TAG	360
С	G	G	A	S	L	T	N	٧	f	G	L	S	S	Α	Υ	L	L	Н	٧	G	N	L	С	L	٧	M	G	С	ī	
ACG	GTA	CTG	CTT	GGC	TGT	GCC	GGG	TGG	TAT	rgg A	GCG	ACT	AAA	GAG	AGC	AGA	GGC	ACG	стс	TTG	TTT	TGC	ATC	CTG	TCA	ATG	GTT	ATT	GTC	450
TGC	CAT	GAC	GAA	CCG	ACA	.cgg	ССС	ACC	ATA	ACCT	CGC	TGA	TTT	стс	TCG	тст	CCG	TGC	GAG	AAC	AAA	ACG	TAG	GAC	AGT	TĄC	CAA	TAA	CAG	, 55
Т	٧	L	L	G	С	Α	G	W	Y	G	А	T	ĸ	E	s	R	G	Т	L	L	F	С	į	L	s	М	٧	ı	٧	
стс	ATC	ATG	GAA	GIT	ACA	GCT	GCC	ACA	GTG	GTC	CTT	CTT	TTC	TTT	CÇA	ATT	GTT	GGA	GAT	GTG	GCC	TTG	GAA	CAC	ACC	TTC	GTG	ACC	CTG	
GAG	TAG	TAC	: C T T	CAA	TGT	CGA	CGG	TGT	CAC	CAG	GAA	GAA	AAG	AAA	GGT	TAA	CAA	ССТ	C T A	CAC	CGG	AAC	CTT	GTG	TGG	AAG	CAC	TGG	GAC	540
L	i	м	ε	٧	Т	Α.	А	T	٧	٧	Ĺ	L	F	F	Р	I	٧	G	D	٧	А	L	Ξ	Н	T	F	٧	T	L	
AGG	AAG	AAT	ŢAC	AGA	GGT	TAC	AAC	GAG	CCA	AGAC	GAC	TAT	тст	ACA	CAG	TGG	AAC	TTG	GTC	ATG	GAG	AAG	CTA	AAG	TGC	TGT	GGG	GTG	AAT	630
TCC	TTC	TTA	ATG	TCT	CCA	ATG	TTG	стс	GGT	CTG	CTG	ΑΤΑ	AGA	TGT	GTC	ACC	TTG	AAC	CAG	TAC	СТС	TTC	GAT	TTC	ACG	ΑĊΑ	ccc	CAC	TTA	-
R	K	N	Υ	R	G	Y	N	ε	P	0	0	Y	S	Т	Q	W	N	L	٨	М	ε	К	L	κ	С	С	G	٧	N	
AAC	TAC	ACA	GAT	TTT	TCT	GGC	TCT	TCC	TTC	GAA	AATG	ACA	ĄCG	GGC	CAC	AÇC	TAC	CCC.	AGG	AGT	TGC	TGT	AAA	TCC	ATC	GGA	AGT	GTG	TCC	720
TTG	ATG	TGT	CTA	AAA	AGA	ccg	AGA	AGG	AAC	GCTT	TAC	TGT	TGC	CCG	GTG	TGG	ATG	GGG	TCC	TCA	ACG	ACA	.TTT	AGG	TAG	CCT	TCA	CAC	AGG	720
N	Y	Т	D	F	S	G	S	S	F	Ε	M	Т	T	G	н	T	Y	ρ	R	S	С	С	K	S	i	G	S	٧	s	
TGT	GAC	GGA	ÇGC	GAT	GTG	TÇT	CCA	AAC	GTO	CATO	CAC	CAG	AAG	GGC	TGT	TTC	CAT	AAA	стс	СТА	AAA	ATC	ACC	AAG	ACT	CAG	AGC	TTC		810
ACA	CTG	CCT	GCG	CTA	CAC	AGA	GGT	TTG	CAC	STAG	GTG	GTC	TTC	CCG	ACA	AAG	GTA	TTT	GAG	GAT	TTT	TAG	TGC	770	TGA	GTC	TC	AAG	TGG	010
С	0	G	R	0	٧	s	Ρ	N	٧	ī	Н	۵	к	G	С	F	н	К	L	L	κ	i	7	ĸ	Т	۵	S	F	Ť	ieg I!
CTG	AGT	GGG	AGC	TCT	CTG	GĢA	GCT	GCA	GTO	GATA	ACAG	AGG	TGC	GGG	тст	cgc	TAT	GTT	GCG	CAG	GCT	GGT	CT1	GAA	CTG	cŢ	GCC	TAA	AĞC	900
GAC	TCA	CCC	TCG	AGA	GAC	CCT	CGA	CGT	CAC	CTAI	rgro	тсс	ACC	ccc	AGA	GCG	ATA	CAA	CGC	GTC	CGA	CCA	GAA	ACTT	GAC	GAC	CGC	ATT		200

HPWA E25													
GCGTCCCTCTGCCTGCCCACTCAGTGGCAACACCCGGGAGCTGTTTTGTCCTTTGTGGAGCCTCAGCAGTTCCCTCTTTCAGAACTCAC	0												
CCGCAGGGAGACGGACGGTGAGTCACCGTTGTGGGCCCTCGACAAAACAGGAAACACCTCGGAGTCGTCAAGGGAGAAAGTCTTGAGTG  TGCCAAGAGCCCTGAACAGGAGCCACCATGCAGTGCTTCAGCTTCATTAAGACCATGATGCTCCTTCAATTTGCTCATCTTTCTGTGT													
	80												
CGGTTCTCGGGACTTGTCCTCGGTGGTACGTCACGAAGTCGAAGTAATTCTGGTACTACTAGGAGAAGTTAAACGAGTAGAAAGACACA	•												
MQCFSFIKTMMILFNLLIFLC													
GTGCAGCCCTGTTGGCAGTGGGCATCTGGGTGTCAATCGATGGGGCATCCTTTCTGAAGATCTTCGGGCCACTGTCGTCCAGTGCCATG	70												
CACGTCGGGACAACCGTCACCCGTAGACCCACAGTTAGCTACCCCGTAGGAAAGACTTCTAGAAGCCCGGTGACAGCAGGTCACGGTAC	70												
G A A L L A V G I W V S I D G A S F L K I F G P L S S S A M													
AGTȚIGICĂACGIGGCIĂCTICCICAICGCAGCCGGCGITGIGGICIȚIGCICITGGȚIICCIGGGCȚGCIAIGGIGCIAAGACIGAG													
TCAAACAGTTGCACCCGATGAAGGAGTAGCGTCGGCCGCAACACCAGAAACGAGAACCAAAGGACCCGACGA	60												
Q F V N V G Y F L I A A G V V, V F A L G F L G C Y G A K T E													
GCAAGTGTGCCCTCGTGACGTTCTTCTTCATCCTCCTCCTCATCTTCATTGCTGAGGTTGCAGCTGCTGTGGTCGCCTTGGTGTACACC	50												
CGTTCACACGGGAGCACTGCAAGAAGAAGTAGGAGGAGGAGTAGAAGTAACGACTCCAACGTCGACGACACAGCGGAACCACATGTGG	50												
S K C A L V T F F I L L L I F I A E V A A A V V A L V Y T													
CAATGGCTGAGCACTTCCTGACGTTGCTGGTAGTGCCTGCC	40												
GTTACCGACTCGTGAAGGACTGCAACGACCATCACGGACGG	40												
T.M. A.E.H.F.L.T.L.L.V.P.A.I.K.K.D.Y.G.S.Q.E.D.F.T.Q.V.W.N.													
CCACCATGAAAGGGCTCAAGTGCTGTGGCTTCACCAACTATACGGATTTTGAGGACTCACCCTACTTCAAAGAGAACAGTGCCTTTCCC													
GGTGGTACTTTCCCGAGTTCACGACACCGAAGTGGTTGATATGCCTAAAACTCCTGAGTGGGATGAAGTTTCTCTTGTCACGGAAAGGG	30												
T T M K G L K C C G F T N Y T D F E D S P Y F K E N S A F P													
CATTCTGTTGCAATGACAACGTCACCAACACAGCCAATGAAACCTGCACCAAGCAAAAAGGCTCACGACCAAAAAGTAGAGGGTTGCTTC													
GTAAGACAACGTTACTGTTGCAGTGGTTGTGTGTGTGTTTTGGACGTGGTTTCCTTTTCCGAGTGCTGGTTTTTCATCTCCCAACGAAG	20												
P F C C N D N V T N T A N E T C T K Q K A H D Q K V E G C F													
ATCAGCTTTTGTATGACATCCGAACTAATGCAGTCACCGTGGGTGG													
TAGTCGAAAACATACTGTAGGCTTGATTACGTCAGTGGCACCCACC	10												
N Q L L Y D I R T N'A V T V G G V A A G I G G L E L A A M N													
GTGTCCATGTATCTGTACTGCAATCTACAATAAGTCCACTTCTGCCTCTGCCACTACTGCTGCCACATGGGAACTGTGAAGAGGCACCC													
CACAGGTACATAGACATGACGTTAGATGTTATTCAGGTGAAGACGGAGACGGTGATGACGACGGTGTACCCTTGACACTTCTCCGTGGG	00												

Päge 1

																									Į-	TT	7	E	FS	عا		
 AAA	AAA	AAC	AAG	GTO	CC	CAC	AG	CAA	AGA	AA,	AGGA	ATA	GG/	TC	AAG	AGAI	TAC	GT(	GGC	rgc	TGC	GCAC	AG	CAA	GCA	TGA	ΑŢ	TÇG,	ATGA	ACT	TCA	90
TTT	TT	TTG	<del>:</del> FTC(	CAC	GG	G T G	TC	GTT	TCT	TT	TCCI	TAT	CC	ΑG	TTC	TCTA	A T G	CA	CCG	ACG.	AC(	CGT	CTC	GTT	CGT	ACT	TA	AGC	TAC	TGA/	AGŤ	50
																										М	N	S	М	T	S	•
											GTG										_											180
GTC	AA	GGC	CAC	CGC	STT	AAC	SAC	ACA	ACC	AC	CAC	CGTC	GGG	G T C	TTA	CCA	ΑΤΑ	GG.	ACA:	CTG	GG	GTC	CTT	AAI	'ACΔ	GAC	ST G	CAC	GGG	GAC.	ATA	
А	٧	Р	٧	A	N		3	٧	L	٧	٧	Α	Ρ	н	N	G	Υ	Р	٧	7	:	Ρ	G	Ĺ	М	S	Н	٧	P	L	Y	
CCAA	AC	AGC	CAG	CC	GC A	AG1	rcc	ACC	TAC	TT	CCT	GGGA	4AC	CC/	ACCT	AGT	TTG	GT	GTC	GAA	TG	TGA	ATG	GG	AGO	СТ	STG	CAG	AAA	GCT	CTG	270
							•				GGA						_						,								,	270
ρ	N		a	ρ				н		٧				Ρ			L	٧					N	G	0	Ρ	٧	0	κ	Α	L	
		000		٠.	C T T		~~		TCI	- A G	ATC	۸۲۲	ΛТΤ	CCI	-r T r	CCT.	<b>Γ</b> Δ(	^ A T	cee	CCT	CG	GCT	CCA	TC.	ATG(	GC G	A C G	GTT	стс	GTA	GGG	
											TAG										_							_				360
3 1 1 1	- 1 1	LLG	111	1 (3)	GAF																						<b>.</b>	.,		.,		
K	_	_	Κ								I																1	٧	L	٧	G	
GAA	TAC	стс	TCT	ΑΤ	TTC	ΑT	ŢC:	TAC	GGA	GGÇ	TTT	CCC	TTC	ŢĢ	GGGA	AGGC	TT	GTG	GTT	TAT	ÇΑ	111	CAC	GΑ	TCT	стс	TCC	GTG	GCA	GCA	GAA	450
CTT	ATG	GAC	AGA	TA	AAC	A T E	AG	ATG	CT	CCC	AAA	GGG	AAG	АC	ccc	rccg	AA	CAC	CAA	ΑТА	ĠT	AAA	GTO	CT	AGA	GAG	AGC	3C A C	CGT	CGT	стт	
ε	Y	L	S	i	Ş	5	F	Υ	G	G	F	Ρ	F	W	G	G	L	٨	, F			Ī	s	G	S	L	S	٧	A	A	ε	
AAT	CAG	SC C A	TAT	TC	TTA	ATT	GC	CTG	CTG	TCI	rggc	AGT	TTG	GG	CTT	GAAC	ΑT	CGT	CAC	TGC	ΑA	TCT	GC.	гст	ĢCA	GTT	GG	AGŢC	:ATA	CTC	тто	540
											CCG			-							$\overline{}$				+							340
N	a	Ρ	Y	S		Y	c	L	L	s	G	s	L	G	L	N	I	١	/ 9	6 A	4	ī	С	S	Α	٨	G	٧	I	L	F	
ATC	AC.	AGA T	rc T A	AΑG	TA	TTC	ÇC	CAC	CCA	TA	TGCC	TAC	ccc	ĢΑ	CTA	TTAT	rcc	TTA	AC G (	сто	gg	GGT	TG	AAC	CCT	GGA	AT.	GGC	SATT	TCT	rggc	: - 630
TAG	TGI	TCT/	AGAT	гтс	ΑТ	AAG	GG	GTG	GGT	ATA	ACGG	ATG	GGC	C T	GAT	AATA	4GG	AA	rgc	GAG	ccc	CAC	AC	TTG	ĠĠĀ	CCT	TΑ	ccd	CTAA	\AGA	ACCC	
1	т	D	L	Ş	5	l	Ρ	Н	٩	Υ	А	Y	Þ	0	ΙΥ	Y	Р		Υ ,	4 1	N.	G.	٧	N	Р	G	М	Α	I	S	G	
GTG	CTO	GCT(	GGT(	CTT	·c r	GCC	TC	CTG	GAG	TT:	TGGC	ATC	:GCA	A T G	cgc	ATC	гтс	CC.	ACT	TTG	GC.	TGC	CAG	TTG	igTC	TGC	TG	TCĄ	ATC/	AAG	CAAT	720
											ACCO																					, ,,,,
			v	-	_	_		,	_	_	G	1	۸	٢					Н		G	۲	a	1	y	· c	c	Ω	s	s	N	
٧	L	L	٧	•	•	Ļ	_	L	C	г	_		^	`		-					-	·		_	•	•	_					_
											AGC										_											
CAC	TC.	ACA	GTA	GA.	۲AG	GT!	ГŢG	TAG	ΑΤΑ	CG	TCG	7770	GGG	TCA	AC T A	GTG	GGG	TC	TTG	GCC	AC	TGG.	AGT	GG1	TGGT	TC/	AΑΤ	AAG	GTC.	ACT	CTA	3
v	c	V	,	,	,	P	N	ı	Υ	Δ	Δ	N	ρ	١	, ,	т	F	•	Ε	P	٧	T	s	Р	2	s	Y	S	S	Ε	i	

Päge 1

																								$\Box$		ユユ	トし	<u>ــم د</u>		
CACG	AG	CAG	GGT	CTC	SGG	CTAC	GTC/	ATG	GC Ģ	TCC	CCG.	гст	GGA	AGA	CTG	AG.	ACT/	AAA	CA	STEA	ATTA	CTI	GTI	TTCA	AGA	AGC	TTC	TGC	TA	90
GTGC	TC	GTC	CCA	GAG	CCC	GAT	CAG	TAC	:GC/	AGG	GGC/	AGA(	3001	rc T	GAC	stc:	TGA	TTT	GG T	CAG	ΓΑΑΊ	GA/	ACAA	AAG	TCI	rcgo	CAAC	SACG		
								м	Α	S	Ρ	S	R	R	L	۵	T	K	Р	٧	í	T	С	F	К	S	٧	L	L .	
ATCT	AC.	ACT	TTT	ATT	TTC	TGG	ATC.	ACT(	GGC	GTT.	ATC	CTT	CTT	GCA	GTT	GGC.	ATT	TGG	GGC.	AAG	GTG	AGC	TG	GAG	AAT	TAC	TTT	гсто	TT:	180
TAGA	TG	TGA.	AAA	TAA.	AAG	ACC	TAG	TGA	ccĠ	CAA	TAG	GAA	GAA(	CGT	CAA	CCG	TAA.	ACC	ccg	TTC	CAC	TÇG	GAC	CTC	TTA	A T G	ΔΑΑΑ	4GAC	AA	
1	Υ	T	F	Į	F	W	1	Т	G	٧	i	L	L	А	٧	G	1	W	G	K	٧	s	L	Ε	N	Y	F	s	L	
TTAA	AT	GAG	ĄAG	GCC	ACC	AAT	GTC	ccc	TTÇ	GTG	стс	ATT	GCT.	ACT	GGT.	ACC	GTC	ATT	ATT	CTT	TTG	GGC.	ACC	TTT	GGT	TGT	TTT	GC F/	ACC	270
AAT1	TA	стс	TTC	cgg	TGG	TTA	CAG	GGG	AAG	CAC	GAG	TAA	CGA	TGA	CCA	TGG	CAG	TAA	TAA	GAA	AAC	CCG	TGG.	AAA	CCA.	ACA	AAA	CGA	rgg	2,0
L	N	Ε	Κ	А	T	N	٧	Ρ	F	٧	L	I	A	T	G	T	٧	Į	I	Ļ	L	G	Т	F.	G	С	F	Α.	т	
TGC	GA	GCT	TCT	GCA	TGG	ATG	CTA	AAA	CTG	TAT	GCA	ATG	TTT	ÇTG	ĄC T	стс	GTT	TTT	TTĢ	GTC	GAA	CTG	GTC	GCT	GCC.	AŢC	GTA	GGA	TTT	260
ACG	GCT	CGA	AGA	CGT	ACC	TAC	GAT	TTT	GAC	ATA	CGT	TAC	AAA	GAC	TGA	<del>. ;</del> GAG	CAA	AAA	AAC	CAG	CTT	GAC	CAG	CGA	CGG	TAG	CAT	CCT.	AAA	360
С	R	А	S	А	w	М	L	K	L	Υ	А	М	F	L	т	L	٧	F	Ļ	٧	ε	L	٧	Α	А	ı	٧	G	F	
GTT	TTC	AGA	CAT	GAG	ATT	AAG	AAC	AGC	TTT	AAG	AAT	AAT	TAT	GAG	AAG	GCT	TTG	AAG	CAG	TAT	AAC	тст	ACA	GGA	GAT	TAT	AGA	AGC	CAŢ	
CAA			٠										+	+		<del></del>		<del></del>					<del>,</del>			<u> </u>				450
	_	_		_				_	_	.,			v	_			,	.,	0	v	A.I	c	т	c	n	٧	B	S	н	
٧	F	R	н	Ε	i	K	N .	• 5	ŀ	ĸ	N	N	T	٤	K	А	Ļ	^	u	•	14	J	•	G	Ü	,		J	••	
GCA						<u> </u>							-										+			<del></del>			<del></del>	540
CGT	CAT	CTG	TTO	TAG	GTT	TTA	TGC	AAC	GTA	ACA	AÇA	CCA	CAG	TGG	CTA	ATA	TCT	TCTA	ACC	TGT	CTA	TGA	TTA	ATA	ATG	AGT	CTT	TTT	CCT	
A	٧	O	κ	. 1	Q	N	Т	L	н	С	С	G	٧	Τ	D	Y	R	0	W	τ	D	Τ	N	Y	Y	S	Ε	κ	G	
TTT	сст	TAAG	AG	TGC	TGT		CTT	GAA	GAT	TG	AC1	CCA	AÇAG	AGA	GAT	GÇA	GAC	CAAA	AGT A	AAC	AAT	GAA	GGT	TGT	ттт	ATA	AAG	GTG	ATG	630
AAA	GGA	TTC	TC	ACC	ACA	TTT	GAA	CTT	CTA	ACA	TGA	\GG1	GTO	TCI	CTA	CGI	СТО	STTI	CAI	TTO	TTA	CTT	CCA	ACA	AAA	TAT	TTC	CAC	TAC	
F	Ρ	K	S	С	С	Κ	L	Ε	0	С	T	ρ	a	R	۵	А	0	K	٧	N	N	ε	G	С	F	I	K	٧	M	
													***		-00	. C T 1		T T C (		- C A /		` A T T		ΔΤΩ		rTr	1600	TAC	TGC	
	_												<del></del>														9000			720
TGG	TAA	ATAT	rc T	CAG	ICI !	IIAL	.CCI	LAC	iCAA	ALG	ILL	IIAA	AAGG	iAAA	100	ILAA	4667	чис	JAAL	العاد							CGG			
T	1	ľ	Ε	S	Ε	М	G	٧	٧	Α	G	i	S	F	G	٧	Α	С	F	0	L	i	G	1	F	Ĺ	А	Y	С	
																														810
GAG	AG	AGC	ACG	GTA	TTG	T T T A	ATTO	GGT	CATA	ACT	CTA	TCA	CATI	rggi	GTT	4CA	TAG	AGA	ccc	GGA	FAAC	GAG	GAGA	ATG	SAAA	ATT(	:CT0	;TAA	ATC	
L	s	R	A	ī	т	N	N	Q	Y	Ε	i	٧	÷	LS	XC.	J	D	N	0:	2	2)									
GGT	CC	CCC	CTG	TGA.	ATT	AGA	4AG	TTG	CTT	GGC	TGG.	AGA.	AÇT	GAC.	AAC	ACT.	ACT	TAC	TGA	TAG	ACC	AAA	4 A A (	CTA	CAC	CAG	TAGO	3770	ATT	900
CCA	GG	GGG	GAC	ACT	TAA	TCT	TTC	AAC	GAAI	CCG	ACC	TET	TGA	CTG	TTG	TGA	TGA	ATG.	ACT.	ATC	TGG'	TTT.	TTT	GAT	GTG	GTC.	ATC	CAAC	TAA	

41 HSBR FOZ	Page 2
AATCAAGATGTATGTAGACCTAAAACTACACCAATAGGCTGATTCAATCAA	
TTAGTTCTACATACATCTGGATTTTGATGTGGTTATCCGACTAAGTTAGTT	+ 990 T
GTTTGCTATGTTCTAAGTCCACCTTCTATCCCATTCATGTTAGATCGTTGAAACCCTGTATCCCTCTGAAACACTGGAAGAGCTAGTA	A + 1080
CAAACGATACAAGATTCAGGTGGAAGATAGGGTAAGTACAATCTAGCAACTTTGGGACATAGGGAGACTTTGTGACCTTCTCGATCAT	
TTGTAAATGAAGTAAAAAAAAAAAAAAAAAAAAAAAAA	
AACATTTACITCATTTITTTTTTTTTTTTTTTTTT	

BNSDOCID: <WO\_\_\_9831799A3\_IA>

HLIAHXO													
ACGAGCATIGCCGCTCTCTCGGTGAGCGCAGCCCCGCTCTCCGGGCCGGGCCTTCGCGGGCCACCGGCCCATGGGCCAGTGCGGCATC													
GTGCTCGTAACGGCGAGAGAGCCACTCGCGTCGGGGGGGG													
ACCTCCTCCAAGACCGTGCTGGTCTTTCTCAACCTCATCTTETGGGGGGGCAGCTGGCATTTTATGCTATGTGGGAGCCTATGTCTTCATC	180												
TGGAGGAGGTTCTGGCACGACCAGAAAGAGTTGGAGTAGAAGACCCCCCGTCGACCGTAAAATACGATACACCCTCGGATACAGAAGTAG													
T S S K T V L V F L N L I F W G A A G I L C Y V G A Y V F I													
	270												
TGAATACTACTGATACTGGTGAAGAAACTTCTACACATGTGCGAGTAGGGACGACATCACTAGTATCGACATCCTCGGGACGAAAAGTAG													
TYOOYOHFFEOVYTLIPAVVIIAVGALLFI													
ATTGGGCTAATTGGCTGCTGCCACAATCCGGGAAAGTCGCTGTGGACTTGCCACGTTTGTCATCATCCTGCTCTTGGTTTTTGTCACA													
TAACCCGATTAACCGACGACACGGTGTTAGGCCCTTTCAGCGACACCTGAACGGTGCAAACAGTAGTAGGACGAGAACCAAAAACAGTGT	360												
I G L I G C C A T I R E S R C G L A T F V I I L L V F V T													
GAAGTTGTTGTAGTGGTTTTGGGATATGTTTACAGAGCAAAGGTGGAAAATGAGGTTGATCGCAGCATTCAGAAAGTGTATAAGACCTAC	450												
CTTCAACAACATCACCAAAACCCTATACAAATGTCTCGTTTCCACCTTTTACTCCAACTAGCGTCGTAAGTCTTTCACATATTCTGGATG	450												
E V V V V I G Y V Y R A K V E N E V D R S ! Q K V Y K T Y													
AATGGAACCAACCCTGATGCTGGCCGGGCTATTGATTATGTACAGAGACAGCTGCATTGTTGTGGAATTCACAACTACTCAGACTGG	540												
TTACCTIGGTTGGGACTACGACGATCGGCCCGATAACTAATACATGTCTCTGTCGACGTAACAACACCTTAAGTGTTGATGAGTCTGACC													
N G T N P D A A S R A I D Y V Q R Q L H C C G I H N Y S D W													
GAAAATACAGATTGGTTCAAAGAAACCAAAAACCAGAGTGTCCCTCTTAGCTGCTGCAGAGAGACTGCCAGCAATTGTAATGGCAGCCTG													
	630												
CTTTTATGTCTAACCAAGTTTCTTTGGTTTTTGGTCTCACAGGGAGAATCGACGACGTCTCTCTGACGGTCGTTAACATTACCGTCGGAC													
ENT D'W FKETKNOS V PLSCCRETASNCNGSL													
GCCCACCCTTCCGACCTCTATGCTGAGGGGTGTGAGGCTCTAGTTGTGAAGAAGCTAGAAGAAATCATGATGCATGTGATCTGGGCCGCA	700												
CGGGTGGGAAGGCTGGAGATACGACTCCCACACTCCGAGATCAACACTTCTTCGATGTTCTTTAGTACTACGTACACTAGACCCGGCGT	720												
A H P S D L Y A E G C E A L V V K K L Q E I M M H V I W A A													
CTGGCATTTGCAGCTATTCAGCTGCTGGGCATGCTGTGTGCTTGCATCGTGTTGTGCAGAAGGAGTAGAGATCCTGCTTACGAGCTCTTC	810												
GACCGTAAACGTCGATAAGTCGACGACCCGTACGACACACGAACGTAGCACAACACGTCTTCCTCATCTCTAGGACGAATGCTCGAGAAG													

1662 (SCOID NO 7)

BNSDOCID: <WO\_\_\_9831799A3\_IA>

HTPBA27	
CACGAGCGCAGAGCTTGGGGCTTCCTTGGTCGCACCCACC	GTC -+ 90
TIGCTCGCGTCTCGAACCCCGAAGGAACCAGCGTGGGTGGTGGACGGAC	
TCTTTCCTGTGGCCAGCCCAGAACTGAAGCGCTGCGGCATGGCGCGCGC	ACC 
AGAAAGGACACEGGTCGGGTCTTGACTTCGCGACGCCGTACCGCGCGCGGGGGGGG	TGG
M A R A C L Q A V K Y L M F A F	N
TGTTCTTCTGGCTGGGAGGCTGTGGCGTGCTGGGTGTCGGCATCTGGCTGG	ССТ
ACAAGAAGACCGACCCTCCGACACCGCACGACCCACAGCCGTAGACCGACC	2/0
	S
TCCCGTCCCTGTCGGCTGCCAACCTGCTCATCACCGGCGCCTTTGTCATGGCCATCGGCTTCGTGGGCTGCCTGGGTGCCATCA	- , 300
AGGGCAGGGACAGCCGACGGTTGGACGAGTAGTAGTGGCCGCGGAAACAGTACCGGTAGCCGAAGCACCCGACGGACCCACGGTAGT	TCC
FPSLSAANLLIITGAFVMAIGFVGCLGA!	K
AGAACAAGTGCCTCCTGCTCACTTTCTTCCTGCTGCTGCTGCTGGTGTTCCTGCTGGAGGCCACCATCGCCATCCTCTTCTCGCCT	ACA 450
TCTTGTTCACGGAGGACGAGTGAAAGAAGGACGACGACGACCACAAGGACGACCTCCGGTGGTAGCGGTAGGAGAAGAAGCGGA	
FNKCLLLTFFLLLLUFLLEAT!A!LFFA	Y
CGGACAAGATTGACAGGTATGCCCAGCAAGACCTGAAGAAAGGCTTGCACCTGTACGGCACGCAGGGCAACGTGGGCCTCACCAACC	CCT
GCCTGTTCTAACTGTCCATACGGGTCGTTCTGGACTTCTTTCCGAACGTGGACATGCCGTGCGTTCCACCCGGAGTGGTTGC	<del></del> 340
T D K ! D R Y A Q Q D L K K G L H L Y G T Q G NV G L i N	А
GGAGCATCATCCAGACCGACTTCCGCTGCTGTGGCGTCTCCAACTACACTGACTG	. 000
CCTCGTAGTAGGTCTGGCTGAAGGCGACGACACCGCAGAGGTTGATGTGACTGAC	TGA
W S I I Q T D F R C C G V S N Y T D W F E V Y N A T R V P	D
CCTGCTGCTTGGAGTTCAGTGAGAGCTGTGGGCTGCACGCCCCGGGCACCTGGTGGAAGGCGCCGTGCTACGAGACGGTGAAGGTG	rggc
GGACGACCTCAAGTCACTCTCGACACCCGACGTGCGGGGGCCGTGGACCACCTTCCGCGGGCACGATGCTCTGCCACTTCCAC	/20
	w
S C C L E F S E S C G L H A P G T W W K A P C Y E I V K V	
TTCAGGAGAACCTGCTGGCTGTGGGCATCTTTGGGCTGTGCACGGCGCTGGTGCAGATCCTGGGCCTGACCTTCGCCATGACCATG	
AAGTCCTCTTGGACGACCGGACACCCGTAGAAACCCGACACGTGCCGCGACCACGTCTAGGACCCGGACTGGAAGCGGTACTGGTAC	AIGA
LQENLLAVGIFGLCTALVQILGLTFAMTM	Y
GCCAAGTGGTCAAGGCAGACACCTACTGCGCGTAGGCCGCCCACCGCCGGCTTCTCTGCCAAAAGGACGCCCACGGGGAGATGGCC	GCAC + 900
CGGTTCACCAGTTCCGTCTGTGGATGACGCGCATCCGGCGGGTGGCGGCCGAAGAGACGGTTTTCCTGCGGGTGCCCCTCTACCGG	
COVVKADIYCA. (SEQID'NO:24)	

	45		HTPRAZT	Page 2
CCACAGCTGCTTTTCCCACCACCAGCTTCGGTGTTCTGC	CCCATGCTGGGAGGAGGAGG	GAGGGACAGGTGC	CTGGAGCCCCGGAA	CC 990
GGTGTCGACGAAAAGGGTGGTGGTCGAAGCCACAAGACG	GGGTACGACCETCCTCCCTCC	CTCCCTGTCCACG	GACCTCGGGGGCCTT	
CTGTTTCTGGAAGGCCCTAGCTCAGGTGGCTTCAGGGCC	TCCGGACCCCCCTGGGAGGG	GTGGCCACGTGCT	GGCTGCGGAACCCAG	GG 1080
GACAAAGACCTTCCGGGATCGAGTCCACCGAAGTCCCGG	AGGCCTGGGGGGGACCCTCCC	CACCGGTGCACGA	CCGACGCCTTGGGTC	
CAGGGGTGGGAGGGGCCTCCAGCACTTTTTATATTTACG	STATTCTCCAAAGCAGTGTTCA	CACGGGAGCCAGC	CTGTGGCCCCCAGCT	TTC 1170
GTCCCCACCCTCCCCGGAGGTCGTGAAAAATATAAATGC	ATAAGAGGTTTCGTCACAAGT	GTGCCCTCGGTCG	GACACCGGGGGTCGA	
CTGGAAAACAGGTTGGCGCTGGAGGAGCCGGGTCTTGGC	ATCCTGGAGGTGGCCCCACTG	gтсставтастсс.	AGGCGGGGCCGTGGA	ACC 1260
GACCTTTTGTCCAACCGCGACCTCCTCGGCCCAGAACCG	TAGGACCTCCACCGGGGTGAC	CAGGACCACGAGG	TCCGCCCCGGCACCT	
			(SEQ I	DIVO:S
CCTCACCTACATTCCATAGTGGGCCCGTGGGGCTCCTGG	TGCATCTTAATAAAGTGTGAG	CAGCAAAAAAAAA		-
GGAGTGGATGTAAGGTATCACCCGGGCACCCCGAGGACC	ACGTAGAATTATTTCACACTC	GTCGTTTTTTTT		345

																						:	<u> </u>		-v		1			
GCGCCGCCGGCCGCAGCATGGGGCGCTTCCGCGGGGGCCTGCGGTGCATCAAGTACCTGCTTGGCTTCAACCTGCTCTTCTGGCTGCGCGCGGCGCCGGGCGCCGGGCGCCGAGTTGGACGAGAAGCCGACCGA														90																
CGC	GGC	GGC	CCG	GCC	STCC	TAC	ccc	GCG	AAC	GCG	ccc	CCG	GAC	GCC	ACG	TAG	TTC	ATO	GAC	GAC	GAA	CCG	AAG	TTG	GAC	GAG	AAG	ACC	GAC	90
							_	0	_		_	_	,	_	_			v		,		C		A.I	1		_	1.1		
						rı	G	π	-	ĸ	Ŀ	G	L	К	L	1	K	1	L	L	L.	U	F	14	L	L	г	₩	L	
GCT	GGA	ATCG	ĢCC	GT(	CATT	GÇT	ттт	GGA	CTA	TGG	TTT	CGG	ŢTC	GGA	GGT	GĊC	ATA	AAAG	GAG	TTA	TCA	TCA	ĢAG	GAC	٩AG	TÇC	CCA	GAG	TAT	100
CGA	CCI	TAGC	CGG	CAC	STAA	CGA	AAA	CCT	GAT	ACC	AAA	GCC	AAG	CCT	CCA	CGG	TAI	TTTC	СТС	ΑΑΤ	AGT	AGT	ĊTC	CTG	TTC	AGG	GGT	стс.	ΔTA	180
	_	•		.,	,		_	_			~	_	_	•			,	ν	_		_	_	_	_	1/	c	0	_	v	
A	G	3	А	٧	1	А	r	G	L	w	,	ĸ	r	G	G	А	i	K	Ę.	L	3	3	_	U	K	3	r	E.	ı	
TTC	TAT	rgra	ĢGG	CTO	STAT	GTT	CTG	GTI	GGA	AGCC	GGC	GCC	ÇTG	SATO	ATG	GÇC	GTO	GGG	TTC	TTC	GGA	TGC	TGC	GGA	GCC	ATG	CGG	GAG		270
AAG	ATA	ACAC	CCC	GAC	CATA	CAA	GAC	CAA	CCT	CGG	ccc	CGG	GAC	TAC	TAC	CGG	CAC	0000	AAG	AAG	CCT	ACG	ACG	CCT	CGG	TAC	GCC	CTC		270
_	v		_		~	·		v	_	٨	c	۸		м	м	٨	V	c	_	_	C	_	r	G	٨	м	D	=	s	
r	,	٧	u	_	,	٧		٧	G	^	u	~	_	11	" "	^	,	u	•		G		_	u		• • •	1	_	J	
CAA	TG	rgre	CTT	GGA	ATCA	TTT	TTT	ACC	TGC	СТС	CTO	GTG	ATA	TTT	GCT	GCT	GAA	AGTA	ACC	ACT	GGA	GTA	TTT	GCT	TTT	ATA	GGC	AAG	GGG	360
GTT	ACA	ACAC	GAA	CC.	TAGI	TAAA	AAA	TGC	ACC	GAG	GAC	CAC	TAT:	ΓΑΑΑ	CGA	CGA	CT.	FCAT	TGG	TGA	CCT	CAT	AAA	CGA	ΑΑΑ	TAT	CCG	TTC	ccċ	
Ω	С	v		G	S	F	F	т	۲	1	1	v	ı	F	Δ	Δ	۶	v	т	т	G	٧	F	А	F	1	G	к	G	
•	ŭ	·	_	•	Ū	•	•	•	Ū	-	_	•	•	•			_	•	•	·	_	·	·			•	_		_	
GTA	GC.	TATO	CGA	CA.	TGTI	CAG	ACC	ATO	TAT	rgaa	GAC	GCT	TAC	CAAT	GAT	TAC	CT.	TAAA	AGAC	AGG	GGA	AAA	GGC	AAT	GGG	ACA	CTC	ATC	ACC	450
CAT	rcg/	ATAG	GCT	GT	ACAA	AGTC	TGG	TAC	ATA	ACTT	CTO	CGA	ATO	STTA	CTA	AT0	GA	ATT	rcra	TCC	CCI	TTT	CCG	TTA	ccc	TGT	GAG	TAG	TGG	
٧	Α	I	R	Н	٧	a	T	М	Y	Ε	Ε	Α	Y	N	Ð	Υ	L	K	0	R	G	κ	G	N	G	T	L	ı	Т	
																							- <del></del>	- <b>-</b> .					*^^	
			+			TGC							-			<del></del>							+						<del>+</del>	540
AA(	GGT	GAGI	TGI	Ĩ A A A	AGTO	CACG	ACA	.C.C.T	TTT	TCTT	TCC	SAGO	CTI	rgto	CAG	GTI	r G G ,	ATGI	FACG	IGG 1	110	CTC	GAA	GAT	CCT	GIG	IIIC	ITA	ACG	
F	н	S	Т	F	a	С	С	G	K	ε	S	S	Ε	Q	٧	Q	Ρ	7	С	ρ	K	ε	L	L	G	н	K	N	С	
٨Τ٢	-C 4 :	TCA/	ΔТ1	EG A I	c v c c	CATA	۸۲۲	AGI	remi	ΓΛΛΩ	ירדו	- r v c	:c Tc	~ A T 7	ree v	ΛΤΊ	rati	ngg:	FATT	.cc4	ΔΤΊ	.ccv	CCT	стс	<b>Δ</b> Γ (	ΔΤΩ	ידד	ממר	ΔTG	
						TAT		+					-			<del> !-</del>						<del></del>	<del></del>	+		<del></del>			+	630
IAC	الد	4611	IAA	4611	. 160	a i A t	IAG	164	ACA,	4110	GAL	الداد	. <b>G</b> A C	a I A A	4C L 1	IAA	A C A I	امادد	A I A A		IAA	1661	CCA	GAL	166	. I A C			, AC	
1	D	٠Ε	ŀ	Ε	T	I	ĺ	S	٧	K	L	a	L	I	G	I	٧	G	I	G	1	Α	G	L	T	i	F	G	M	
ATA	ATT	CAGO	ATO	ag Ti	CCTO	TGC	TGT	GC	SATA	ACGA	AAC	TCA	AC G A	4GA1	rgrg	ATA	ATG.	AAG	CTAC	TTO	TAC	ATG	AAA	ATT	GCA	ATO	TAA	AGC	TTT	
			+			GACG		-					<del></del>			<del></del>							+			-+-		<del></del>	<del>+</del>	720
17				. <b>.</b>	JUAN	1,,,,,					110															`				
I	F	S	М	٧	L	С	С	Α	I	Я	N	S	R	D	٧	1	•	(	CC	L G		71	V)	$\mathcal{L}$	5					
CA	TAC	CAAA	TG.	гтс		,	_	_				٠~١	$\sim$													ر۔				
GT/	ATG	GTT	ACA	AAG	734	+ (	SC	9	J	, O	N	Ũ,	,4	$\mathcal{I}$																

HAI DQ59	
AGTGTTTATGGGACTAAAAAACTTTTAACACCTTTTTAGGGGAAATATTTTGGTCCTATACAAAACATGTAAATATGCTTTATTACTTTC	
TCACAAATACCCTGATTTTTTGAAAATTGTGGAAAAATCCCCTTTATAAAACCAGGATATGTTTTGTACATTTATACGAAATAATGAAAG	
ATTTTCTGACCCTGCTGTAAACTACTGCAACCCTCACATCCCTCAAAGGGACTTTTATGTCAAACTCTTCTGTTTCTCCAAATATAAGGA	180
TAAAAGACTGGGACGACATTTGATGACGTTGGGAGTGTAGGGGAGTTTCCCTGAAAATACAGTTTGAGAAGACAAAGAGGTTTATATTCCT	
AAAAAGACTAAAGCAAGAGATCTGGCAGTTGAAAATTGTGGGGAAAGAGAGATTTGTATGGGCACTGTATCTATGAAATACCTCATACTTAC	- 2/0
TTTTTCTGATTCGTTCTCTAGACCGTCAACTTTTAACACCCTTTCTCTTAAACATACCCGTGACATAGATACTTTATGGAGTATGAATG	i
GTTTACATGTTTTCCTAACTTTTGTATTTTCTTGTATAGCCACCTAGAGAATTCTTCATAGATTAAGAACTACAGTTTTCACCACTTA	- 360
CAAATGTACAAAAGGATTGAAAAAACATAAAAAGAACATATCGGTGGATCTCTTAAGAAGTATCTAATTCTTGATGTCAAAAAGTGGTGAAT	•
ACATAAGTAAAACAAAGTCCTTCATAATTTAACCATTAGCATCTTTGGCCAAACCAAAATAAAGAAAAGCATCTTCTCCTAGTTGTGTGT	
TGTATTCATTTTGTTTCAGGAAGTATTAAATTGGTAATCGTAGAAACCGGTTTGGTTTATTTCTTTTCGTAGAAGAGGATCAACACACAC	4
GGGCAACAGAAACAAGTTAAGGAAACAAAAATACTTATATATA	
CCCGTTGTCTTTGTTCAATTCCTTTGTTTTTATGAATATATGTGTCTTGTTTTTATTACAAGAAAAATACGTTTAGGGGACACTTTT	7
AAAATTITCAATGITTAAAAAAAAAAAAAAAAAAAAAAAA	
TITTAAAAGTTACAAATTITTTTTTTTTTTTTTTTTTTTT	

															48	3													Pa	age 1
	HHFEIK40  TCGGCACGAGCTGCGGGGGGGCGGGGGGGCCCCGGGAGCCGCGCTCTCAGTCTCTAGGCGCAGTCCCTTCGCCGCTTCCGG  AGCCGTGCTCGACGCCCGCCGCCGACCCGCCGGGGCCCTCGGCGCGAGAGTCAGAGAGATCCGCGTCAGGGAAGCGGCGAAGGCC																													
TTC	GGC	ACG	AGC	TGC	GGG	CGG	TGGC	GCG	SCT	GGG	CGGC	ccc	GG	SAG	CGG	CGC	TCT	CAG	TCT	CTC	TAG	SCG	CAG	rcc	TT	CGCC	CGC 1	TTCC	GG	
AAG	CCG	TGC	tcg.	ACG	CCC	GCC	ACC	GCC	CGA!	CCC	SCC	GGG	GC C	CTE	GGC	GCG.	AGA	GTC	AGA	GAG	ATC	CGC	GTC	AGG	GAA	GCG	GCGA	AAGG	CC.	90
AGC	ccc	TGG	ÇAG	GGC	CCA	GAA	GCCA	ATG	300	CAC	TAT	٩AG	ACT	GAG	CAG	GAC	GAC	TGG	CTG	ATC.	ATC	TAC	TTG	AAG	TAT	TTA	STCI	TTTG	TC	180
TCG	GGG	ACC	GTC	CCG	GGT	CTT	CGG	TAC	CGG	G T G	ATA	TC	TGA	CTC	GTC	CŤG	CTG	ACC	GAC	TAG	TAG.	ATG.	AAC	TTC.	ATA	AAT	GAGA	AAAC	AG	100
								М	Α	н	Y	K	Т	E	Q	D	D	W	L	l	i	Υ	L.	K	Υ	Ļ	Ŀ	F	V	
TTC	AAC	TTC	TTC	TTC	TGG	GTC	GGG	GGA	GCA	GCCI	GTC	TG	GCT	GTG	GGC.	ATC	TGG	ACC	ÇTG	GTG	GAG	AAG	AGT	GGC	TAC	CTC.	AGC(	GTCC	TG	
AAG	TTG	AAG	AAG	AAG	ACC	CAG	CCC	CCT	CGT	CGG	CAG	GAC	CGA	CAC	CCG	TAG	ACC	TGG	GAC	CAC	СТС	TTC	TCA	ccg	ATG	GAG	TCG	CAGG	AC	270
F	N	F	F	F	W	٧	G	G	Д	А	٧	L	Α	٧	G	ĺ	W	T	L	٧	Ε	K	S	G	Υ	L	s	٧	L	
GCC	TCC	AGC	ACC	7 7 7	GCC	GCC	TCC	GCC	TAC	ATC	CTC.	ATC	TTT	GCG	GGC	GTA	CTT	GTC	ATG	GTG	ACC	GGC	TTC	CTG	GGC	TTC	GGT	GCCA	ATC	
			<del></del>										+					+	÷				+			<del></del>		CGG1		360
Α	s	s	T	F	Α	А	s	Α	Υ	i	L	I	F	Α		٧	L	٧	М	٧	Т	G	F	L	G	٣	G	A	Į	
стс	TGG	GAG	CGG	AAG	GGC	TGC	CTC	TCC.	AÇG	TAT	TTC	TGC	CTG	TTG	CTC	GTC	ATC	TTC	CTG	GTT	GAG	CTG	GTG	GCG	GGA	GTC	CTG	GCCC	CAT	
GAG	ACC	СТС	GCC	TTC	CCG	ACG	GAG	AGG	TGC	ATA	AAG	A C G	GAC	AAC	GAG	CAG	TAG	AAG	GAC.	CAA	стс	GAC	CAC	CGC	CCT	CAG	GAC	CGGC	<del></del>	450
	١.,	_	R	¥	G	۲		s	т	Y	t.	۲	1	ı	L	v	ı	F	ł	v	F	1	٧	Α	G	v	L	Α	н	
GTG	" TAT	TAC		AGG		AGT	GAT	-	CTG	AAG	CAG	CAC	TTG	AAC											GAG	CAC	. – GCA	GATO		
CAC	ATA	AATG	GTC	TCC	GAC	TCA	СТА	CTT	GAC	TTC	GTC	GTG	AAC	TTG	GCC	TGA	GAC	CGA	CTC	TTG	ATG	ССС	GTC	GGC	стс	GTG	CGT	CTAC	GTG	540
٧	Y	Y	۵	R	L	S	D	Ε	L	K	Q	н	L	N	R	ī	L	А	Ε	N	Υ	G	۵	ρ.	ε	н	Α	D	н	
GCC	TCA	AGTG	GAC	CGA	стс	CAG	CAG	GAT	TTC	AAG	TGC	TGC	GGA	AGC	AAC	AGC	TCA	GCC	GAC	TGG	CAG	CAC	AGC	ACG	TAC	ATC	CTG	TTG	cee	
CGG	AGI	CAC	CTG	GCT	GAG	GTC	GTC	CTA	AAG	TTC	ACG	ACG	CCT	TCG	TTG	TCG	AGT	CGC	CTG	ACC	GTC	GTG	TCG	TGC	ATG	TAG	GAC	AAC	GCC	630
А	s	٧	0	R	L	Q	a	D	F	K	С	С	G	S	N	S	s	Α	D	W	Q	н	s	T	Υ	t	L	L	R	
GAC	GCC	GAG	GGC	cac	CAG	GTG	CCC	GAC	AGC	TGC	TGC	AAG	ACA	GTG	GTG	GCG	cgc	TGC	GGC	CAG	CGG	GCC	CAC	ccc	TCC	AAC	ATC	TAT	AAG	
сто	CGC	GCTC	CCG	GCC	GTC	CAC	GGG	CTG	TCG	AÇG	ACG	TTC	TGT	CAC	CAC	CGC	GCG	ACC	CCG	GTC	GCC	CGG	GTG	GGG	AGG	TTG	TAG	ATA		720
Ε	Α	Ε	G	R	Q	٧	Р	O	S	С	С	K	т	٧	٧	A	R	С	G	Q	R	Α	н	Р	s	N	l	Y	K	
GTO	GAG	GGGA	AGGC	TGC	стс	ACC	AAG	CTG	GAG	CAG	TTC	CTG	GCC	GAC	CAC	сте	сто	CTI	ATG	GGG	GCA	GTG	GGC	ATC	GGG	GŢG	GCC	TGC	СТĢ	0.0
ÇAC	CTO	CC1	1000	ACC	GAG	TGG	TTC	GAC	CTC	GTC	AAG	GAC	CGG	CTG	GTG	GAC	GAC	GAA	TAC	ccc	CGT	CAC	CCG	TAG	ccc	CAC	CGG	ACG	GAC	810
٧	Ε	G	G	С	L	Т	. K	L	Ε	a	F	L	Α	0	н	L	L	L	М	G	Α	٧	G	i	G	٧	Α	С	L	
CAC	SAT	CTG	GGG	ATO	GTT	CTC	ACC	TGC	TGC	TTG	CAC	CAG	AGG	CTC	CAG	ccc	CAI	ודדו	TAC	TAA	TGG	CAA	CCA	сст	CCT	CTT	CCA	ACT	GCC	<b></b>
								-															-					TGA	+	900
a	ı	С	G	М	٧	Ł	T	С	С	L	н	Q	R	L	a	R	н	F	Y	į	(S	ي س	I	D	· N	10	`.d	26	)	

CCTCAAGACAACATGTGGCACATGCCATCTGCAAGG
GGAGTTCTGTTGTACACCGTGTACGGTAGACGTTCC

936 SCO IDNO: 11

BNSDOCID: <WO\_\_\_9831799A3\_IA>

					HGBGV89																									
AGC	ГΤА	CTT	ŢCA	стс	ACC	GÇC	TGT	ССТ	TCC	TGA	CAC	CTC	ACC.	ATG	TGT	AÇG	GGA	AAA	TGT	GCC	CGC	TGT	GTG	GGG	СТС	TÇC	CTC.	ATTA	4CC	90
TCG	AAT	GAA	AGT	GAG	TGG	cĠĠ	ACA	GGA	AGG	ACT	GTG	GAG	TGG	TAC	ACA	TGC	ССТ	TTT	ACA	CGG	GCG	ACA	CAC	CCC	GAG	AGG	GAG	TAA		90
														М	С	Т	G	K	С	Α	R.	С	٧	G	L	s	L	I	τ.	
ר דר:	ran	CTC	GTC	TGC	ΔΤΤ	ere	GCC.	ΔΔ۲	פרנ	רדר	r Tai	rtai	2 T A :	ררד	ΔΔΤι	ccc	G A G	Δει	TCC	TGG	ΔΓΓ	ΔΔΓ	۸۲۲	ልልሮ	<b>Γ</b> Δ Τ	רדר	46C	TTG	~ ^ ^	
			+	+		<del></del>		<del></del>	<del>+</del>							+		+					+			-+		AAC	<del></del>	180
GAG.	400	970	CAU	~00	1100	CAC	cou			محم	unc.	<b>U</b> A C 1	-71	uun	110			100	700	~~~			,	,,,	G 1 7	iana	100.	776	311	
L	С	L	٧	С	i	٧	Α	N	Α	L	L	L	٧	Ρ	Ν	G	Ε	Τ	S	W	T	N	T	N	Н	L	S	Ĺ	a	
GTC	TGG	стс	ĄTG	GGC	GGC	TŢC	ATT	GGC	GGĢ	GGC	CTA.	ATG	ŞΤΑ	CTG	TGT	CÇA	GGG	ATT	GCĄ	GCC	GTT	CGG	ĢCA	GGG	GGC	:AAG	GGC	TGC	TGŢ	270
CAG	ACC	GAG	TAC	CCG	CCG	AAG	TAA	CCG	ccc	CCG	GAT	TAC	CAT	GAC	ACA	GGT	ССС	TAA	CGT	CGG	CAA	GCC	CGT	ССС	CCC	TTC	CCG	ACG	ACA	270
v	W	L	М	G	G	F	ī	G	G	G	L	М	٧	L	С	Р	G	i	Α	Α	٧	R	Α	G	G	к	G	С	С	
				-01				T00							***	***	T.C.O.	000		000		C T T	007	000						
			+	+		-+-		<del></del>	+		<del></del>					-+-		+	<del></del> +				+	+		TAC				
LLA	LGA	LLL	ALG	ALA		III	انا انا	ALG	166	IAC	GAL	. فا با فا	AGL	LAG	AAG.	AGG	AGL	LGL	AAG	LLL	LAL	GAA	LLA	466	IAU	1A 1 G	ACG			
G	Α	G	С	С	G	N	R	С	R	M	L	R	S	٧	F	S	S	Α	F	G	٧	L	G	Α	i	Y	С	L		
GTG	тст	GGA	GCT	GGG	стс	CGA	AAT	GGA	ССС	AGA	TGC	TTA.	ATG	AAC	GGC	GAG	TGG	GGC	TAC	CAC	TTC	GAA	GAC	ACC	GC	:GGA	GCT	TAC		
CAC	AGA	CCT	CGA	ССС	GAG	GCT	TTA	CCT	GGG	TCT	ACG	AAT	TAC	TTG	ccg	CTC	ACC	CCG	ATG	GTG	AAG	CTT	CTG	TGG	CGC	CCT	CGA	ATG		450
٧	s	G	А	G	L	R	N.	G	Р	R	С	L	М	N	G	Ε	W	G	Υ	н	F	Ε	D	Т	А	G	А	Y	L	
								***													c = c	+ + ^	T.C.O.	C T C	C T (				T.C.C	
			+					+	-+				+	+				+	+				+					GCC.	+	540
GAG	116	GLG	IGA	LIA I	ALL	LIA	GLL	ALG	L 1 L	الالالا	فافافا	ББА	نانانا	LAL	LAG	GGG	ALC	IIA	LAC	الال	GAG	AAG	AGL	GAL	GAC	.CAC		CGG.	AGG	
Ł	N	R	Τ	L	W	D	R	С	E.	А	P	ρ	R	٧	٧	P	W	N	٧	T	L	F	S	Ļ	L	٧	A	Α	S	
TGC	CTG	GAG	ATA	GTA	CTG	TGT	GGG	ATC	CAG	CTG	GTG	AAC	GCG	ACC	ATT	GĢT	GTC	TTC	TGC	GGC	GAT	TGC	ĄGG	AAA	AAA	\CAG	GAC	ACA	CCT	630
ACG	GAC	стс	TAT	CAT	GAC	ACA	ССС	TAG	GTC	GAC	CAC	TTG	CGC	TGG	TAA	CCA	CAG	AAG	ACG	CCG	CTA	ACG	TCC	TTT	TT	rgto	CTG	TGT	GGA	630
С	L	ε	I	٧	L	С	G	ı	a	L	٧	N	A	Т	ı	G	٧	F	С	G	٥	С	R	K	K	a	0	Т	Р	
					_																		. <del>.</del> .					CTC	T.C.A	
	<del></del>		+			<del>}</del>		+					+	+		<del></del>		<del></del>	<del>+</del>				+					CTC		720
											CGA	GGA	AGG	ACC	IGC	GAG	IGA	GGG	IAAC	GAG	CGA		IAI	116	ALI	3 A A A	ال العالما	GAG.	AGI	
Н	. (	S	eG	) J	D	N	0:	Ü	+	)																				
AAA	AAA	AAA	AAA	AAA	AAC	: _ =		10	00	<u>-</u>	<del>-</del> -	フィ	(~	٠ (	1	7		٠												
TTT	<del></del>	TIT	<del> </del>	777		73	8	ت	Y	ν,	レし	ノバ	4C	′ .	احد	J														

Page 1

														STTG															810
TTTI	cc.	ATTA	TCA	AGA	CAT	GGC	CAT	СТА	тст	ΆΑΑ	TAT	TAI	TATO	CAAC	TGT	GTT	AGA	CTT	GAG	GGC	AAT	ATT	GAA	AAT	GAT	GGT	GCT.	ттс	
I	R	٧	٧	м	۵	L	s	ĸ	i	L	С	G	s	Y	s	٧	ī	F	Q	Р	G	I	[	. (	SC	0	ID	) N	10;
GTAG	TCI	CAT	CAG	TAC	GTT	GAT	AGG	TTC	TAT	GAC	ACA	CCT	TCG	ATA	AGT	CAC	TAG	AAG	GTC	GGA	ССТ	TAG	TAA	ACT	TAT	TCC	TGT		,20
CATO	AGA	AGTA	GTC	ATG	CAA	C T A	тсс	AAG	ATA	CTG	TGT	ĞGA	AGC	TAT	TÇA	GTG	ATC	TTC	CAG	ССТ	GGA	ATC	ATT	TGA	ATA	AGG	ACA	AAA	720
С	L	ε	Ρ	Α	н	٧	٧	Ε	W	N	ı	I	L	F	S	I	L	I	T	L	S	G	L	Q	٧	i	1	С	
CACG	GAC	CTT	GGA	CGT	GTA	CAA	CAC	стс	ACC	TTG	TAG	TAA	AAT	AAA	AGG	TAA	GAG	TAT	TGG	GAG	TCA	ccc	GAA	GTT	CAC	TAG	TAG	ACG	
GTGC					<del>- i -</del>					_+-		+	<del></del>				+					+			-				630
G	Ρ	Y	C.	R	T	L	D	G	W	ε	Y	A	F	Ε	G	T	Α	G	R	F	Ļ	T	0	S	S	I	W	l	
TCCC	GGT	ATA	ACG	GCG	TGG	GAA	CTA	CCG	ACC	CTC	ATA	CGA	AAA	CTT	CCG	TGA	CGA	CCT	GCA	AAG	GAA	TGT	CTA	AGA	TCG	TAT.	ACC:	TAA-	-
AGGG					<del></del>		+					+										1					,		540
. Y	٧	T	L	L	S	I	I	F	S	S	L	G	i	Α	F	<b>S</b>	G	Y	C	L	٧	ı	S		L	G	L,	٧	
HAIA	LAL	. 1 14 1	GAL	GAL	AG I	IAA	IAG	AAA	AGA	AGG	g A G				-	AUA1	1	., .,		,		, ,			, , , , ,	- CA	1		•
ATAT FATA		-1			<del></del>					-,-					<del>- ; -</del>			<del></del>				<del></del>			<del></del>				450
,			_	, . <del>.</del> .						_			_		···	 TCT:				- - T C (	· TC	A T C	TCT	-	TTC	CCT	~ T T C	: T.C	
		м		r	v	т	т	v	,				Ε		N	N	N	Y	ĸ	С	С	۵	s	Ε	N	С	S	K :	- 1
STAG		+			<del></del>			$\overline{}$				<del></del>										+	-					_	360
CATC	ATC	ATC	CTT	Λ T Λ:	CTA	ΛΓΛ	۸۲۵	GTT	rtt	CTG	CTA	c t G	GAG	ΔΔΤ	ΔΔΤ.	ΔΑΟ	AAC.	TATA	AAA'	TGT	rge	CAG	AGT	GAA.	AAC	TGC	AGC A	AAA	
L	Y	F	ρ	N	G	۵	Т	,S	Υ	Α	S	S	N	K	L	ī	N	Υ	٧	W	Υ	F	ε	G	ı	С	F	s	
TAAC	ΑТА	AAG	GGC	TTA	CCC	GTT	TGA	AGG	ATA	CGT	AGG	TCG	TTA	TTT	GAG	TGG	FTGA	TGC	AC	ACCA	ATA,	AAA	CTT	CCT	TAG.	AC A	AGA	GT	270
ATTG	TAT	TTC	CCG.	4ΑΤ	GGG	CAA	ACT	TCC	TAT	GCA	TCC.	AGC	AAT	AAA	CTC	ACCA	AACI	TAC	TG'	rggi	ΓΑΤ	TTT	GAA	GGA,	ATC	TGT:	TCT	CA	
			M	G	S	R	K	С	G	G	С	L	s	С	L	L	i	Р	L	Α	L	W	s	ĺ	i	٧	N	ı	
GTC	CCG	TTT	TAC	CC.	AGA	GCC	TTC	ACA	CCT	CCG.	ACG	GAT	TCA	ACA.	AAC	GACT	(AAC	GCG	AA	CGTC	AA	ACC.	TCA	TAT:	TAG	CACT	TGT	ΑŤ	
CAG	GGC	AAA.	ATG	GG.	тçт	CGG	AAG	TGT	GGA	GGC	TGC	ÇTA	AGT	TGT	TTG	TGA	TTC	cgo	TTO	CAC	: T T	rgg,	AGT/	ATA	ATC	STGA	ACA	TA	180
TGC	161	C I A	ACA.	300	3,70		<u> </u>	010							• • • • • • • • • • • • • • • • • • • •												•		
							Δ :	ΔιΔ	4614	ı ι Δ :	41:1:1	1111	A   - A	144 A I	5 I A1		I I A L		AL I	166	. L. Uz /	l Li A l	. !!:!	a La A L		LAL	. L. A.A	UA	

ACGTAAACCACAAATAAACATTTTTTAAACGTCAGGAGTGACGTTCATATGGTGGGAAGGTAAATCATACAAAAAATTCATTAT

WO 98/31799 PCT/US98/00959

52

Paige 2

AGAAACCTTTATGCAATATGTATATTGCAACATTATTTAATATTCTGGAAAATTGGAAACACCCCAAAATTCTAACTCAAA

TCTTTGGAAATACGTTATACATATAACGTTGTAATAAAATTATAAGACCTTTTAACCTTTGTGGGGTTTTAAGATTGAGTTT

BNSDOCID: <WO\_\_\_9831799A3\_IA>

Page 1

# HJACE54

TT	TGI	G	GAG	ĢGC	AG	CAG	AGA	GT.	ACC	:CA	GCT	GGA	CA	TCC	TTT	CC	TGC	TG.	AŢG	AGC	ccc	AG	GCT	GGA	GG	TGC	CC	TGC	TCA	CAT	GCT	CTT	222	90
AA	AC.	C	CTC	CCC	тс	GTC.	TCT	CA	TGC	3GT	CGA	CCT	GT	AGG.	٩AA	GG	ACG	AC	TAC	TCG	GGC	STC	CGA	СТ	CC.	ACG	GG	ACG	AGT	GTA	kC G A	GAA	GGG	30
																			М	S	Р	R	L	Ε		٧	P	С	s	н	А	L	Р.	
				i		TGG									<del></del>		—		<del></del>		+		+							<del></del>		<del></del>	$\rightarrow$	180
GT	ccc	A	GAG	AG	GG	ACC	CGT	CC.	AG1	ΓAG	TAT	CAT	GC	CCC	TGA	CC	AGA	AAC	GTT	CTC	GG	CTT	CGT.	AΑΑ	ΑT	GAC	AC	TCG	GAC	TCC	CTG:	GTC	CGA	
۵	(	3	L	s	Ρ	G	Q		٧	I	i	٧	R	G	£	-	٧	L	Q	Ε	Ρ	K	Н	F	•	Т	٧	S	L	R	D	Q	Α	
GC	CCA	A T	GCT	cc.	rgr	GAC	ACT	CA	GG(	GCC	TCC	TTC	GC	AGA	CĄG	A A	CTC	TG	GÇC	TGG	AT(	стс	çcg	CTG	GG	GG	AG	AAG	AAA	CTC	SATO	TCA	GCĊ	270
CG	GG	ГΑ	CGA	ĠG,	CA	CTG	TGA	GΤ	cca	GG	AGG	AAC	CG	TCT	GTC	: T T	GAC	SAC	CGG	ACC	TAC	GAG	GGC	GAC	cc	CCC	GTC	TTC	TT	GAC	TAG	AGT	CGG	
А	ı	4	Α	ρ	٧	T	L		R	Α	S	F	А	ם	F	₹	Т	L	Α	W	i	S	R	W	ı	G	۵	к	κ	L	i	s	Α	
۲۲	ст.	דר	רדר	тт.	ΓΤΔ	CCC	CCA	GA	GA.	TTC		GAC	GT	GCT	GC 1	rcc	TGI	ГTС	CAG	GAG	GG	AGG	GCT	GAA	\GC	TG	3C G	стс	:AA1	rggr	GÇAC	GGG	CTG	
_				+		GGG				+									<del></del>		<del></del>		+										$\rightarrow$	360
_		_		_		_			_	_	_	_						_	^	_	_	^		v	,		٨		NI.	C	0	C	,	
Р	1	F	L	F	Y	Р	Q		ĸ	۲	۲	£	٧		L	-	L	۲	u	£	G	G		N	•	_	A	L	IN	u	u	u	_	
GG	GG	cc	ACC	AG	CAT	GAA	CCA	GC	AG	GCC	CTG	GAC	C A	GCT	GÇ (	GG	AGG	стс	CGG	ATO	AG	TGG	AAG	TGT	rcc	AG	СТС	TAC	TG	FGT	CCAC	TCC	TGA	450
CE	CC	GG	TGG	TC	GTA	CTT	GGT	CG	TC	CGG	GAC	CTO	GT	CGA	cĠ	cc	CTC	GAG	GCC	TAG	TC.	ACC	TTC	ACA	AGG	TC	GAG	ATO	ACA	4CAI	GGTO	AGG	ACT	: TO
G		A	т	s	М	ı N	a	l	a	Д	L	ε	C	ı L	F	₹	E	L	R	i	s	G	s S	١	,	a	L	Y	С	٧	Н	s		NO: 29
ΔΩ	GA.	TG	GTT	cc.	AGG	AAA	TAC	CG	CA	GAA	AAAC	CAAC	SAG	TCA	GC	CAC	TC	ccc	AGC	GCC	cc.	ACT	CTC	сто	cc	CT	CAT	TAA	AC	CAT	CCAC	CTG	AAC	
_				<del></del>		TTT				+					<del></del>								+				+							540
																													•					
ΑC	CA	GC	A C A	TC.	AGG	GCC	TGG	TT	CA	ССТ	гсто	GGG	GT C	CACG	ΑĢ	AC1	ΓGΑ	GTC	TAC	CAGO	AG	стт	ŢGG	GCC	CTG	AG	GGA	AGG	CA	CAA	GAG	rgc A	AAG	630
TG	GT	CG	TGI	AG	TCC	CGG	ACC	AA	GT	GGA	4GAC	CCC	AC	STGC	TC	TG	ACT	CAG	ATO	STC	:TC	GAA	ACC	CGC	GAC	TC	ccı	TCC	GT	377	CTC	ACGT	TTC	000
G T	TC	СТ	CGA	AC	тст	GCA	CCI	TC	CT	CCA	ACC/	AGG	4GC	CTG	GG.	A T	ATG	GCT	rcc/	ATC	rgc	CTT	CAG	GGG	CCT	GG	AC 1	rgc/	ACT	JAC.	AGA	GCA	AGT	720
CA	AG	GΑ	GCT	ΤÌG	AGA	CGT	GGA	AG	GA	GGT	rggi	TCC	TC	GGAC	CC.	TA'	TAC	CGA	AGG.	TAGA	ACG	GAA	GTC	CCC	GGA	CC	TGA	ACC.	ΓGΑ	GTG	TCT	CGT	TCA	
<b>C</b> 1		T A	C 4 (	~ T A	۸ ۲ ۸	AAAG	A T A	CT	. c c	A A /	4 A T /	۸ C A	ν Т (	ece T	Τ.	۸ ۸ ۱	2 4 4	TGI	ree:	TCA:		ΔΤΊ	CTT	ΤΔ.	TTA	TT	ΤΔ.	r r r z	ΔΤΤ	TGT	GGTI	CAAA	TAA	
		-				TTTC	<del>i</del>												-+				+				+							810
C.A	AC	A I	C 1 (	Aί	161	1 1 L	IAI	uА	نا دا،	111	I I A	i Gri	ıΑί	4 فا ما د	1 A	1 11	. I I	MLF	4661	MG 17	4 H H	. I A F	10 A P	A 17	44 i	MA	n 1 /		. ~ ^		5571			
Αī	ΓΔΑ	ΑT	`ΑΑ(	GGT	TAI	ГТТА	ттт.	ΓΑΑ	AAA	AA	ΑΑΑ	AAA.	AA	444	AA.	AΑ	ААА	AAA	<b>4</b> AA	AAA	AΑ		_ //	<b>-</b> -	<u>_</u>	—	· _		<u> </u>	14	†_	١		
-		-		<del></del>		AAAT									-				-+-	TTT	<del></del>	869	5 C	50	ب	T	(سا-	(N	<i>\( \)</i>		٠,	'		

HROAD63

Päge 1

CACGAGAGACGACATC	AGAGATGAC	GGACA	GCATT	GCTG	стсс	TTGC	AGC'	CTG	GCT	GTG	GCT	ACA	GGG	CCA	GCC	CTT	ACC	:CTG	cac	TGC	90
GTGCTCTCTGCTGTAG	CTCTACTO	CCTGT	CGTAA	CGAC	GAGG	AACG	TCG	GGAC	CGA	CAĊ	CGA	TGT	CCC	GGT	CGG	GAA	TGC	GAC	GCG	ACG	30
	M F	R T	A L	L	Ļ	L A	. Α	L	Α.	٧	Α	Т	G	Ρ	A	L	T	L	R	С	•
ACGTGTGCACCAGCTC	CAGCAACT(	GCAAG	CATTC	TGTG	GTCT	GCCC	GGC	CAGO	TCT	cgç	TTC	TGC	AAG	ACC	ACG	AAC	AÇA	GTG	GAG	CCT	180
TGCACACGTGGTCGAG	GTCGTTGAG	CGTTC	GTAAG	ACAC	CAGÁ	CGGG	ccè	STCG	AGA	GCG	AAG	ACG	TTC	ŤGG	TGC	TTG	TĠI	[CAC	CTC	GGA	, ,
H V C T S S	S N (	с к	н s	٧	٧	C F	РА	s	s	R	F	С	K	T	T	N	T	٧	Ε	Ρ	
TGAGGGCTTCCCCGAA	AGTCTGGG	ACCAĢ	GTCCA	GGTG	GGCA	TGGA	ATG	CTGA	TGA	CTT	GGA	GCA	GGC	çcc	ACA	GAC	cco	CACA	GAG	GAT	270
ACTCCCGAAGGGGCTT	TCAGACCC	TGGTC	CAGGT	CCAC	CCGT	ACCI	TAC	GAC 1	ACT	GAA	CCT	CGT	CCG	GGG	TGT	CTG	GĠ	STGT	стс	CTA	
L R A S P K	v w (	o	v a	٧	G	м 8	. c		(3	Sca		II	1 (	40	); .	3C		)			
AAGCCACCCACAGAG	GATGCAGC	CCCCA	.GCTGC	ATGG	AAGG	TGG	AGGĄ	CAGA	AGC	CCŢ	GTG	GAT	ccc	ÇGG	ATI	TCA	CĄ	CTCC	TTC	TGŢ	20
TTCGGTGGGGTGTCTC	CTACGTCG	GGGGT	CGACG	TACC	TTCC	ACC	rcct	GTC	TCG	GGA	CAC	CTA	GGG	ĠCC	TAA	AGT	GT	GAGO	SAAC	ACA	36

HMW6546

Page 1

CCGGCCTCAAGATCAGCATGGCCCAGAACTTGAAGGACTTGAACGACTTGGCGGCCCCATCTCCGCCCGGGGCCCCGGGGCAGGGCCCCGGGGCAGGGCCCCGGAGGCCCCGGAGGCCCCGGAGGCCCCGGGGCCCCGGGGCCCCGGGGCCCCGGGGCCCCGGGG
HAAQNLKODLAAGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGG
M A Q N L K D L A G R L P A G P R G M G T A L K  CTGTTGCTGGGGGCCCGCGCGCCCCTACGGTGTGCCGAATCTGTGTTCACCCGTGGAAGGCGGCCCAAGGCCATCTTCTTCAATCGG GACAACGACCCCCGGCGCGCCGCGGCCCTACGGTGTGCCGAATCTGTGTTCACCCGTGGAAGGCGGGCACAGAGCCATCTTCTTCAATCGG GACAACGACCCCCGGCGCGCGCGCGCGCGCCACCGGATGCCACACGGCCTTAAGACACAAGTGGCACCTTCCGCCCGTGTCCGGTAGAAGAAGATAGCC  L L L G A G A V A Y G V R E S V F T V E G G H R A I F F N R  ATCGGTGGAGTGCAGCAGGACACTATCCTGGCCGAGGGCCTTCACTTCACTTCAGGATCCCTTGGTTCCAGTACCCCATTATCTATGACATTCGG TAGCCACCTCACGTCGTCCTGTGATAGGACCGGCTCCCGGAAGTGAAGTCCTTGGGTTCCAGTACCCCCATTATCTATGACATTCGG TAGCCACCTCACGTCGTCCTGTGATAGGACCGGCTCCCAGAGAGCCTTCACTTCAGGAACCCAAGGTCATGGGGTAATAGATACTGTAAGCC  I G G V Q Q D T I L A E G L H F R I P W F Q Y P I I Y D I R  GCCAGACCTCGAAAAATCTCCTCCCCCTACAGGCTCCAAAGACCTACAGATGGTGAAATATCTCCCTGCGAGTGTTGTCTCGACCCAATGCT CGGTCTGGAGCTTTTTAGAGGAAGGGGATGTCCGAGGTTTCTGGATGTCTACCACTTATAGAGGGACGCTCAAACAGAGCTGGGTTACGA A R P R K I S S P T G S K D L Q M V N I S L R V L S R P N A  CAGGAGCTTCCTAGCATGTACCAGCGCCTAGGGCTGGACTACGAGGAACGAGTGTTGCCCGCTCCATTGTCAACAGAGGTGCTCAAGAGTTGT GTCCTCGAAGGATCGTACCAGCGCCCTAGGGCTGGACTACGAGGAACGAGTTGTCCCACCAACCA
CIGTIGCIGGGGCCCGCGGCCCCGGGCCCCGGTGCCTACGGTGGCCGAATCTGTGTTCACCGTGGAAGGCGGCACAGAGCCATCTTCTTCAATCGG  GACAACGACCCCCGGCCGGCCGCGGCACCGGATGCCACACGGCGTTAGACACAAGTGGCACCTTCCGCCCGTGTCTCGGTAGAAGAAGATTAGCC  L L G A G A V A Y G V R E S V F T V E G G H R A I F F N R  ATCGGTGGAGTGCAGCAGGACACTATCCTGGCCGGAGGCCTTCACTTCAGGATCCCTTGGTTCCAGTACCCCATTATCTATGACATTCGG  TAGCCACCTCACGTCGTCCTGTGATAGGACCGGCTCCCGGAAGTGAAGTCCCTTGGTTCCAGTACCCCATTATCTATGACATTCGG  TAGCCACCTCAACGACTCCTGTGATAGGACCGGCTCCCGGAAGTGAAGTCCTAGGGAACCAAGGTCATGGGGTAATAGATACTGTAAGCC  I G G V Q Q D T I L A E G L H F R I P W F Q Y P I I Y D I R  GCCAGACCTCGAAAAATCTCCTCCCCTACAGGCTCCAAAGACCTACAGATGGTGAATATCTCCCTGCGGAGTGTTGTCTCGACCCAATGCT  CGGTCTGGAGCTTTTTAGAGGAGGGGGATGTCCGAAGGTTTCTGGATGTCTACCACTTATAGAGGGACGCTCACAACAAGAGCTGGGTTACGA  A R P R K I S S P T G S K D L Q M V N I S L R V L S R P N A  CAGGAGCTTCCTAGCATGTACCAGGGCTGAGGCTGGACTACGAGGAACGAGTGTTGCCGTCCATTGTCAACACAAGAGCTGGGTTACCAC  G E L P S M Y Q R L G L D Y E E R V L P S I V N E V L K S V  GTGGCCAAAGTTCAATGCCTCACAGCTGATCACCCAGGGGCCCAAGGACCAACAAGTGCTCCACAAGAGTCCCGGTTCCACAA  CACCGGTTCAAAGTTACCGGAGTGCACCAACAAGAGCCCAAGGGCCCAAGGAC  V A K F N A S Q L I T Q R A Q V S L L I R R E L T E R A K D  TTCAGCCTCATCCTGGATGATGTGGCCATCACAAGAGCTGAGCTTTAGCCCGAGAGTACACAGGTGCTCTAGGAAGCCCAAACAAGTGGCCCAG  AAGTCCGGAGTAGGACCTACTACACCGGTTAGTCCCGACTGAGCTTTAGCCCGAGAGTACACAGCTGCTGTTAGAAGCCAAACAAGTGGCCCAG  AAGTCCGGAGTAGGACCTACTACACCGGTTAGTCCCGACTGAACCAAACAAGTGGCCCAACAAACA
CAGGACCTCCGAAAAATCTCCTCCCCTACAGGCTCCCAAAGACCTACCAGGGTGAATATCTCCCTGGGAGGTGTTCTCGGCCAATGCT  ATCGGTGGAGCTTTTTTAGAGGAGGGGTCCCACAGGGTTCCCAAAGACCTACCAAGAGTGGAACCTACACGGGAACCTACCT
CAGGACCTCCGAAAAATCTCCTCCCCTACAGGCTCCCAAAGACCTACCAGGGTGAATATCTCCCTGGGAGGTGTTCTCGGCCAATGCT  ATCGGTGGAGCTTTTTTAGAGGAGGGGTCCCACAGGGTTCCCAAAGACCTACCAAGAGTGGAACCTACACGGGAACCTACCT
ATCGGTGGAGTGCAGCAGGACACTATCCTGGCCGAGGGCCTTCACTTCAGGATCCCTTGGTTCCAGGTACCCCATTATCTATGACATTCGG TAGCCACCTCACGTCGTCGTGATAGGACCGGCTCCCGGAAGGTGAAGTCCTTGGGTTCCAAGGGCTAATGGTAAGCC  I G G V Q Q D T I L A E G L H F R I P W F Q Y P I I Y D I R  GCCAGACCTCGAAAAATCTCCTCCCCTACAGGCTCCAAAGACCTACAGATGGTGAATATCTCCCTGGGGAGTGTTGCTCCAACACAGGTTTCTCGACCCAATGCT CGGTTGGAGGCTTTTTAGAGGAGGGGGATGTCCGAGGTTTCTGGATGTTCTACCACTTATAGAGGGACGCTCCAAAGACCTGGGTTACGA  A R P R K I S S P T G S K D L Q H V N I S L R V L S R P N A  CAGGAGCTTCCTAGCATGTACCAGCGCCTAGGGCTGGACTACGGAGGAACGAGTGTTGCCGTCCATTGTCAACGAGGTGCTCAAGAGTGTG GTCCTCGAAGGATCGTACATGGTCGGGGATCCCGACCTGATGCTCCTTGGTCACAACGGCGGAGCTACAGAGTGTG GTCCTCGAAGGATCGTACATGGTCGCGGATCCCGACCTGATGCTCCTTGGTCACAACGGCGGAGCTACAGAGTGTG GTCCTCGAAGGATCGTACATGGTCGGGGATCCCGACCTGATGCTCCTTGGTCACAACGGCGGAGCTGACAGAGGGCCCAAGGAC Q E L P S M Y Q R L G L D Y E E R V L P S I V N E V L K S V  GTGGCCAAGTTCAATGCCTCACAGCTGATCACCCAGCGGGCCCCAGGTATCCCTTGTTGATCCGCCGGGAGCTGACAGAGAGGGCCCAAGGAC CACCGGTTCAAGTTACGGAGTGTCGACTAGTGGGCCCCAGGGTCCCTGGTTGATCCCCCGGGGGCCCTGACAGGAGGGCCCAAGGAC  V A K F N A S Q L I T Q R A Q V S L L I R R E L T E R A K D  TTCAGCCTCATCCTGGATGATGGGCCATCACAGAGCTGAGCTTAGGCCGAAACCAAGCTGCTCTAGCACAACAACAAGTGGCCCCAG AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCCGACTCGAAATCGGCTCTCATGTGTCGACCAAACAACAAGTGGCCCCAG
TAGCCACCTCACGTCGTCCTGTGATAGGACCGGCTCCCGGAAGTGAAGTCCTAGGGAACCAAGGTCATGGGGTAATAGATACTGTAAGCC  I G G V Q Q D T I L A E G L H F R I P W F Q Y P I I Y D I R  GCCAGACCTCGAAAAATCTCCTCCCCCTACAGGCTCCAAAGACCTACAGATGGTGAATATCTCCCTGCGGAGTGTTGTCTCGACCCAATGCT CGGTCTGGAGCTTTTTAGAGGGAGGGGATGTCCGAGGTTTCTGGATGTCTACCACTTATAGAGGGACGCTCACAACAGAGCTGGGTTACGA  A R P R K I S S P T G S K D L Q H V N I S L R V L S R P N A  CAGGAGCTTCCTAGCATGTACCAGCGCCTAGGGCTGGACTACGAGGAACGAGTGTTGCCCACTTGTCAACGAGGTGCTCAAGAGTGTG GTCCTCGAAGGATCGTACATGGTCGCGGATCCCGACCTGATGCTCCTTGCTCACAACGGCAGGTAACAGTTGCTCCACGAGGTTCTCACAC  Q E L P S H Y Q R L G L D Y E E R V L P S I V N E V L K S V  GTGGCCAAGTTCAATGCCTCACAGCTGATCACCCAGCGGGCCCAGGTATCCCTGTTGATCCGCCGGGAGCTGACAGAGAGGGCCAAGGAC CACCGGTTCAAGTTACGGAGTGCCCACAGCTGATCACCCAGGGGCCCAAGGACCGGCCCTCGACTGTCTCTCCCGGTTCCTG  V A K F N A S Q L I T Q R A Q V S L L I R R E L T E R A K D  TTCAGCCTCATCCTGGATGATGGCCCATCACAGAGCCTGACCTTTAGCCGAAACCAAGTGCCCCAGGTTCCTCACAACCAGGCCCCAGGTTCCTCGGTTCCTCGGTTTCTCCCGGTTCCTC  AAGTCGGAGTAGGACCTACTACACCCGGTAGTCTCCTGAAATCGGCTCTCTATGTGTCGACAGCACAAACAA
TAGCCACCTCACGTCGTCCTGGATAGGACCGGCTCCCGGAAGTGAAGTCCTAGGGAACCAAGGTCATGGGGTAATAGATACTGTAAGCC  I G G V Q Q D T I L A E G L H F R I P W F O Y P I I Y D I R  GCCAGACCTCGAAAAATCTCCTCCCCCTACAGGCTCCAAAGACCTACAGATGGTGAATATCTCCCTGCGAGTGTTGTCTCGACCCAATGCT CGGTCTGGAGCTTTTTAGAGGGAGGGGATGTCCGAGGTTTCTGGATGTCTACCACTTATAGAGGGACGCTCACAACAGAGCTGGGTTACGA  A R P R K I S S P T G S K D L Q H V N I S L R V L S R P N A  CAGGAGCTTCCTAGCATGTACCAGCGCCTAGGGCTGGACTACGAGGAACGAGTGTTGCCCGTCCATTGTCAACGAGGTGCTCAAGAGTGTG GTCCTCGAAGGATCGTACATGGTCGCGGGATCCCGACCTGATGCTCCTTGCTCACAACGGCAGGTAACAGTTGCTCCACGAGGTTCTCACAC  Q E L P S H Y Q R L G L D Y E E R V L P S I V N E V L K S V  GTGGCCAAGTTCAATGCCTCACAGCTGATCACCCAGCGGGCCCAGGTATCCCTGTTGATCCGCCGGGAGCTGACAGAGAGGGCCAAGGAC CACCGGTTCAAGTTACGGAGTGCCCTAACAGCTGACCCAGGGGCCCAAGGACCGACC
GCCAGACCTCGAAAAATCTCCTCCCCTACAGGCTCCAAAGACCTACAGATGGTGAATATCTCCCTGCGAGTGTTGTCTCGACCCAATGCT CGGTCTGGAGCTTTTTAGAGGAGGGGGATGTCCGAGGTTTCTGGATGTCTACCACTTATAGAGGGACGCTCACAACAGAGCTGGGTTACGA  A R P R K I S S P T G S K D L Q M V N I S L R V L S R P N A  CAGGAGCTTCCTAGCATGTACCAGCGCCTAGGGCTGGACTACGAGGAACGAGGTGTTGCCGTCCATTGTCAACGAGGTGCTCAAGAGTGTG GTCCTCGAAGGATCGTACCAGGCGCCTAGGGCTGGACCTGATGCTCCTTGCTCACAACGGCAGGTAACAGTTGCTCCAACGAGTTCTCAACA  Q E L P S M Y Q R L G L D Y E E R V L P S I V N E V L K S V  GTGGCCAAGTTCAAAGTTAACGCAGCTGATCACCCAGCGGGCCCAGGTATCCCTGTTGATCCGCCGGGAGCTGACAAGAGAGGGCCAAGGAC CACCGGTTCAAGTTAACGGAGTGTCGACCTAGTGGGCCCAGGGTCCATAGGGAACAACTAGGCGGCCCTCGACTGTCTCCCCGGTTCCTG  V A K F N A S Q L I T Q R A Q V S L L I R R E L T E R A K D  TTCAGCCTCATCCTGGATGATGTGGCCATCACAGAGCTGAGGTTTAGCCGAGAGTACACAAGCTGCTGTAGAAGCCAAACAAGTGGCCCAGG AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCCGACTGTTTTAGCCGAGAGTACACAAGCTGCTTTTTCACCGGGTC AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCCGACTGTCTCCACGGGTC CACCGGTTCATCCTGGATGATGTGGCCCATCACAGAGCTGAACTAGGCGACAACCAAGTGGCCCAG AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCCGACTGTTTTTCACCGGGTC  AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCCGACTGCTAATGTGTCGACGACATCTTCGGTTTGTTCACCGGGTC  AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCCGACTGCAAATCAGGCTCCTCATGTGTCGACGACATCTTCGGTTTGTTCACCGGGTC
CGGTCTGGAGCTTTTTAGAGGAGGGGATGTCCGAGGTTTCTGGATGTCTACCACTTATAGAGGGACGCTCACAACAGAGCTGGGTTACGA  A R P R K I S S P T G S K D L Q M V N I S L R V L S R P N A  CAGGAGCTTCCTAGCATGTACCAGCGCCTAGGGCTGGACTACGAGGAACGAGTGTTGCCGTCCATTGTCAACGAGGTGCTCAAGAGTGTG  GTCCTCGAAGGATCGTACATGGTCGCGGATCCCGACCTGATGCTCCTTGCTCACAACGGCAGGTAACAGTTGCTCCACGAGTTCTCACAC  Q E L P S M Y Q R L G L D Y E E R V L P S I V N E V L K S V  GTGGCCAAGTTCAAGTGCCTCACAGCTGATCACCCAGCGGGCCCAGGTATCCCTGTTGATCCGCCGGGAGCTGACAGAGAGGGCCAAGGAC  CACCGGTTCAAGTTACGGAGTGTCGACTAGTGGGTCGCCCGGGTCCATAGGGACAACTAGGCGGCCCTCGACTGTCTCTCCCCGGTTCCTG  V A K F N A S Q L I T Q R A Q V S L L I R R E L T E R A K D  TTCAGCCTCATCCTGGATGATGTGGCCATCACAGAGCTGAGCTTTAGCCGAGAGTACACAGCTGCTGTAGAAGCCAAACAAGTGGCCCAG  AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCGACTCGAAATCAGGCTCTCTATGTGTCGACGACAACTTCTCGGTTTGTTCACCGGGTC  720
CGGTCTGGAGCTTTTTAGAGGAGGGGATGTCCGAGGTTTCTGGATGTCTACCACTTATAGAGGGACGCTCACAACAGAGCTGGGTTACGA  A R P R K I S S P T G S K D L Q M V N I S L R V L S R P N A  CAGGAGCTTCCTAGCATGTACCAGCGCCTAGGGCTGGACTACGAGGAACGAGTGTTGCCGTCCATTGTCAACGAGGTGCTCAAGAGTGTG  GTCCTCGAAGGATCGTACATGGTCGCGGATCCCGACCTGATGCTCCTTGCTCACAACGGCAGGTAACAGTTGCTCCACGAGTTCTCACAC  Q E L P S M Y Q R L G L D Y E E R V L P S I V N E V L K S V  GTGGCCAAGTTCAAGTGCCTCACAGCTGATCACCCAGCGGGCCCAGGTATCCCTGTTGATCCGCCGGGAGCTGACAGAGAGGGCCAAGGAC  CACCGGTTCAAGTTACGGAGTGTCGACTAGTGGGTCGCCCGGGTCCATAGGGACAACTAGGCGGCCCTCGACTGTCTCTCCCCGGTTCCTG  V A K F N A S Q L I T Q R A Q V S L L I R R E L T E R A K D  TTCAGCCTCATCCTGGATGATGTGGCCATCACAGAGCTGAGCTTTAGCCGAGAGTACACAGCTGCTGTAGAAGCCAAACAAGTGGCCCAG  AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCGACTCGAAATCAGGCTCTCTATGTGTCGACGACAACTTCTCGGTTTGTTCACCGGGTC  720
CAGGAGCTTCCTAGCATGTACCAGCGCCTAGGGCTGGACTACGAGGAACGAGGTGTTGCCGTCCATTGTCAACGAGGTGCTCAAGAGTGTG  GTCCTCGAAGGATCGTACATGGTCGCGGGATCCCGACCTGATGCTCCTTGCTCACAACGGCAGGTAACAGTTGCTCCACACGAGTTCCTCACAC  Q E L P S M Y Q R L G L D Y E E R V L P S I V N E V L K S V  GTGGCCAAGTTCAATGCCTCACAGCTGATCACCCAGCGGGCCCAGGTATCCCTGTTGATCCGCCGGGAGCTGACAGAGAGGGCCAAGGAC  CACCGGTTCAAGTTACGGAGTGTCGACTAGTGGGTCGCCCGGGTCCATAGGGACAACTAGGCGGCCCTCGACTGTCTCCCCGGTTCCTG  V A K F N A S Q L I T Q R A Q V S L L I R R E L T E R A K D  TTCAGCCTCATCCTGGATGATGTGGCCATCACAGAGCTGAGCTTTAGCCGAAGTACACAGCTGCTGTAGAAGCCAAACAAGTGGCCCAG  AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCGACTCGAAATCGGCTTCCATGTGTCGACGACCATCTTCGGTTTGTTCACCGGGTC  720
GTCCTCGAAGGATCGTACATGGTCGCGGATCCCGACCTGATGCTCCTTGCTCACAACGGCAGGTAACAGTTGCTCCACGAGTTCTCACAC  Q E L P S M Y Q R L G L D Y E E R V L P S I V N E V L K S V  GTGGCCAAGTTCAATGCCTCACAGCTGATCACCCAGCGGGCCCAGGTATCCCTGTTGATCCGCCGGGAGCTGACAGAGAGGGCCAAGGAC CACCGGTTCAAGTTACGGAGTGTCGACTAGTGGGTCGCCCGGGTCCATAGGGACAACTAGGCGGCCCTCGACTGTCTCTCCCCGGTTCCTG  V A K F N A S Q L I T Q R A Q V S L L I R R E L T E R A K D  TTCAGCCTCATCCTGGATGATGTGGCCATCACAGAGCTGAGCTTTAGCCGAGAGTACACAGCTGCTGTAGAAGCCAAACAAGTGGCCCAG AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCGACTCGAAATCGGCTCTCATGTGTCGACCACAACTTCTCGGTTTGTTCACCGGGTC
GTCCTCGAAGGATCGTACATGGTCGCGGATCCCGACCTGATGCTCCTTGCTCACAACGGCAGGTAACAGTTGCTCCACGAGTTCTCACAC  Q E L P S M Y Q R L G L D Y E E R V L P S I V N E V L K S V  GTGGCCAAGTTCAATGCCTCACAGCTGATCACCCAGCGGGCCCAGGTATCCCTGTTGATCCGCCGGGAGCTGACAGAGAGGGCCAAGGAC  CACCGGTTCAAGTTACGGAGTGTCGACTAGTGGGTCGCCCGGGTCCATAGGGACAACTAGGCGGCCCTCGACTGTCTCTCCCCGGTTCCTG  V A K F N A S Q L I T Q R A Q V S L L I R R E L T E R A K D  TTCAGCCTCATCCTGGATGATGTGGCCATCACAGAGCTGAGCTTTAGCCGAGAGTACACAGCTGCTGTAGAAGCCAAACAAGTGGCCCAG  AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCGACTCGAAATCGGCTCTCATGTGTCGACGACAACTTCTCGGTTTGTTCACCGGGTC
GTGGCCAAGTTCAATGCCTCACAGCTGATCACCCAGCGGGCCCAGGTATCCCTGTTGATCCGCCGGGAGCTGACAGAGAGGGCCAAGGAC CACCGGTTCAAGTTACGGAGTGTCGACTAGTGGGTCGCCCGGGTCCATAGGGACAACTAGGCGGCCCTCGACTGTCTCTCCCCGGTTCCTG  V A K F N A S Q L I T Q R A Q V S L L I R R E L T E R A K D  TTCAGCCTCATCCTGGATGATGTGGCCATCACAGAGCTGAGCTTTAGCCGAGAGTACACAGCTGCTGTAGAAGCCAAACAAGTGGCCCAG AAGTCGGAGTAGGACCTACACCGGTAGTGTCTCGACTCGACTCGAAATCGGCTCTCATGTGTCGACCAGACATCTTCGGTTTGTTCACCGGGTC
CACCGGTTCAAGTTACGGAGTGTCGACTAGTGGGTCGCCCGGGTCCATAGGGACAACTAGGCGCCCTCGACTGTCTCCCCGGTTCCTG  V A K F N A S Q L I T Q R A Q V S L L I R R E L T E R A K D  TTCAGCCTCATCCTGGATGATGTGGCCATCACAGAGCTGAGCTTTAGCCGAGAGTACACAGCTGCTGTAGAAGCCAAACAAGTGGCCCAG  AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCGACTCGAAATCGGCTCTCATGTGTCGACGACATCTTCGGTTTGTTCACCGGGTC  720
CACCGGTTCAAGTTACGGAGTGTCGACTAGTGGGTCGCCCGGGTCCATAGGGACAACTAGGCGCCCTCGACTGTCTCCCCGGTTCCTG  V A K F N A S Q L I T Q R A Q V S L L I R R E L T E R A K D  TTCAGCCTCATCCTGGATGATGTGGCCATCACAGAGCTGAGCTTTAGCCGAGAGTACACAGCTGCTGTAGAAGCCAAACAAGTGGCCCAG  AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCGACTCGAAATCGGCTCTCATGTGTCGACGACATCTTCGGTTTGTTCACCGGGTC  720
V A K F N A S Q L I T Q R A Q V S L L I R R E L T E R A K D  TTCAGCCTCATCCTGGATGATGTGGCCATCACAGAGCTGAGCTTTAGCCGAGAGTACACAGCTGCTGTAGAAGCCAAACAAGTGGCCCAG AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCGACTCGACATCGGCTCTCATGTGTCGACGACATCTTCGGTTTGTTCACCGGGTC
AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCGACTCGAAATCGGCTCTCATGTGTCGACGACATCTTCGGTTTGTTCACCGGGTC
AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCGACTCGAAATCGGCTCTCATGTGTCGACGACATCTTCGGTTTGTTCACCGGGTC
CAGGAGGCCCAGCGGGCCCAATTCTTGGTAGAAAAAGCAAAGCAGGGAACAGCGGCAGAAAATTGTGCAGGCCGAGGGTGAGGCCGAGGCT
GTCCTCCGGGTCGCCCGGGTTAAGAACCATCTTTTTCGTTTCGTCCTTGTCGCCGTCTTTTAACACGTCCGGCTCCCACTCCGGCTCCGA
QEAGRAGFLYEKAKQEGRUKLYUAEGEAEA
0.000.0
GCCAAGATGCTTGGAGAAGCACTGAGCAAGAACCCTGGCTACATCAAACTTCGCAAGATTCGAGCAGCCCAGAATATCTCCAAGACGATC GCCAAGATGCTTGGAGAACCTCTTCGAGCACGATCTTGGAACCGATCTAAAACTTCGCAAGATTCTGAAGCTCTTAAGAGGTTCTGCTAG

HMW F546

Page 2

GCCACATCACAGAATCGTATCTATCTCACAGCTGACAACCTTGTGCTGAACCTACAGGATGAAAGTTTCACCAGGGGAAGTGACAGCCTC
GGTGTAGTGTCTTAGCATAGATAGAGTGTCGACTGTTGGAACACGACTTGGATGTCCTACTTTCAAAGTGGTCCCCTTCACTGTCGGAG

A T S Q N R I Y L T A D N L V L N L Q D E S F T R G S D S L

ATCAAGGGTAAGAAATGAGCCTAGTCACCAAGAACTCCACCCCCACAAGAAGTGGATCTGCTTCTCCAGTTTTTGA
TAGTTCCCATTCTTTACTCGGATCAGTGGTTCTTGAGGTGGGGGTGTTCTTCACCTAGACGAAGAGTCAAAAACT

(SEQIDNO:16

IKGKK (SODIDNO: 3)

57 hMF. WO6

Page 1

GGC																														00
CCG	GCT	CTA	CTG	TAG	TGA	T.T.C	ACC	GGC	TAG	ACG	TGT	стс	GTC	CGG	TCC	TCG	TTG	GTG	TĠT	ccc	AAG	GAC	GTG	TAC	CTG	ACG	CTC	TAG	TT	90
																								М	O	С	Ε	į	κ.	
GGG	CGC	ccc	TGC	TGC	ATC	GGC	ACC	AAG	GĢC	AGC	TGT	GAG	ATC	ACC	ACC	ÇGG	GAA	TAC	TGT	GAG	TTC	ATG	CAC	GGC	TAT	TTC	CAT	GAG	GĄ	180
ccc	GCC	GGG	ACG	ACC	TAG	cco	TGC	TTC	ccc	TCG	ACA	CTC	TAG	TGC	TGG	GCC	CTT	ATG	ACA	CTC	AAG	TAC	GTG	CCG	ATA	AAG	GTA	CTC	CT	100
G	R	P	С	С	I	G	T	κ	G	s	С	Ε	i	T	T	R	Ε	Y	С	Ε	F	М	н	G	Υ	F	Н	Ε	Ε	
AGC	ACA	ACTO	TGC	TCC	CAG	GTG	AGG	SC G A	GGC	AGG	CCT	GGA	GTA	GTO	GAG	GAG	AGG	ACG	CTG	GGC	ATO	GCA	GCC	TGC	TGG	GGC	CGC	GGC	.TC	
TCG	TGT	r G A C	ACG	AGG	GTC	CAC	TCC	GCT	CCC	TCC	GGA	CCT	CAT	CAC	СТС	CTC	TCC	TGC	GAC	ccc	TAC	CGT	CGG	ACG	ACC	CCG	GCC	ccc	AG	270
A	T	L	С	s	۵	٧	R	R	G	R	Р	G	v	٧	Ε	Ε	R	T	L	G	М	Α	Α	С	W	G	R	G	s	
ACG	+					+					<del></del>					+			-+-		<del></del>	1				+			-+	360
, , ,	3107	100		1017	·CAG		CGC		_		-	_	_			, can	_	4.0					_	_	-	_				
R	Ť	Р	S	н	٧	G	Α	S	D	S	G	С	F	W	G	А	E	н	Н	M	Р	ı	Р	R	C	i	٧	L	D	
CAA	GTO	TGI	TGG	GCT	GCT	GCC	TTC	СТС	AAC	CCT	GAG	GTC	CCA	GAT	CAG	TTT	TAC	AGG	TCT	GGC	TGT	CTC	TTT	TCC	TAC	ATG	TTO	GGT	AA	450
GTT	CAC	CACA	ACC	CGA	CGA	ĊGG	AAG	GAG	TTO	GGA	CTC	CAG	GGT	CTA	GTC	AAA	ATG	TCC	AĠA	CCG	ACA	GAG	AAA	AGG	ATG	TAC	AAC	CCA	TT	
κ	٧	С	W	A	A	Ά	F	L	N	Р	Ε	٧	P	D	Q	F	Y	R	s	G	С	Ļ	F	s	Y	М	L	G	κ	
GAG	TCC	TC/	ATG	ccc	ccg	AAC	:000	ACC	:cçı	GTG	ATG	GAC	ACC	CAC	GCG	ĢAC	ccc	TGG	GGA	AAG	GTT	cci	GGG	CÇA	GGG	TAT	GGI	CGG	тç	540
CTC	CAGO	SAGI	TAC	GGC	GGC	TTG	GGC	TG	GGA	CAC	TAC	CTG	TGG	GTC	CGC	CTG	GGG	ACC	cct	TTC	CAA	GGA	ccc	GGT	ccc	ΑTA	CCA	GCC	:AG	540
R	s	S	M	Ρ	Ρ	N	Р	T	Ρ	٧	М	D	τ	Q	A	0	P	W	G	K	٧	Ρ	G	Ρ	G	Y	G	R	s	
CAA	сто	secq	AAG	ACT	ACT	GCT	rcci	[GAA	AG TO	STCT	GGA	TGA	AGG	cçc	ста	ÇCT	GGT	GTG	TÇC	стс	ccc	CAC	TGT	GGG	TGC	ACT	GCC	сто		630
GTT	GAC	GGG	TTC	TGA	TGA	CGA	\GGA	ACTI	CAC	AGA	ACCT	ACT	TCC	GGC	GAC	GGA	CCA	CAC	AGG	GAG	GGG	GTO	ACA	ccc	ACC	TGA	CGC	GAC		030
N	L	٩	ĸ	T	т	A	Ρ	ε	٧	s	G	. '	(SE	<b>(</b>	IC	7	<b>1</b> 0:	36	2)	)										
TGT	: 1 T C	STGO	GTC	. T T T	TCA	AAT	GAC	CATO	:001	rga <i>a</i>	AGGG	GAC	CTG	GAC	GAA	AGGT	GGT	CCG	GCT	GGC	ACC	CTA	TCG	C _		<u> </u>	9	I!	γC	10);
ACA																									704					

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

3. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an additional sheet
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and 12301 Parklawn Drive Rockville, Maryland 20852 United States of America	l country)
Date of deposit May 16, 1997	Accession Number 209053
C. ADDITIONAL INDICATIONS (leave blank if not app	olicable) This information is continued on an additional sheet
D DESIGNATED STATES FOR WHICH INDICATIO	ONS ARE MADE (if the indications are not for all designated States)
·	
. SEPARATE FURNISHING OF INDICATIONS (leave	
he indications listed below will be submitted to the Internati Jumber of Deposit")	tional Bureau later (specify the general nature of the indications e.g., "Accession of the indications e.g., "Accession of the indications e.g.,"
For receiving Office use only	For International Bureau use only
For receiving Office use only  This sheet was received with the international application of the internation of the i	

2 3 FED 1998

# INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

	WIFO FOT
A. The indications made below relate to the microorganism referred on page13, line	ed to in the description  able 1
B. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an additional sheet
Name of depositary institution	
American Type Culture Collection	
Address of depositary institution (including postal code and countr 12301 Parklawn Drive Rockville, Maryland 20852 United States of America	(ע
Date of deposit  May 16, 1997	Accession Number 209053
C. ADDITIONAL INDICATIONS (leave blank if not applicable	This information is continued on an additional sheet
In respect of those designations in which of the deposited microorganism will be mathe mention of the grant of the European application has been refused of withdrawn the issue of such a sample to an expert managed (Rule 28(4) EPC).	nde available until the publication of patent or until the date on which the or is deemed to be withdrawn, only by
D. DESIGNATED STATES FOR WHICH INDICATIONS AF	RE MADE (if the indications are not for all designated States)
	ъ.
E. SEPARATE FURNISHING OF INDICATIONS (leave blan	sk if not applicable)
The indications listed below will be submitted to the International B Number of Deposit")	Sureau later (specify the general nature of the indications e.g., "Accession
For receiving Office use only	For International Bureau use only
This sheet was received with the international application	This sheet was received by the International Bureau on:
Authorized office verte simple participal Specialist	Authorized officer

Form PCT/RO/134 (July 1992)

# INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism	
on page 13 ,/i/d/_	Table 1
B. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an additional sheet
Name of depositary institution  American Type Culture Collection	
Address of depositary institution (including postal code and 12301 Parklawn Drive Rockville, Maryland 20852 United States of America	country)
Date of deposit  May 16, 1997	Accession Number
	209034
C. ADDITIONAL INDICATIONS (leave blank if not app	olicable) This information is continued on an additional sheet
D. DESIGNATED STATES FOR WHICH INDICATION	NS ARE MADE (if the indications are not for all designated States)
E. SEPARATE FURNISHING OF INDICATIONS (leav	ve blank if not applicable)
The indications listed below will be submitted to the International Number of Deposit")	ional Bureau later (specify the general nature of the indications e.g., "Accession
For receiving Office use only	For International Bureau use only
This sheet was received with the international application	11
Authorized officer Carol Specialist APD-PCT Operations	Authorized officer

# 59/2

# INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

REC'D	2 3	FEB	1998
WIPO	PC	T	

A. The indications made below relate to the microorganism reference on page13	
B. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an additional sheet
Name of depositary institution  American Type Culture Collection	
Address of depositary institution (including postal code and count 12301 Parklawn Drive Rockville, Maryland 20852 United States of America	(ry)
Date of deposit May 16, 1997	Accession Number 209054
C. ADDITIONAL INDICATIONS (leave blank if not applicable	This information is continued on an additional sheet
EUROPE In respect of those designations in whi of the deposited microorganism will be the mention of the grant of the Europea application has been refused or withdrawby the issue of such a sample to an expethe sample (Rule 28(4) EPC).  D. DESIGNATED STATES FOR WHICH INDICATIONS A	n patent or until the date on which wn or is deemed to be withdrawn, only ert nominated by the person requesting
E. SEPARATE FURNISHING OF INDICATIONS (leave blan	nk if not applicable)
	Bureau later (specify the general nature of the indications e.g., "Accession
For receiving Office use only	For International Bureau use only
Authorized officer  Author	This sheet was received by the International Bureau on:  Authorized officer

What is claimed is:

5

15

20

25

35

1. An isolated polynucleotide comprising a nucleotide sequence that has at least 80% identity over its entire length to a nucleotide sequence encoding the receptor polypeptide of SEQ ID NO:Y; or a nucleotide sequence complementary to said isolated polynucleotide.

60

- The polynucleotide of claim 1 wherein said polynucleotide comprises the nucleotide sequence contained in SEQ ID NO:X encoding the receptor polypeptide of
   SEQ ID NO:Y.
  - 3. The polynucleotide of claim 1 wherein said polynucleotide comprises a nucleotide sequence that is at least 80% identical to that of SEQ ID NO:X over its entire length.
  - 4. The polynucleotide of claim 3 which is polynucleotide of SEQ ID NO:  $\mathbf{X}$ .
    - 5. The polynucleotide of claim 1 which is DNA or RNA.
  - 6. A DNA or RNA molecule comprising an expression system, wherein said expression system is capable of producing a receptor polypeptide comprising an amino acid sequence, which has at least 80% identity with the polypeptide of SEQ ID NO:Y when said expression system is present in a compatible host cell.
    - 7. A host cell comprising the expression system of claim 6.
- 8. A process for producing a receptor polypeptide comprising culturing a host of claim 7 under conditions sufficient for the production of said polypeptide and recovering the polypeptide from the culture.
  - 9. A process for producing a cell which produces a receptor polypeptide thereof comprising transforming or transfecting a host cell with the expression system of claim 6 such that the host cell, under appropriate culture conditions, produces a receptor polypeptide.
  - 10. A receptor polypeptide comprising an amino acid sequence which is at least 80% identical to the amino acid sequence of SEQ ID NO:Y over its entire length.

- 11. The polypeptide of claim 10 which comprises the amino acid sequence of SEQ ID NO:Y.
- 5 12. An antibody immunospecific for the receptor polypeptide of claim 10.
  - 13. A method for the treatment of a subject in need of enhanced activity or expression of receptor polypeptide of claim 10 comprising:
  - (a) administering to the subject a therapeutically effective amount of an agonist to said receptor; and/or
  - (b) providing to the subject polynucleotide of claim 1 in a form so as to effect production of said receptor activity in vivo.
- 14. A method for the treatment of a subject having need to inhibit activity or expression of the receptor polypeptide of claim 10 comprising:
  - (a) administering to the subject a therapeutically effective amount of an antagonist to said receptor; and/or
  - (b) administering to the subject a nucleic acid molecule that inhibits the expression of the nucleotide sequence encoding said receptor; and/or
  - (c) administering to the subject a therapeutically effective amount of a polypeptide that competes with said receptor for its ligand.
    - 15. A process for diagnosing a disease or a susceptibility to a disease in a subject related to expression or activity of the receptor polypeptide of claim 10 in a subject comprising:
    - (a) determining the presence or absence of a mutation in the nucleotide sequence encoding said receptor polypeptide in the genome of said subject; and/or
- (b) analyzing for the presence or amount of the receptor polypeptide30 expression in a sample derived from said subject.
  - 16. A method for identifying agonists to the receptor polypeptide of claim 10 comprising:
    - (a) contacting cells produced by claim 9 with a candidate compound;
- 35 and

20

25


(b) determining whether the candidate compound effects a signal generated by activation of the receptor polypeptide.

- 17. An agonist identified by the method of claim 16.
- 18. The method for identifying antagonists to the receptor polypeptide of claim 10 comprising:
  - (a) contacting said cell produced by claim 9 with an agonist; and
- (b) determining whether the signal generated by said agonist is diminished in the presence of a candidate compound.
  - 19. An antagonist identified by the method of claim 18.


15

5

- 20. An isolated receptor polynucleotide comprising a nucleotide sequence selected from the group consisting of:
- (a) a nucleotide sequence having at least 80% identity to a nucleotide sequence encoding the receptor polypeptide expressed by the cDNA insert deposited at the ATCC; and
- (b) a nucleotide sequence complementary to the nucleotide sequence of (a).
- 21. A recombinant host cell produced by a method of Claim 9 or a membrane thereof expressing a receptor polypeptide.



F16.1A



F 6.1





FIG. 10



-16.2A





FIG.28



SUBSTITUTE SHEET (RULE 26)




FIG. 4A







FIG.4B




F16.5A





HIG.6A



		1.77/#/		
HTPBA27.aa NAG-2.aa	HTPBA27.aa NAG-2.aa	HIPBA27.aa NAG-2.aa	HTPBA27.aa NAG-2.aa	HTPBA27.aa NAG-2.aa
1 40 WLAATQGS WLAATQGS	80   80   1 G A I K E N K   L G A I K E N K	120 RYAQQDLK RYAQQDLK	160 Y T D W F E V Y Y T D W F E V Y	ZOC TVKVWLQE TVKVWLQE
20 30 FWLGGCGVLGVGIWLAATQGS FWLGGCGVLGVGIWLAATQGS	70 MAIGFVGC MAIGFVGC	110 120 FAYTDKIDRYAQQDLK FAYTDKIDRYAQQDLK	150 160 QTDFRCCGVSNYTDWFEVY QTDFRCCGVSNYTDWFEVY	180 190 200 LHAPGTWWKAPCYETVKVWLOE LHAPGTWWKAPCYETVKVWLOE
FNLFFWLGGCGVLGVGIWLAATQGS	LLILTGAFVMAIGFVGCLGAIKENK	100 LEATIATLF LEATIAILF	140 NAWSIIQTDFRCCGVSNYTDWFEVY NAWSIIQTDFRCCGVSNYTDWFEVY	SCGLHAPGTWWKAPCYETVKVWLQE
10 QAVKYLMFAI QAVKYLMFAI	SFPSLSAAN!	90 FLLLLLLVFL1 FLLLLLVFL1	130 TOGNVGLT TOGNVGLT	170 CCLEFSE CCLEFSE
MARACL (	41 FATLSS 41 FATLSS	SI CLLLTF1 81 CLLLTF1	121 KGLHLYG 121 KGLHLYG	161 NATRVPDS 161 NATRVPDS

FIG.7




FIG.84






FIG. 8B



F16.9A



F.0




FIG. 10





FIG. 11A







FIG. 12A



SUBSTITUTE SHEET (RULE 26)

PLR GNIVKKDCAESCTPSYTLQGOV
D L C N E


FIG. 13

30 L. L. G. A. G. HMMSS46.aa	LLGAG B-cell Recpt Associated Pr.aa	IFFNRIGGVOO HWGS46.aa IFFNRIGGVOO B-cell Recpt Associated Pr.aa	RPRKI B-cell Recpt Associated Pr.aa	120 LELPSM HMWGS46.aa ELPSM B-cell Recpt Associated Pr.aa	150 1 HWGS46.aa
1 10 1 D L A G R L	DLAGRL - 40	R E S V F T V E G G H R A R E S V F T V E G G H R A	70 80 ILAEGLHFRIPWFQYPIIYDIRAF ILAEGLHFRIPWFQYPIIYDIRAF	100 K D L Q M V N I S L R V L S R P N A Q K D L Q M V N I S L R V L S R P N A Q	RLGLDYEERVLPSTVNEVLKSVVAKFNA HMGS46.aa
MAQNLK	M	31 AVAYGV 31 AVAYGV	a DTILAE a DTILAE	91 SSPTGS 91 SSPTGS	121 Y Q R L G L

FIG. 14A

HMMSS46.aa B-cell Recpt Associated Pr.aa	HMMSS46.aa B-cell Recpt Associated Pr.aa	HMKS46.aa B-cell Recpt Associated Pr.aa	HMMCS46.aa B-cell Recpt Associated Pr.aa	HMGS46.aa B-cell Recpt Associated Pr.aa
160 170 180 151 SQLITQRAQVSLLIRRELTERAKDFSLILD 151 SQLITTORAQVSLLIRRELTERAKDFSLILD	190 200 210 31 DVAITELSFSREYTAAVEAKQVAQQEAQRA HMMGS46.aa 81 DVAITELSFSREYTAAVEAKQVAQQEAQRA B-cell Req	220 230 240  1 QFLVEKAKQEQRQKIVQAEGEAEARKMLGE HMMCS46.aa  1 OFLVEKAKQEQROKIVQAEGGEAEARKMLGE B-cell Recpt Associated Pr.aa	250 260 270  ALSKNPGYIKLRKIRAAQNISKTIATSQNR HAMGS46.aa  ALSKNPGYIKLRKIRAAQNISKTIATSQNR B-cell Rec	280 290 300 I Y L T A D N L V L N L Q D E S F T R G S D S L I K G K K . I Y L T A D N L V L N L Q D E S F T R G S D S L I K G K K
15	181	# # # # # # # # # # # # # # # # # # #	241 241	271 271 271

FIG. 14B



#### INTERNATIONAL SEARCH REPORT

HONA	LOE	AHUH	REP

Internatio	nt	Application No
P	5	98/00959

				Application No 98/00959
A. CLASS IPC 6	C12N15/12 C12N15/85 C12N5/ C12Q1/68 A61K38/17 A61K48			D7K16/28
According t	o international Patent Classification (IPC) or to both national classifi	ication and IPC		
	SEARCHED			
Minimum de IPC 6	CL2N CO7K C12Q A61K	ation symbols)		
Documenta	tion searched other than minimum documentation to the extent that	such documents are	e included in the fields	searched
	ata base consulted during the international search (name of data b	ase and, where pra	ctical, search terms us	sed)
Category *	ENTS CONSIDERED TO BE RELEVANT  Citation of document, with indication, where appropriate, of the re			Colores An alaim Ala
Calcyc.,	ORALISH OF GOODINGHICK WITH INGIGENCY WITH THE CAPPAGE MAN TO THE CONTROL OF THE	18vani hassadas		Relevant to claim No.
Х	DATABASE GENBANK Accession No. T48852, 8 February "EST70319 from Stratagene place XP002066898 see the whole document			1-12,20
X	DATABASE GENBANK Accession No. G14442, 4 January "STS SHGC-10055" XP000206689 see the whole document	1996		1-3,20
		-/		
		·		
X Furth	er documents are listed in the continuation of box C.	Patent fa	mily members are lists	ed in annex.
"A" docume consider of filing de "L" docume which i citation "O" docume other m" "P" docume later th.	nt which may throw doubts on priority claim(s) or socied to establish the publication date of another or or other special reason (as specified) or referring to an oral disclosure, use, exhibition or neans at published prior to the international filing date but an the priority date claimed	or priority data cited to under cited to under cited to under cited to under cannot be control to each courrent is ments, such in the art.	retand the principle or articular relevance; the naidered novel or can ventive step when the articular relevance; the naidered to involve an combined with one or combination being observed.	ith the application but theory underlying the e claimed invention not be considered to document is taken alone e claimed invention inventive step when the more other such docurrious to a person skilled ant family
	ctual completion of the international search  June 1998	Date of mailin	g of the international s 1 6. 09. 19	·

Form PCT/ISA/210 (second sheet) (July 1992)

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentiaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo ni, Fax: (+31-70) 340-3016

Authorized officer

Galli, I

# INTERNATIONAL SEARCH REPORT

International Application No CT/US 98/00959

C1/0S 98/00959		·
	nuation) DOCUMENTS CONSIDERED TO BE RELEVANT	<u> </u>
Relevant to claim No.	Citation of document, with indication, where appropriate, of the relevant passages	Category '
1-21	ZANNETTINO A.C.W. ET AL.: "A powerful technique for isolating genes encoding surface antigens using retroviral expression cloning" J. IMMUNOL., vol. 156, no. 2, 15 January 1996, pages 611-620, XP002066893 see abstract	A .
1-21	ADAMS M D ET AL: "COMPEMENTARY DNA SEQUENCING: EXPRESSED SEQUENCE TAGS AND HUMAN GENOME PROJECT" SCIENCE, vol. 252, no. 5013, 21 June 1991, pages 1651-1656, XP000645049 cited in the application see abstract	A
1-21	SIMMONS D. & SEED B.: "Isolation of a cDNA encoding CD33, a differentiation antigen of myeloid progenitor cells" J. IMMUNOLOGY, vol. 141, no. 8, 15 October 1988, pages 2797-2800, XP002066894 see abstract see figure 4	Α
1-21	CROCKER P.R. ET AL.: "Sialoadhesin, a macrophage sialic acid binding receptor for haemopoietic cells with 17 immunoglobulin-like domains" EMBO J., vol. 13, no. 19, 3 October 1994, pages 4490-4503, XP002066895 see abstract see figure 2	A .
1-21	KELM S. ET AL.: "The Sialoadhesins: a family of sialic acid-dependent cellular recognition molecules within the immunoglobulin superfamily" GLYCOCONJ. J., vol. 13, no. 6, December 1996, pages 913-926, XP002066896 see the whole document	A .
1-21	DATABASE GENBANK Accession No. U71382, 12 November 1996 PATEL N. ET AL.: "OB binding protein - 1" XP002066901 cited in the application see the whole document	Α
	Accession No. U71382, 12 November 1996 PATEL N. ET AL.: "OB binding protein - 1" XP002066901 cited in the application see the whole document	Α

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

# INTERNATIONAL SEARCH REPORT

PC 98/00959

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT			
Category <sup>5</sup>	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.	
P,X	DATABASE GENBANK Accession No. AA344713, 21 April 1997 "EST50650, H. sapiens cDNA similar to CD33 antigen" XP002066900 see the whole document	1-12,20	
P,A	TAKEI Y. ET AL.: "Molecular cloning of a novel gene similar to myeloid antigen CD33 and its specific expression in placenta" CYTOGENET. CELL GENET., vol. 78, 1997, pages 295-300, XP002066897 see abstract see figure 1	1-21	



ter onal application No.
PCT/US` 98/ 00959

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)
This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:
1. X Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely: Remark: Although claims 13 and 14 are directed to a method of treatment of the human/animal body, and claim 15 (as far as in vivo methods are envisaged) is directed to a diagnostic method, the search has been carried out and based on the alleged effects of the compound/composition.  2. Claims Nos.: because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).
Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)
This International Searching Authority found multiple inventions in this international application, as follows: see further information sheet
As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. X No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:  see further information sheet, subject 1.
Remark on Protest  The additional search fees were accompanied by the applicant's protest.  No protest accompanied the payment of additional search fees.

Form PCT/ISA/210 (continuation of first sheet (1)) (July 1992)

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: (1-21) - partial

An isolated polynucleotide encoding the HMACR70 cell surface receptor (Seq. IDs 1,18), polynucleotides complementary thereto, fragments thereof, homologs thereof.

An expression system comprising said polynucleotides, a process to produce a cell harbouring said expression system. Said cell or membrane thereof expressing said receptor.

A process for the production of the HMACR70 polypeptide comprising culturing said cell.

Said HMACR70 polypeptide or homologs. An antibody directed against said polypeptide.

Therapeutic and diagnostic applications.

A method to identify agonists or antagonists to said receptors. Said agonists and antagonists.

2. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 2,3,19

3. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 4,20

4. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 5,21

5. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 6,22

6. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 7,23

7. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 8,24

### FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

- 8. Claims: (1-21) partial

  Idem as subject matter 1 but limited to Seq. IDs 9,10,25
- 9. Claims: (1-21) partial

  Idem as subject matter 1 but limited to Seq. IDs 11,26
- 10. Claims: (1-21) partial
   Idem as subject matter 1 but limited to Seq. IDs 12,27
- 11. Claims: (1-21) partial
   Idem as subject matter 1 but limited to Seq. IDs 13,28
- 12. Claims: (1-21) partial
   Idem as subject matter 1 but limited to Seq. IDs 14,29
- 14. Claims: (1-21) partial
   Idem as subject matter 1 but limited to Seq. IDs 16,31
- 15. Claims: (1-21) partial

  Idem as subject matter 1 but limited to Seq. IDs 17,21

THIS PAGE BLANK (USPTO)